

Original Paper

Availability and Accessibility of Student-Specific Weight Loss Programs and Other Risk Prevention Health Services on College Campuses

Sarah Lynch¹, MPH; Sharon Hayes², PhD; Melissa Napolitano³, PhD; Katrina Hufnagel⁴, BS

¹Department of Psychology, University of Colorado Denver, Denver, CO, United States

²Keiser University, Department of Psychology, Port St. Lucie, FL, United States

³The George Washington University, Department of Prevention and Community Health; Department of Exercise Science, Milken Institute of Public Health, Washington, DC, United States

⁴The George Washington University, Department of Prevention and Community Health, Milken Institute of Public Health, Washington, DC, United States

Corresponding Author:

Sarah Lynch, MPH

Department of Psychology

University of Colorado Denver

1200 Larimer Street, 5th Floor

Denver, CO, 80204

United States

Phone: 1 702 845 3238

Fax: 1 303 556 3520

Email: sarah.lynch@ucdenver.edu

Abstract

Background: More than one third of college students who are overweight or obese are in need of weight loss programs tailored to college students. However, the availability and accessibility of these programs is unknown.

Objective: The aim of this study is to examine the availability and ease of access to weight loss programs for students at 10 universities with the largest undergraduate enrollment.

Methods: The 10 public universities with the largest student bodies with a mean (SD) undergraduate enrollment of 41,122 (7657) students were examined. The websites of the universities were assessed to determine the availability of weight loss programs. Services for high-risk health needs common to university campuses (ie, alcohol and other drugs, victim services, sexual health, and eating disorders) were searched.

Results: Of the universities searched, 3 (30%, 3/10) offered weight loss programming, however, none met the predetermined criteria. Comparatively, all schools (100%, 10/10) offered no-cost and continual enrollment programming for the other high-risk health needs.

Conclusions: There are limited weight loss services available to undergraduate students compared with other university services. Collaboration between existing college health service providers is suggested for the delivery of appropriate programming for overweight and obese undergraduates wanting to lose weight.

(*JMIR Public Health Surveill* 2016;2(1):e29) doi: [10.2196/publichealth.5166](https://doi.org/10.2196/publichealth.5166)

KEYWORDS

weight loss; student health services; students; universities

Introduction

College campuses provide a foundation for the development of lifelong behaviors for over 70% of young adults between the ages of 18 and 24 who enroll in college [1]. During the

undergraduate years, young adults must adjust to multiple personal, social, and academic demands while trying to establish or maintain healthy lifestyle behaviors [2]. This adjustment can be challenging particularly when it comes to establishing healthy eating and exercise patterns [2]. Services offered on university campuses are a convenient resource for undergraduate students

looking for assistance with the college adjustment. Undergraduate students' need for university-based health services has increased over the past 5 years, as evidenced in a 2013 survey of college counselors [3].

Services for high health risk needs, like alcohol and other substance use, eating disorders, sexual health services, and victim services are generally recognized as critical to ensure the health of undergraduate students and other university members. However, the prevalence of the high health risk needs requiring services on college campuses varies dramatically. For example, the prevalence of undergraduate students with alcohol use disorders ranges between 20.4% [4] to 31% [5]. According to the Centers for Disease Control and Prevention, 1 in 4 sexually active young adults between the ages of 19 to 24 years have a sexually transmitted disease [6]. Rates of sexual assault in undergraduates range from 1.9% to 7% in college populations [7]. With regards to eating disorders, 1.2% of undergraduate females report being diagnosed and/or treated for anorexia [7], 1% for bulimia [7], and 8.4% for binge eating disorder [8]. An additional 9% of young women meet Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria for specified or unspecified feeding or eating disorder [9,10]. Comparatively, approximately 35% percent of college students are overweight or obese [7].

As with the services for high-risk needs described above, colleges have the unique opportunity to address weight and weight-related topics to further ensure the health of their student body. Average weight gain from college matriculation to graduation ranges from 3.5 pounds [11] to 9.5 pounds [12]. Of note, upon college matriculation, 22% of college students are already overweight, and 12% are obese [7]. With one-third of undergraduate students already at risk for negative health and psychosocial outcomes associated with obesity [13,14], relatively small weight gains can increase an individual's risk of having a body mass index (BMI) in the overweight or obese category [15]. These small gains may increase an individual's risk for type 2 diabetes, cardiovascular disease, and depression [16-19].

Despite this public health concern, there remains a need on campuses for weight management services tailored to undergraduates [20,21]. Behavioral health services for the treatment of obesity have lagged behind other on-campus high-risk health behavior services, including drug and alcohol use, risky sexual behavior, eating disorders, and victim services [22]. A 2013 survey of university-based registered dietitians found that over half of participating university dietitians felt it was important for their respective universities to offer structured weight loss programming for students [23]. An integrative approach to the treatment of overweight and obesity could provide an opportunity for counseling centers and health centers to collaborate by promoting the application of evidenced-based practices and access to mental health resources that individuals may need while losing weight. However, it is unclear what services are currently available for overweight and obese undergraduate students wanting to lose weight. No systematic assessment of weight loss programs specific to college students and offered on university campuses could be located in the literature.

Research suggests that weight gain [24] and access to weight-related services [25] are concerns for undergraduate students. A recent study reviewing causes of stress referenced in undergraduate student Facebook pages found weight gain was referenced in 10% of posts [24]. A qualitative study of undergraduate students found that students who felt they had greater access to fitness facilities and healthy eating options were more likely to use those services [25]. Despite this evidence, it remains unclear as to what weight loss services are available and accessible to overweight and obese undergraduate students wanting to lose weight.

This study examines the availability and ease of access of weight loss programs for undergraduate students compared to existing services for other high-risks health needs (ie, alcohol and other drugs, eating disorders, victim services, and sexual health).

Methods

Search Methodology

The 10 public universities with the largest undergraduate student bodies, with a mean (SD) undergraduate enrollment of 41,122 (7657.4) students, were identified using the 2013 US News and World Report College Guide [26]. Because undergraduate students may be more comfortable in programming specifically for students of their same age and demographic, particularly ones that do not include graduate students or faculty who may be their professors or advisors, we selected to focus on weight loss programs tailored specifically to undergraduate students. Similarly, health programming for other high-risk health services are traditionally tailored exclusively to the needs of undergraduate students; as such, we decided to use this same metric when examining weight loss programming. The Internet is ranked as the top source of gathering health-related information for college students [27]. Thus, a Web-based search was elected to search for university weight loss programs. Additionally, college students report trusting health information originating from their respective universities and university health centers [7,27]. For this reason, only websites with edu links were reviewed.

Structured searches of university pages using standardized search terms were conducted over a period of 4 months (July 2013 to October 2013). Search terms were typed into the university's main page by 3 independent coders (2 undergraduate and 1 graduate research assistants). Phone calls were placed to university health, counseling, and wellness centers to confirm the Web search findings.

Search terms were initially selected through a brief survey of undergraduate students that asked students to identify what term they would use to search for weight loss services on their campus. The suggested search terms were reviewed by 2 clinical psychologists and a registered dietitian with experience in the development and delivery of undergraduate weight loss programming, as well as by 2 undergraduate and 2 graduate research assistants. A total of 15 key terms were used to search for student-specific weight loss programs using the search box found on the home page of the respective universities: diet program, weight loss program, losing weight, fitness program,

nutrition counseling, nutritionist, dietician, dietitian, exercise facility, weight watchers, student weight loss, student diet program, student weight loss program, weight management, and student weight management .

Search terms for traditional high-risk health services offered on university campuses (ie, alcohol and other drugs, victim services, sexual health, eating disorders) were included to compare the availability of these services to student weight loss services. These terms were also typed in the search box found on the home page of the respective universities.

Search terms were additionally typed into the search box on the student health services page and university counseling pages to ensure no programs were not found due to website-derived search algorithms unknown to the research team.

Establishment of Inter-Rater Reliability

The aforementioned clinical psychologists established criteria prior to the webpage reviews. In order to be coded as a weight loss program, the program must have been undergraduate student-specific, offered on campus, no additional cost, and delivered by university-funded treatment providers. Results were also coded as “related weight loss services” (missing one or more of the weight loss program criterion, fee-based weight loss program, nutrition counseling, personal training, research on college student weight loss or community-based weight loss program), or no weight loss program (group fitness programs, academic research with a focus outside the scope of college student weight loss, nutrition and physical activity handouts).

Two undergraduate and one graduate research assistant piloted the search terms on 4 large (undergraduate student enrollment greater than 30,000 students) test universities, which were randomly selected from a list of universities from the US News and World Report [26]. These 4 schools were not included in the final university sample pool. Terms were typed into the university search engine found on the university’s home page. Research assistants took a screen shot of the results and a screen shot of the first page of each link clicked on, in order to capture the content of the website at the time reviewed. After completion of coding, screen shots and coding of the content were reviewed and discussed between the coders. Coding decisions were confirmed under the direction of 2 clinical psychologists who have experience in the development and delivery of college weight loss treatments.

Results

Weight Loss Programs

Although the predetermined criteria were not met, the following 3 universities offered some weight loss programming: (1) University of Minnesota offered a weight loss program for their undergraduate students; however, the 12-week program for undergraduates cost US \$250, (2) Florida International University offered an 8-week program for undergraduate students inspired by the Biggest Loser for US \$80, and (3) University of Texas-Austin offered a 12-week program exclusively to graduate students for US \$60. Phone calls placed to university health, counseling, and wellness centers verified the results. The aforementioned programs had discrete

enrollment periods. Two other universities offered short-term programs, but there was no confirmed plan to continue those regularly beyond the planned dates of the one program. For the purpose of this paper those programs were not included.

Accessibility of Weight Loss Programs

The results of our findings are shown in Table 1. Coders spent a mean (SD) of 102.2 (21.0) minutes searching for weight loss programs using the 15 search terms, and a mean (SD) of 8.2 (6.8) minutes per search term. Comparatively, coders spent a mean (SD) of 20.5 (2.9) minutes searching for 4 traditional university services and 5.1 (0.7) minutes per search term. Although the predetermined criteria were not met, 3 universities offered student weight loss programs. Comparatively, all of the universities searched (100%, 10/10) offered free, university-sponsored, student-specific services for sexual health and victim services. We found that 90% (9/10) of the universities offered free, university-sponsored student-specific services for alcohol and other drug treatments; one school required a fee for assessment and treatment. In addition, 90% (9/10) universities offered free, university-sponsored student-specific services for eating disorders; one school provided screening and referred out to the community for treatment.

Discussion

Principal Findings

Three schools offered weight loss programming, however, none met the predetermined criteria. Comparatively, all schools offered no-cost and continual enrollment programming for the other high-risk health needs. Given the current prevalence of overweight and obesity among college students and the demonstrated weight gain during college, there is a need for obesity-related services on campuses. Universities have competing wellness priorities for their students, and they may designate resources for those behavior and health risks that have more immediate negative consequences such as suicide or alcohol-related death or injury [28-30]. Obesity treatment may be a lower priority as the consequences may not manifest until long after students leave the university. However, our findings indicate that some colleges are beginning to offer services specific to the treatment of obesity in their college populations.

The focus of weight management on college campuses may reflect changes in health services mandated by the Affordable Care Act (ACA). Students are now able to remain on their parents’ insurance until the age of 26 years old [31]. The extended insurance coverage is coupled with an increased focus on obesity prevention and weight loss under the ACA, which requires insurance providers to provide regular BMI screening, nutrition counseling, and other weight management services for no or little cost to the insured [32]. The 35% prevalence rate of overweight and obesity in undergraduate students [7] is comparable to or exceeds the prevalence rates of high-risk behaviors that have regularly provided no-cost services (10% for eating disorder [7-10], 20-30% for alcohol and other drug use [4,5], 25% of sexually transmitted disease [6], and 1-7% for sexual assault [7]). Given the shift in focus that we have seen nationally related to weight management due to the ACA, colleges have the opportunity to mirror these new initiatives.

Table 1. The largest public universities and the health services offered (N=10).

University	Number of students ^a	Weight loss program	Victim services	Sexual health	Alcohol and other drugs	Eating disorder treatment
University of Central Florida	51,269	0 ^b	X ^c	X	X	R ^d (only screening offered)
Texas A&M	44,315	0	X	X	X	X
Ohio State	44,201	0	X	X	X	X
Penn State	40,085	0	X	X	R (fee)	X
University of Texas at Austin	39,979	R (fee; only for graduate students)	X	X	X	X
Florida International University	39,045	R (fee)	X	X	X	X
Arizona State University	38,735	0	X	X	X	X
Michigan State	37,988	0	X	X	X	X
University of Minnesota	34,449	R (fee)	X	X	X	X
University of Florida	33,168	0	X	X	X	X

^aAcademic year 2013-2014.

^b0 is no program.

^cX is a program that meets established criteria.

^dR is a related program that does not meet established criteria for program.

Despite these opportunities, colleges and universities are faced with a number of demands in terms of academic and health-related priorities and limited funds. Examining the barriers and facilitators on college campuses to delivering weight loss programming is an important step in understanding the broad scale implementation and acceptability of these programs. One on-going project, funded by the National Institutes of Health (DK 100916), is examining the efficacy of using social media and text messaging for delivering weight loss programming to college students on 2 different campuses [33]. In addition to examining weight loss and metabolic outcomes (ie, glucose, blood pressure, lipids), one primary aim of this project is to evaluate the implementation feasibility and sustainability infrastructure on college campuses for delivering weight loss programs tailored for students. This evaluation will include a cost effectiveness assessment as well as interviews with key administrative and student leaders on campus to provide data on both the facilitators and challenges to offering tailored weight management programs.

A 2011 analysis of the American College Health Association-National College Health Assessment (ACHA-NCHA) database found that the top two most trusted sources of health information were health center medical staff and health educators [34]. However, this same study found that these two sources of information were also the most under-utilized method of gathering health information [34]. While it is unclear why these on-campus resources are underutilized, it is possible that the nature of the Web search (eg, timely, cumbersome) may have been a barrier to utilization. Students relying on the Internet alone might struggle to find information about on-campus services. The cost of these programs may also reduce the likelihood that students will utilize the resources. We found weight loss programs exclusively for

undergraduate students were available for a fee (US \$60-\$250) at 2 of the 10 universities included in this Web-based search (University of Minnesota and Florida International University).

For these and other reasons, a multidisciplinary approach to disseminating and marketing a weight loss program specific to college students is critical. Collaboration with the counseling center may be particularly important. The is concern that weight loss services on college campuses may promote disordered eating and/or eating disorders in a vulnerable college population [35]. Dieting is prevalent on college campuses: over 60% of college females and 36% of college males report they are trying to lose weight, and 48% of college females and 28% of college males reported dieting to lose weight [7]. Overweight and obese young adults who desire to lose weight are just as likely to develop an eating disorder as the general population [36], yet eating disorder treatment for this population is often over looked [23,37].

There is literature suggesting that there may be a common underpinning for both excessive weight gain and eating disorders. These shared experiences and characteristics are dieting, media use, body image dissatisfaction, and weight-related teasing [38,39]. Thus, it is not weight loss per se, but a combination of maladaptive behaviors and factors that may put an overweight or obese individual at risk for an eating disorder. College women, in particular, may use a combination of healthy and unhealthy weight loss practices [40]. Thus, a multimodal approach of social, environmental, and individual strategies coupled with media literacy and advocacy is needed for overweight and obese students wanting to lose weight [23,41]. It is critical that university health providers delivering weight loss programs for students be aware of the possibility of underlying eating disorders that may complicate the students' efforts to lose weight so that the appropriate treatment can be

delivered to students [23,37]. The key is to provide education and evidence-based programs to teach safe, effective, and healthy ways of losing weight.

While many university health officials and administrators recognize the need for an environment that promotes healthy weight loss practices and relevant programming, there are a number of challenges faced by these key leaders on campus. There are competing priorities for resources, time, as well as other behavioral and health needs with more immediate consequences [28,29,42]. Unfortunately, outsourcing weight loss programming is likely not the solution to implementing weight loss programs for undergraduate students as traditional weight loss programs for adults. Young adults who attend weight loss programs tailored to adult populations attend fewer sessions and lose less weight than adults attending weight loss programs [43]. Many universities likely already have necessary components that would allow them to offer a systematic program to prevent and treat obesity. The key is to integrate these available services and personnel to provide education and evidence-based programs that are attuned to the potential for disordered eating behaviors in this population of overweight and obese students wanting to lose weight [40]. Programming that promotes a campus environment of positive body image and self-esteem may help overweight and obese and normal weight students who want to lose weight. For example, The Body Positive is a body image and health curriculum for high school students that promote physical and mental health through

workshops, videos, and campus events. These types of programs could provide an interim solution prior to the launching of sustained weight loss program offerings.

Limitations

This study provides only a snapshot of the availability and ease of access to weight loss services on college campuses. Only the websites of the ten largest public universities were reviewed, limiting the search and possibly ignoring student weight loss programs available at smaller and/or private universities. Universities often update their webpages so that recent content is pushed to the top of searches, so it is possible that the results presented here were influenced by search mechanisms unknown to the study authors. Lastly and importantly, this study was not able to determine the extent of the utilization of available weight loss services by university students.

Conclusions

This study examined the availability and ease of access to weight loss programs for college students. The results highlight that there are limited weight loss services available and accessible to college students when compared to other university services, such as alcohol and other drugs, eating disorders, sexual health, and victim services.

Advocacy from student groups and collegiate administrators is called for to provide treatment and reduce stigma regarding accessing resources specific to weight loss for overweight and obese students wanting to lose weight.

Acknowledgments

We would like to acknowledge Stephanie Goldstien and Jana Howard, who helped with the conceptualization of this project. Additionally, we would like to thank Hillary DeCarl for her assistance in coding the webpages.

This project was partially funded by an internal grant from The George Washington University Food for Thought Initiative.

Conflicts of Interest

None declared.

References

1. National Center for Education Statistics. Digest of Education Statistics. Washington, DC: U.S. Department of Education; 2014. Percentage of recent high school completers enrolled in 2-year/4-year colleges, by income level: 1975 through 2012 URL: https://nces.ed.gov/programs/digest/d13/tables/dt13_302.30.asp [WebCite Cache ID 6hkvs2ZYA]
2. Cluskey M, Grobe D. College weight gain and behavior transitions: male and female differences. *J Am Diet Assoc* 2009 Feb;109(2):325-329. [doi: [10.1016/j.jada.2008.10.045](https://doi.org/10.1016/j.jada.2008.10.045)] [Medline: [19167962](https://pubmed.ncbi.nlm.nih.gov/19167962/)]
3. Gallagher R. National Survey of College Counseling Centers 2013. Alexandria, VA: The International Association of Counseling Services, Inc; 2013. URL: <http://www.collegecounseling.org/wp-content/uploads/Survey-2013-4-yr-Directors-1.pdf> [WebCite Cache ID 6hkw1FVww]
4. Blanco C, Okuda M, Wright C, Hasin D, Grant B, Liu S, et al. Mental health of college students and their non-college-attending peers: results from the National Epidemiologic Study on Alcohol and Related Conditions. *Arch Gen Psychiatry* 2008 Dec;65(12):1429-1437 [FREE Full text] [doi: [10.1001/archpsyc.65.12.1429](https://doi.org/10.1001/archpsyc.65.12.1429)] [Medline: [19047530](https://pubmed.ncbi.nlm.nih.gov/19047530/)]
5. Knight JR, Wechsler H, Kuo M, Seibring M, Weitzman ER, Schuckit MA. Alcohol abuse and dependence among U.S. college students. *J Stud Alcohol* 2002 May;63(3):263-270. [Medline: [12086126](https://pubmed.ncbi.nlm.nih.gov/12086126/)]
6. Centers for Disease Control/Prevention. Surveillance and Information Systems Branch. 2014 Dec 16. Division of STD/HIV Prevention URL: <http://www.cdc.gov/std/stats13/> [accessed 2016-05-18] [WebCite Cache ID 6hbGIm1Zz]
7. American College Health Association. National College Health Assessment II: Spring 2014 Reference Group Executive Summary. Hanover, MD; 2014 May 01. National College Health Assessment II: Spring 2014 Reference Group Executive Summary URL: http://www.acha-ncha.org/docs/ACHA-NCHA-II_ReferenceGroup_ExecutiveSummary_Spring2014.pdf [WebCite Cache ID 6hkwMGZZO]

8. Napolitano M, Himes S. Race, weight, and correlates of binge eating in female college students. *Eat Behav* 2011 Jan;12(1):29-36. [doi: [10.1016/j.eatbeh.2010.09.003](https://doi.org/10.1016/j.eatbeh.2010.09.003)] [Medline: [21184970](https://pubmed.ncbi.nlm.nih.gov/21184970/)]
9. Smink FR, van Hoeken D, Oldehinkel A, Hoek H. Prevalence and severity of DSM-5 eating disorders in a community cohort of adolescents. *Int J Eat Disord* 2014 Sep;47(6):610-619. [doi: [10.1002/eat.22316](https://doi.org/10.1002/eat.22316)] [Medline: [24903034](https://pubmed.ncbi.nlm.nih.gov/24903034/)]
10. Stice E, Marti C, Rohde P. Prevalence, incidence, impairment, and course of the proposed DSM-5 eating disorder diagnoses in an 8-year prospective community study of young women. *J Abnorm Psychol* 2013 May;122(2):445-457 [FREE Full text] [doi: [10.1037/a0030679](https://doi.org/10.1037/a0030679)] [Medline: [23148784](https://pubmed.ncbi.nlm.nih.gov/23148784/)]
11. Holm-Denoma J, Joiner T, Vohs K, Heatherton T. The “freshman fifteen” (the “freshman five” actually): predictors and possible explanations. *Health Psychol* 2008 Jan;27(1 Suppl):S3-S9. [doi: [10.1037/0278-6133.27.1.S3](https://doi.org/10.1037/0278-6133.27.1.S3)] [Medline: [18248103](https://pubmed.ncbi.nlm.nih.gov/18248103/)]
12. Racette S, Deusinger S, Strube M, Highstein G, Deusinger R. Changes in weight and health behaviors from freshman through senior year of college. *J Nutr Educ Behav* 2008;40(1):39-42. [doi: [10.1016/j.jneb.2007.01.001](https://doi.org/10.1016/j.jneb.2007.01.001)] [Medline: [18174103](https://pubmed.ncbi.nlm.nih.gov/18174103/)]
13. Huang TT, Shimel A, Lee RE, Delancey W, Strother ML. Metabolic Risks among college students: prevalence and gender differences. *Metab Syndr Relat Disord* 2007 Dec;5(4):365-372. [doi: [10.1089/met.2007.0021](https://doi.org/10.1089/met.2007.0021)] [Medline: [18370807](https://pubmed.ncbi.nlm.nih.gov/18370807/)]
14. Huang T, Kempf A, Strother M, Li C, Lee R, Harris K, et al. Overweight and components of the metabolic syndrome in college students. *Diabetes Care* 2004 Dec;27(12):3000-3001. [Medline: [15562226](https://pubmed.ncbi.nlm.nih.gov/15562226/)]
15. Morrell JS, Byrd-Bredbenner C, Quick V, Olfert M, Dent A, Carey GB. Metabolic syndrome: comparison of prevalence in young adults at 3 land-grant universities. *J Am Coll Health* 2014;62(1):1-9. [doi: [10.1080/07448481.2013.841703](https://doi.org/10.1080/07448481.2013.841703)] [Medline: [24313691](https://pubmed.ncbi.nlm.nih.gov/24313691/)]
16. Dietz WH. Overweight in childhood and adolescence. *N Engl J Med* 2004 Feb 26;350(9):855-857. [doi: [10.1056/NEJMp048008](https://doi.org/10.1056/NEJMp048008)] [Medline: [14985480](https://pubmed.ncbi.nlm.nih.gov/14985480/)]
17. Freedman DS, Patel DA, Srinivasan SR, Chen W, Tang R, Bond MG, et al. The contribution of childhood obesity to adult carotid intima-media thickness: the Bogalusa Heart Study. *Int J Obes (Lond)* 2008 May;32(5):749-756. [doi: [10.1038/sj.ijo.0803798](https://doi.org/10.1038/sj.ijo.0803798)] [Medline: [18227845](https://pubmed.ncbi.nlm.nih.gov/18227845/)]
18. Cornette R. The emotional impact of obesity on children. *Worldviews Evid Based Nurs* 2008;5(3):136-141. [doi: [10.1111/j.1741-6787.2008.00127.x](https://doi.org/10.1111/j.1741-6787.2008.00127.x)] [Medline: [19076912](https://pubmed.ncbi.nlm.nih.gov/19076912/)]
19. Whetstone L, Morrissey S, Cummings D. Children at risk: the association between perceived weight status and suicidal thoughts and attempts in middle school youth. *J Sch Health* 2007 Feb;77(2):59-66; quiz 98. [doi: [10.1111/j.1746-1561.2007.00168.x](https://doi.org/10.1111/j.1746-1561.2007.00168.x)] [Medline: [17222156](https://pubmed.ncbi.nlm.nih.gov/17222156/)]
20. Nelson MC, Story M, Larson NI, Neumark-Sztainer D, Lytle LA. Emerging adulthood and college-aged youth: an overlooked age for weight-related behavior change. *Obesity (Silver Spring)* 2008 Oct;16(10):2205-2211 [FREE Full text] [doi: [10.1038/oby.2008.365](https://doi.org/10.1038/oby.2008.365)] [Medline: [18719665](https://pubmed.ncbi.nlm.nih.gov/18719665/)]
21. Laska MN, Pelletier JE, Larson NI, Story M. Interventions for weight gain prevention during the transition to young adulthood: a review of the literature. *J Adolesc Health* 2012 Apr;50(4):324-333 [FREE Full text] [doi: [10.1016/j.jadohealth.2012.01.016](https://doi.org/10.1016/j.jadohealth.2012.01.016)] [Medline: [22443834](https://pubmed.ncbi.nlm.nih.gov/22443834/)]
22. Sparling PB. Obesity on campus. *Prev Chronic Dis* 2007 Jul;4(3):A72 [FREE Full text] [Medline: [17572976](https://pubmed.ncbi.nlm.nih.gov/17572976/)]
23. Patience N. Weight Management Programs on Campus: What's Out There and Working? *Weight Management Matters* 2013 Dec;12(3):1-5.
24. Moreno MA, Jelenchick LA, Egan KG, Cox E, Young H, Gannon KE, et al. Feeling bad on Facebook: depression disclosures by college students on a social networking site. *Depress Anxiety* 2011 Jun;28(6):447-455 [FREE Full text] [doi: [10.1002/da.20805](https://doi.org/10.1002/da.20805)] [Medline: [21400639](https://pubmed.ncbi.nlm.nih.gov/21400639/)]
25. Walsh JR, Hebert A, Byrd-Bredbenner C, Carey G, Colby S, Brown-Esters ON, et al. The development and preliminary validation of the behavior, environment, and changeability survey (BECS). *J Nutr Educ Behav* 2012;44(6):490-499. [doi: [10.1016/j.jneb.2012.05.002](https://doi.org/10.1016/j.jneb.2012.05.002)] [Medline: [23140563](https://pubmed.ncbi.nlm.nih.gov/23140563/)]
26. US News and World Report. US News and World Report. 2013. 10 universities with the largest undergraduate student populations URL: <http://www.usnews.com/education/best-colleges/the-short-list-college/articles/2015/09/29/10-universities-with-the-highest-undergrad-enrollment> [accessed 2016-05-18] [WebCite Cache ID 6hbHopyK3]
27. Kwan MY, Arbour-Nicitopoulos KP, Lowe D, Taman S, Faulkner GE. Student reception, sources, and believability of health-related information. *J Am Coll Health* 2010;58(6):555-562. [doi: [10.1080/07448481003705925](https://doi.org/10.1080/07448481003705925)] [Medline: [20452932](https://pubmed.ncbi.nlm.nih.gov/20452932/)]
28. Clay R. Mental health issues in college on the rise: APA is working with the White House to help address the problem. *Monitor on Psychology* 2013;44(11):54.
29. Novotney A. APA partners to review college student mental health. *Monitor on Psychology* 2014;45(8):40.
30. DeAngelis T. New insights on college mental health: A network of data will offer a national picture of student well-being. *Monitor on Psychology* 2009;40(10).
31. Saloner B, Lê Cook B. An ACA provision increased treatment for young adults with possible mental illnesses relative to comparison group. *Health Aff (Millwood)* 2014 Aug;33(8):1425-1434. [doi: [10.1377/hlthaff.2014.0214](https://doi.org/10.1377/hlthaff.2014.0214)] [Medline: [25092845](https://pubmed.ncbi.nlm.nih.gov/25092845/)]
32. U.S. Department of Health and Human Services. U.S. Department of Health and Human Services. 2015. Preventive care benefits: preventive health services for adults URL: <https://www.healthcare.gov/coverage/preventive-care-benefits/> [accessed 2016-05-19] [WebCite Cache ID 6hcxGsVu0]

33. U.S. National Institutes of Health. 2015. Social media obesity treatment for college students URL: <https://clinicaltrials.gov/ct2/show/NCT02342912?id=DK100916&rank=1> [accessed 2015-12-12] [WebCite Cache ID 6hkyL13hd]
34. Vader A, Walters S, Roudsari B, Nguyen N. Where do college students get health information? Believability and use of health information sources. *Health Promot Pract* 2011 Sep;12(5):713-722. [doi: [10.1177/1524839910369995](https://doi.org/10.1177/1524839910369995)] [Medline: [21282492](https://pubmed.ncbi.nlm.nih.gov/21282492/)]
35. Stice E, Presnell K, Shaw H, Rohde P. Psychological and behavioral risk factors for obesity onset in adolescent girls: a prospective study. *J Consult Clin Psychol* 2005 Apr;73(2):195-202. [doi: [10.1037/0022-006X.73.2.195](https://doi.org/10.1037/0022-006X.73.2.195)] [Medline: [15796626](https://pubmed.ncbi.nlm.nih.gov/15796626/)]
36. Lebow, Sim LA, Kransdorf LN. Prevalence of a history of overweight and obesity in adolescents with restrictive eating disorders. *J Adolesc Health* 2015 Jan;56(1):19-24. [doi: [10.1016/j.jadohealth.2014.06.005](https://doi.org/10.1016/j.jadohealth.2014.06.005)] [Medline: [25049202](https://pubmed.ncbi.nlm.nih.gov/25049202/)]
37. Lebow J, Sim L, Kransdorf L. Prevalence of a history of overweight and obesity in adolescents with restrictive eating disorders. *J Adolesc Health* 2015 Jan;56(1):19-24. [doi: [10.1016/j.jadohealth.2014.06.005](https://doi.org/10.1016/j.jadohealth.2014.06.005)] [Medline: [25049202](https://pubmed.ncbi.nlm.nih.gov/25049202/)]
38. Spear B. Does dieting increase the risk for obesity and eating disorders? *J Am Diet Assoc* 2006 Apr;106(4):523-525. [doi: [10.1016/j.jada.2006.01.013](https://doi.org/10.1016/j.jada.2006.01.013)] [Medline: [16567146](https://pubmed.ncbi.nlm.nih.gov/16567146/)]
39. Haines J, Neumark-Sztainer D. Prevention of obesity and eating disorders: a consideration of shared risk factors. *Health Educ Res* 2006 Dec;21(6):770-782 [FREE Full text] [doi: [10.1093/her/cyl094](https://doi.org/10.1093/her/cyl094)] [Medline: [16963727](https://pubmed.ncbi.nlm.nih.gov/16963727/)]
40. Hayes S, Napolitano MA. Examination of weight control practices in a non-clinical sample of college women. *Eat Weight Disord* 2012 Sep;17(3):e157-e163. [Medline: [23086250](https://pubmed.ncbi.nlm.nih.gov/23086250/)]
41. Irving L, Neumark-Sztainer D. Integrating the prevention of eating disorders and obesity: feasible or futile? *Prev Med* 2002 Mar;34(3):299-309. [doi: [10.1006/pmed.2001.0997](https://doi.org/10.1006/pmed.2001.0997)] [Medline: [11902846](https://pubmed.ncbi.nlm.nih.gov/11902846/)]
42. Shuchman M. Falling through the cracks--Virginia Tech and the restructuring of college mental health services. *N Engl J Med* 2007 Jul 12;357(2):105-110. [doi: [10.1056/NEJMp078096](https://doi.org/10.1056/NEJMp078096)] [Medline: [17625121](https://pubmed.ncbi.nlm.nih.gov/17625121/)]
43. Gokee-LaRose J, Gorin A, Raynor H, Laska M, Jeffery R, Levy R, et al. Are standard behavioral weight loss programs effective for young adults? *Int J Obes (Lond)* 2009 Dec;33(12):1374-1380 [FREE Full text] [doi: [10.1038/ijo.2009.185](https://doi.org/10.1038/ijo.2009.185)] [Medline: [19786967](https://pubmed.ncbi.nlm.nih.gov/19786967/)]

Abbreviations

ACA: Affordable Care Act

BMI: body mass index

Edited by G Eysenbach; submitted 24.09.15; peer-reviewed by L Quintiliani, K Kattelmann; comments to author 28.10.15; revised version received 02.12.15; accepted 12.02.16; published 08.06.16

Please cite as:

Lynch S, Hayes S, Napolitano M, Hufnagel K

Availability and Accessibility of Student-Specific Weight Loss Programs and Other Risk Prevention Health Services on College Campuses

JMIR Public Health Surveill 2016;2(1):e29

URL: <http://publichealth.jmir.org/2016/1/e29/>

doi: [10.2196/publichealth.5166](https://doi.org/10.2196/publichealth.5166)

PMID: [27278261](https://pubmed.ncbi.nlm.nih.gov/27278261/)

©Sarah Lynch, Sharon Hayes, Melissa Napolitano, Katrina Hufnagel. Originally published in *JMIR Public Health and Surveillance* (<http://publichealth.jmir.org>), 08.06.2016. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Public Health and Surveillance*, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.