

Review

Randomized Controlled Trials of Technology-Based HIV/STI and Drug Abuse Preventive Interventions for African American and Hispanic Youth: Systematic Review

David Córdova¹, PhD; Frania Mendoza Lua¹, MSW; Laretta Ovadje¹, PhD; Ethan Hong²; Berenice Castillo¹, MSW; Christopher P Salas-Wright³, PhD

¹School of Social Work, University of Michigan, Ann Arbor, MI, United States

²College of Literature, Science, and the Arts, University of Michigan, Ann Arbor, MI, United States

³School of Social Work, Boston University, Boston, MA, United States

Corresponding Author:

David Córdova, PhD
School of Social Work
University of Michigan
1080 S. University Ave.
Ann Arbor, MI, 48109
United States
Phone: 1 (734) 763 6201
Fax: 1 (734) 763 3372
Email: cordovad@umich.edu

Abstract

Background: HIV/sexually transmitted infections (STIs) and drug abuse remain significant public health concerns in the United States, and African American and Hispanic youth are disproportionately affected. Although technology-based interventions are efficacious in preventing and reducing HIV/STI and licit/illicit drug use behaviors, relatively little is known regarding the state of the science of these interventions among African American and Hispanic youth.

Objective: The aim of this review is to identify and examine randomized controlled trials (RCTs) of technology-based HIV/STI and/or drug abuse preventive interventions for African American and Hispanic youth.

Methods: We searched electronic databases (ie, PubMed, Proquest, PsycINFO, Ebscohost, Google Scholar) to identify studies between January 2006 and October 2016. RCTs of technology-based interventions targeting African American and Hispanic youth HIV/STI risk behaviors, including sexual risk, licit and illicit drug use, and HIV/STI testing were included.

Results: Our search revealed a total of three studies that used an RCT design and included samples comprised of >50% African American and/or Hispanic youth. The follow-up assessments ranged from two weeks to six months and the number of participants in each trial ranged from 72 to 141. The three interventions were theory-driven, interactive, and tailored. The long-term effects of the interventions were mixed, and outcomes included reductions in sex partners, licit drug use, and condomless anal sex acts.

Conclusions: Although technology-based interventions seem promising in the prevention of HIV/STI and drug abuse among African American and Hispanic youth, more research is needed.

(*JMIR Public Health Surveill* 2017;3(4):e96) doi: [10.2196/publichealth.7129](https://doi.org/10.2196/publichealth.7129)

KEYWORDS

technology; HIV; prevention; drug use; adolescents

Introduction

HIV/sexually transmitted infections (STIs) and drug abuse remain major public health concerns in the United States. Although youth in general (defined here as adolescents 13-25 years of age) are at increased risk of HIV/STI, African American

and Hispanic youth are disproportionately affected [1,2]. African American and Hispanic youth disproportionately engage in sexual risk and licit/illicit drug use behaviors [3,4], which increase their risk for STI and HIV infection [5,6]. However, HIV testing—a National Institutes of Health HIV/acquired immune deficiency syndrome (AIDS) research priority

[7]—among African American and Hispanic youth remains underutilized [8]. Furthermore, many youths are not routinely screened for asymptomatic STIs, as recommended by the Centers for Disease Control and Prevention [2]. Youth experience multiple barriers to access HIV/STI preventive services, and exploring the use of technology as a platform to improve the delivery and uptake of such services is warranted.

Given the ubiquity of technology among youth, technology-based interventions provide an ideal platform for the dissemination of HIV/STI and drug abuse preventive interventions to populations disproportionately affected by HIV/STIs, including African American and Hispanic youth [6,9,10]. However, a gap in knowledge persists in the literature with respect to the state of the science of technology-based HIV/STI and drug abuse preventive interventions. The purpose of this review is to identify randomized controlled trials (RCTs) of technology-based HIV/STI and/or drug abuse preventive interventions for African American and Hispanic youth.

The Prevalence of HIV/STI Risk Behaviors and HIV/STIs Among African American and Hispanic Youth

African American and Hispanic youth are disproportionately at risk of engaging in HIV/STI risk behaviors, including licit and illicit drug use and sexual risk behaviors [4,11-13]. Alcohol remains the most widely used licit drug among youth [4]. National surveillance data from the *Monitoring the Future* survey indicate that 9.5% of Hispanic 8th grade youth report past 30-day alcohol use, relative to 8.2% of non-Hispanic white and 6.9% of African American 8th grade youth, respectively [4]. By 10th grade, however, these trends change such that 23.0% of non-Hispanic white youth report past 30-day alcohol use, compared to 21.4% of Hispanic and 13.9% of African American youth, respectively [4]. Furthermore, 39% of non-Hispanic white youth report past 30-day alcohol use, compared to 34.9% of Hispanic and 21.8% of African American 12th grade youth, respectively [4]. Marijuana remains the most widely used illicit drug among youth [4]. Approximately 17.5% of Hispanic and 17.2% of African American 8th grade youth report lifetime marijuana use, compared to 10.0% of non-Hispanic white 8th grade youth, respectively [3]. Furthermore, 35.2% of Hispanic and 33.0% of African American 10th grade youth report lifetime marijuana use, relative to 28.9% of non-Hispanic white 10th grade youth, respectively [4]. Moreover, 50.3% of Hispanic 12th grade youth report lifetime marijuana use, relative to 43.6% of non-Hispanic white and 41.8% of African American 12th grade youth, respectively [4]. Licit and illicit drug use behaviors increase the risk of sexual risk behaviors, which in turn increase the vulnerability to STI and HIV infections [14]. National data from the *Youth Risk Behavior Surveillance* survey indicate that 44.4% Hispanic 9th-12th grade youth report condomless sex during last sexual intercourse, compared to 43.2% of non-Hispanic white youth [5]. Furthermore, 22.8% of Hispanic and 21.8% of African American 9th-12th grade youth report alcohol or drug use before last sexual intercourse, relative to 19.3% of non-Hispanic whites [5].

Due (in part) to the aforementioned risk behaviors, youth are at disproportionate risk of STI and HIV infections. Youth between the ages of 15 and 24 represent 25% of the sexually experienced population, and comprise nearly 26% and 50% of new HIV infections and new STIs, respectively [1,2]. African American and Hispanic youth are disproportionately affected by HIV/STIs. In 2014, an estimated 9731 youth were diagnosed with HIV in the United States; 78% of those diagnoses occurred in African American and Hispanic youth [3,5]. Indeed, the majority of these infections are among young men who have sex with men (YMSM). Importantly, 86% of young females acquire HIV through heterosexual sex [3,15,16]. Furthermore, African American and Hispanic female youth are disproportionately affected by STIs, which increase the risk of HIV infection [5].

HIV and STI Testing Among African American and Hispanic Youth

Despite the disproportionate prevalence of HIV/STI risk behaviors and infections among African American and Hispanic youth, HIV testing remains low. Approximately 16.6% and 11.2% of African American and Hispanic 9th-12th grade youth report having ever been tested for HIV (not including tests when donating blood), respectively [3,5]. Although the low prevalence of HIV testing is alarming, even more disconcerting is the fact that between 2013 and 2015, HIV testing among youth has significantly decreased [5]. Concern for the downward trend in HIV testing among youth is underscored by recent shifts in the HIV/AIDS research priorities issued by the National Institutes of Health, in conjunction with practice recommendations issued by the US Preventive Services Task Force, which call for the urgent need for sound science and practice efforts aimed at improving HIV testing in youth [7,17,18].

Beyond the need to improve youth HIV testing in general, and in African American and Hispanic youth in particular, increasing STI testing remains a federal priority [7]. Screening for STIs is particularly important in preventing and reducing the transmission and acquisition of HIV, because the presence of an STI increases the risk of HIV infection [2]. Unfortunately, 60% of youth do not know they are infected with an STI, and many youths are only being tested based on their perceived risk and are not routinely screened for asymptomatic STIs [2,16]. However, this perceived risk is based, in part, on youth disclosing their HIV/STI risk behaviors to their clinician, and clinicians initiating patient-client risk communication. Researchers have shown that youth in clinical settings are not always forthcoming with respect to their HIV/STI risk behaviors, and some clinicians do not engage in youth-clinician HIV/STI risk communication [19]. Indeed, clinician-youth HIV/STI risk communication is brief or nonexistent, and some clinicians report discomfort or lack of HIV/STI risk communication training, which may partially explain the low rates of HIV and STI testing among African American and Hispanic youth [20,21].

Harnessing the Power of Technology in the Prevention of HIV/STIs and Drug Abuse Among African American and Hispanic Youth

Technology is ubiquitous in the lives of many youths in the United States. For example, among youth aged 13-17 years, 73% and 58% have access to a smartphone or tablet, respectively [22]. Relative to non-Hispanic whites, African American youth are more likely to report smartphone ownership [22]. Thanks to the prevalence of these mobile devices, many youths utilize mobile technology to access the Internet and download apps. In fact, 74% of youth 12-17 years of age report accessing the Internet via mobile devices, and 58% report downloading apps to their smartphone or tablet [22].

Given the widespread use of mobile technology, it should not be surprising that mobile health (mHealth) interventions—the science and practice of health promotion through technology—are burgeoning [23]. In fact, data indicate that between 2010 and 2015 the overall app growth was 38.1% compounded annually over the past five years, compared to 41.9% for health-related apps [24]. In 2015, the number of health-related apps surpassed 165,000 [25]. Among adult populations, national data indicate that 36% of smartphone or tablet owners report having health-related apps on their devices [26]. Of those adults with apps, 60% report the usefulness of mHealth apps in achieving health behavior goals, 35% report helpfulness for medical care decision-making, and 38% report usefulness in asking their clinician new questions or seeking a second opinion [26]. However, limited data are available among youth populations. Extending the findings from adult populations, technology-based interventions may reduce some of the challenges that youth face that keep them from participating in face-to-face interventions (eg, transportation), offer a more cost-effective option, improve and extend the quality of care, and customize and prioritize the needs and preferences of youth [26,27]. In fact, HIV and drug abuse prevention scientists suggest that technology-based interventions show promise in preventing and reducing HIV and drug abuse among youth populations, but much work remains [28,29]. However, the literature on technology-based HIV/STI and drug abuse preventive interventions among youth remains limited, and even less is known about racial and ethnic minority populations.

A systematic review that focuses on African American and Hispanic youth populations is important for several reasons. First, African American and Hispanic youth are at disproportionate risk of engaging in HIV/STI risk behaviors (including licit and illicit drug use and sexual risk behaviors), acquiring STI and HIV infections, and HIV/STI testing is underutilized in these populations. Therefore, the identification of rigorously tested, culturally congruent, preventive interventions is warranted. Second, African American and Hispanic populations encompass important biological, psychological, social, historical, political, and cultural heterogeneity that underscores the need to develop and test culturally congruent preventive interventions [30-32]. In fact, researchers have shown how culturally specific preventive interventions may lead to optimal behavioral outcomes among

African American and Hispanic youth [33,34]. Third, researchers conducting intervention research with African American and Hispanic youth populations have employed inclusion criteria with predominately African American or Hispanic samples in the identification of culturally congruent, efficacious preventive interventions [35,36]. Despite decades of prevention and intervention research, a review of RCTs comprised of primarily African American or Hispanic youth samples (ie, >70%) identified only 10 face-to-face interventions [37]. Unfortunately, the literature on technology-based interventions that are focused on African American and Hispanic youth is relatively limited. Therefore, the purpose of this review is to identify and examine RCTs of technology-based HIV/STI and/or drug abuse preventive interventions among African American and Hispanic youth.

Methods


Data Sources and Search Criteria

In the present review, we employed the methodological guidelines established for conducting systematic reviews [38-40], including the search, retrieval, selection, and coding of published articles. We searched electronic databases for relevant citations between January 2006 and October 2016, including PubMed, Proquest, PsycINFO, Ebscohost, and Google Scholar. Search terms included: “Intervention,” “Prevention,” “mHealth,” “eHealth,” “Telemedicine,” “HIV,” “STD (Sexually Transmitted Disease),” “STI,” “Risk Behavior,” “Substance Use,” “Substance Abuse,” “Adolescent,” “Web,” “Internet,” “Online,” “Mobile,” and “Phone.”

Study Selection

For the purposes of this review, we included articles if they reported on preventive interventions delivered through mobile technologies or Web-based platforms that targeted HIV/STI risk behaviors (ie, sexual risk, and licit and illicit drug use), and HIV or STI testing. Additionally, studies had to employ an RCT design and focus on youth 13-25 years of age. Following the methodological approach employed by Szapocznik et al [37], we initiated the present review of RCTs with samples comprised of primarily African American or Hispanic youth (ie, >70%). This approach resulted in only one study that met the criteria. Therefore, the inclusion criteria in the present review was revised to a >50% threshold. We excluded articles that were nonrandomized trials, narrative reviews, observational studies, qualitative studies, treatment approaches (eg, medication adherence), face-to-face only interventions, did not include behavioral outcomes (eg, intentions, beliefs, knowledge), or adult populations. Additionally, we reviewed the references of the potential articles identified to ensure that any studies that may meet inclusion criteria of the present review were included.

Following the methodological approach for conducting systematic reviews [38-40], we screened and excluded articles that did not meet inclusion criteria (eg, nonrandomized trials, intervention was not technology-based, study did not include behavioral outcomes). Once studies were excluded that did not meet inclusion criteria, the full text of the remaining studies was reviewed and three were included for the purposes of this review (Figure 1).

Figure 1. Primsa flow diagram.

Data Extraction and Interrater Reliability

We conducted an interrater reliability check. Three of the authors (DC, LO, FML) independently screened and coded the articles that were identified as having fit the search criteria. Results were compared and any discrepancies regarding which articles met inclusion criteria were discussed until consensus was reached. We extracted the following information into a spreadsheet database: method of delivery (eg, mobile application, Web-based); author; publication year; study title; targeted sample; sample size; study site; study phase (eg, efficacy, effectiveness); and study findings.

Results

As shown in [Table 1](#), our search revealed three RCTs, all of which were published in 2013 or later. The studies examined the preliminary efficacy [41], efficacy [42], and comparative effectiveness [43] of a preventive intervention. Two studies

used an intention-to-treat approach [41,43], one study used an attention control condition [42], and type of randomization was documented in two of the studies [41,42]. Sample sizes ranged from 72 [42] to 141 participants [43]. The longitudinal designs ranged from two weeks [43] to six months [42] follow-up, and two trials were registered with ClinicalTrials.gov [41,43]. The primary outcomes across the three studies included reductions in past 30-day sex partners [43], number of cigarettes smoked in the past 30 days [42], and rate of condomless anal sex acts [41].

Marsch et al [43] examined the comparative effectiveness of an interactive, customizable, Web-based intervention on sexual risk behaviors and the potential theoretical underpinnings by which change occurs (eg, skills acquisition), relative to a traditional educator-delivered intervention, among youth in an outpatient community-based substance abuse treatment clinic. Employing an intention-to-treat approach, 141 youth were randomized to either the experimental or control condition, and assessed at baseline and two-weeks postintervention.

Table 1. Randomized controlled trials of HIV/STI and/or drug abuse preventive interventions with African American and Hispanic youth.

Author	Platform	Sample	n	Study Stage	Study Design	Results
Marsch et al, 2015	Internet (Web)	Youth aged 12-18 years; 51% Hispanic, 43% Black	141	Effectiveness	RCT; face-to-face intervention or Web-delivered therapeutic education system. Follow-up conducted two weeks postbaseline.	Participants in both conditions reported reductions in past 30-day sex partners at two-week follow-up
Mason et al, 2015	Mobile (texting)	Youth aged 14-18 years; 90.8% African American	72	Efficacy	RCT; personalized motivational text messages or general health text control condition. Follow-up conducted at one, three, and six months postbaseline.	Participants in the experimental group had decreased past 30-day cigarette smoking, compared to control at six-month follow-up
Mustanski et al, 2013	Internet (Web)	Youth aged 18-24 years; 47% Hispanic, 12.7% African American	102	Preliminary efficacy	RCT; experimental condition or an online didactic HIV knowledge control condition. Follow-up conducted at six and twelve weeks.	Participants in the experimental condition reported a 44% lower rate of unprotected anal sex acts at the 12-week follow-up

Marsch et al [43] provide descriptive statistics whereby participants could indicate both their race and ethnicity, with 44.6% identifying as Black and 51.0% Hispanic, respectively. Participants in the experimental condition received a self-directed, interactive, Web-based intervention. Guided by a community reinforcement approach to behavior therapy [44], the Web-based intervention aims to prevent and reduce HIV risk behaviors by improving skills, including decision-making and negotiation skills. The Web-based intervention is comprised of a computerized risk behavior assessment, followed by a total of 26 tailored modules. Using a computer at the clinic, the intervention was delivered in 60-minute sessions and the time for completing each module ranged between 10 and 30 minutes. On average, participants completed the sessions in approximately three sessions. Participants assigned to the control condition received two group or individual sessions, each of which lasted approximately one hour. Based on the National Institute on Drug Abuse HIV prevention principles in drug-using populations, the sessions focused on the epidemiology of HIV, promotive strategies to prevent and reduce HIV risk behaviors (including sexual risk and drug use), and HIV testing information [45]. Additionally, participants received a 20-minute youth-centered HIV prevention video. Findings indicated that participants in both conditions reported reductions in past 30-day sex partners ($P=.05$), however no significant differences in condition x time were observed. Furthermore, findings suggested that participants in both conditions reported improvements in condom use skills ($P<.001$), and this difference did not significantly differ across conditions two-weeks postintervention [43].

Mason et al [42] examined the efficacy of a text-based motivational interviewing intervention, relative to an attention control condition, on reducing smoking among a sample of predominately African American (90.8%) youth. Using a respondent-driven sampling design, 72 youth were randomized to either the experimental or attention control condition via block randomization, and were assessed at baseline, 1, 3, and

6-months postbaseline. Participants in the experimental condition received a text-based intervention that delivered 30 interactive text messages and three booster text messages over a five-day period that focused on rapport building, tobacco use feedback, social network information and feedback, and summary and plans with respect to the prevention of tobacco use [42]. By integrating motivational interviewing and social network counseling, the intervention aims to prevent and reduce tobacco use by increasing psychological precursors of behavior change, and reducing risk in close peer social networks. The attention control condition consisted of 30 health-based texts, including diet, exercise, and study habits. Findings indicated that participants in the experimental condition reduced the number of cigarettes smoked in the past 30 days, relative to the attention control condition. The results yielded a large and significant effect size, $F(1,55)=4.39$, $P<.01$, $\eta^2=0.17$ [42]. Furthermore, youth in both conditions increased their readiness to change and willingness to refuse cigarettes from their friends, and decreased the number of close friends who were daily smokers, however no significant differences across conditions were found [42].

Mustanski et al [41] examined the preliminary efficacy of the *Keep It Up!* intervention, compared to an online didactic HIV knowledge condition. Using an intention-to-treat approach, 102 predominately African American (12.7%) and Hispanic (47%) YMSM were randomized via a computerized algorithm stratified by race, and assessed at baseline, immediate postintervention, prior to a 6-week booster session, and 12 weeks postintervention [41]. Participants in the experimental condition received the *Keep It Up!* intervention. Informed by the information motivation behavioral skills model of HIV risk behavior change, *Keep It Up!* consists of 7 tailored, interactive modules delivered across three sessions and completed at least 24 hours apart. *Keep It Up!* aims to prevent and reduce condomless sex via improvements in HIV knowledge, safer-sex self-efficacy, beliefs and intentions to use condoms, and condom use errors.

Table 2. Cochrane summary of findings for randomized controlled trials of HIV/STI and/or drug abuse preventive interventions with African American and Hispanic youth.

Author	Outcomes	Sample	Number of participants followed-up (studies)	Quality of the evidence (grade)	Anticipated absolute effects	
					Risk with control	Risk difference with technology intervention
Marsch et al, 2015	Reduction of 30-day sex partners	Youth aged 12-18 years; 51% Hispanic 43% African American	74 (1 RCT)	High	The mean reduction of 30-day sex partners was 0	Mean difference 5.67 higher (5.2 higher to 6.14 higher)
Mason et al, 2015	Number of cigarettes smoked in the past 30 days	Youth aged 14-18 years; 90.8% African American	72 (1 RCT)	High	The mean number of cigarettes smoked in the past 30 days was 0	Mean difference 0.6 lower (1.45 lower to 0.25 higher)
Mustanski et al, 2013	Condomless anal sex acts	Youth aged 18-24 years; 47% Hispanic, 12.7% African American	102 (1 RCT)	High	The mean condomless anal sex acts was 0	Mean difference 2.5 lower (6.19 lower to 1.19 higher)

Participants in the control condition received an HIV knowledge intervention consisting of 7 nontailored, noninteractive modules delivered across three sessions. The didactic intervention included HIV information and facts, including condom use, and the epidemiology of HIV and STIs in YMSM. Findings indicate that, relative to the control condition, participants in the experimental condition reported a 44% lower rate of unprotected anal sex acts at the 12-week follow-up (Risk Ratio=.056, $P<0.05$). Participants in both conditions showed a large increase in HIV knowledge at postintervention (Cohen's d for *Keep It Up!*=0.75; control=0.87), however no significant differences were found across conditions [41]. As shown in Table 2, we computed Cochrane summary of findings to designate the score for each criterion in each article reviewed. All three articles that were reviewed received a high grade with respect to the quality of the evidence.

Discussion

The present systematic review indicates that although advances in technology-based HIV/STI and drug abuse preventive interventions for African American and Hispanic youth have been made, only three studies that utilized an RCT design and included licit and illicit drug use or sexual risk behavior outcomes were identified. To the best of our knowledge, this is among the first systematic reviews focused on the state of the science on technology-based HIV/STI and drug abuse preventive interventions for African American and Hispanic youth. From this review, we identified four important gaps in the literature. First, there remains the need to employ methodological and statistical rigor, including the use of RCTs, to develop a fuller understanding of the effects of technology-based interventions on HIV/STI risk behaviors. Second, studies examining the long-term effects of technology-based HIV/STI and drug abuse preventive interventions are limited. Third, the identification of how, and for whom, technology-based prevention interventions do and do not work are needed. Finally, a dearth of rigorously tested technology-based HIV/STI and drug abuse preventive interventions for African American and Hispanic youth exist. Addressing these important gaps in the literature may be helpful in moving the field forward.

There remains a need to employ rigorous methodological and statistical analytic approaches to test the effects of technology-based HIV/STI and drug abuse preventive interventions. The RCT remains the standard when examining the impact of interventions [46], yet we identified only three studies that employed an RCT design. Therefore, future research should use RCT designs when examining the effects of technology-based interventions on HIV/STI and drug use among African American and Hispanic youth. Although studies examining the effects of technology-based HIV/STI and drug abuse preventive interventions using RCT designs in the United States are limited, perhaps we can draw from international studies aimed at moving the field forward. In fact, during the review process, we identified five studies conducted internationally that have employed an RCT design when examining HIV/STI risk behavioral outcomes [47-51]. For example, Ybarra et al [47] examined the effects of CyberSenga, a five-hour online healthy sexuality program on past 90-day

condom use and abstinence over a six-month period. A total of 366 youth aged 12 years and older from Mbarara, Uganda, were randomized via parallel group with adaptive randomization to CyberSenga, CyberSenga with booster sessions, or a control group [47]. Although no statistically significant results were found among the main outcomes, study findings may help inform the development and testing of future technology-based interventions. For example, it seems that abstinent youth in the CyberSenga group benefited most in the short-term, whereas sexually active participants in the CyberSenga with booster group benefited long-term [47]. Future research to examine whether (and extent to which) these findings hold true in other youth populations is needed.

The long-term effects of technology-based HIV/STI and drug abuse preventive interventions are not well understood. Of the three studies included in this review, a relatively short-term follow-up assessment (ie, range from 2 weeks to 6 months) was used. Therefore, developing a more complete understanding of the long-term effects of technology-based HIV/STI and drug abuse preventive interventions on behavioral outcomes is essential to moving the field forward. Importantly, variations in the types of drug use behaviors and vulnerability to HIV infection exist. For example, alcohol use behaviors alone are not a risk factor for HIV infection. However, alcohol use behaviors increase the risk of engaging in sexual risk behaviors, which in turn enhances vulnerability to HIV infection [52]. Injection drug use (IDU), conversely, is a direct and efficient mechanism for the transmission and acquisition of HIV [53]. Despite this knowledge, we did not identify one single study with IDU as a behavioral outcome. Given the recent rise in opioid use in the United States and links to HIV infections [54], the development and testing of technology-based interventions focused on opioid use behaviors may be warranted.

Beyond the need to establish the long-term efficacy of technology-based interventions, there also remains the need to understand the pathways through which such interventions work. Indeed, scientists have underscored the necessity of developing and testing theory-driven interventions, which can aid in identifying the mechanisms through which preventive interventions have an impact on HIV/STI testing and risk behaviors among youth. Although our review suggests that investigators have developed theory-driven technology-based preventive interventions, the testing of the theoretical underpinnings that guide these interventions is limited. In fact, our review did not reveal a single study that tested for mediation. Therefore, future research should examine the pathways through which technology-based interventions may have an effect on HIV/STI risk behaviors and testing.

There also remains a need to develop a more complete understanding of the populations for which preventive interventions do and do not work [55,56]. That is, investigators examining both malleable and nonmalleable sample and contextual study characteristics may have great utility in working toward optimally efficacious technology-based interventions. For example, some studies indicate that the effects of technology-based interventions varied as a function of gender, suggesting a potential need for gender-specific interventions [48,51,57,58].

Although stark racial and ethnic disparities exist in the incidence of new STI and HIV infections among youth [1,2], our review revealed that the majority of studies did not include samples comprised of >50% racial and ethnic minority populations. Given the facts that African American and Hispanic youth disproportionately engage in HIV/STI risk behaviors [4,5], are disproportionately affected by STI and HIV infections [1,2], and underutilize HIV and STI testing [5], more research on the development and testing of technology-based HIV/STI interventions focusing on African American and Hispanic youth is warranted. Moreover, because African Americans (12%) and Hispanics (13%) are more likely to report being smartphone-dependent compared to non-Hispanic whites (4%) [22], mobile-technology interventions utilizing smartphones may be an effective approach to engaging and retaining African American and Hispanic youth in prevention services.

Findings from the present review may have implications for practice in at least two ways. First, improving clinician-delivered HIV/STI preventive services, including screening for HIV/STI risk behaviors and status, remains a federal priority [18]; however, clinicians face numerous challenges, including interpersonal and structural-level factors. Technology-based interventions have the potential to overcome many of the challenges associated with providing HIV/STI preventive services, including improving youth-clinician HIV/STI risk communication [27], providing access to HIV/STI prevention programs [28], and supporting and enhancing traditional prevention models by supplementing face-to-face intervention efforts [59]. Second, technology-based interventions have the potential to improve clinicians' delivery of targeted, tailored, and cost-effective HIV/STI preventive services. For example, some technology-based interventions include a youth risk assessment [43,60], which can be used to deliver targeted and tailored HIV/STI prevention services based on the youth's specific risk behaviors.

Findings from the present review should be interpreted in light of an important limitation. Specifically, we did not include quasi-experimental studies in our review. RCTs may not always be a feasible intervention design, and well-designed quasi-experimental studies may have added to the evidence-base of the present review. We did not include quasi-experimental studies in our review for several reasons. First, the key distinction between quasi-experimental studies and RCTs is the random assignment of participants to either the treatment or

control group. This distinction is particularly important during analyses given the increased potential of nonequivalence between groups, yielding challenges in comparisons. Second, the aim of the present review is to establish the efficacy of technology-based HIV/STI and drug abuse preventive interventions for African American and/or Hispanic youth. As such, it was pertinent that our review focus on RCTs, which has been established as the primary method that allows for better causation implications. Finally, in our systematic review, we followed the methodological approach of including only RCTs, as employed in other studies in medical Internet research [61,62].

Conclusions

In summary, although important advances have been made in the development and testing of technology-based HIV/STI and drug abuse preventive interventions, few interventions that target *both* sexual risk and licit/illicit drug use behaviors—behaviors that often co-occur [5]—and even fewer interventions aimed at improving HIV and STI testing among youth exist [27]. Importantly, efficacious interventions for African American and Hispanic youth are limited. The results from this review demonstrate the promise of technology-based interventions to prevent and reduce new STI and HIV infections among African American and Hispanic youth. Although there is some empirical support for technology-based HIV/STI preventive interventions, much work remains. The present review identified four critical gaps, including the need to develop a more complete understanding of: (1) the effects of technology-based interventions on HIV/STI risk behaviors and testing through cutting-edge methodological and statistical approaches, including the use of RCTs; (2) the long-term effects of technology-based interventions; (3) the pathways through which such interventions work; and (4) for whom these interventions do and do not work. For example, there have been a relatively large number of studies focused on establishing the feasibility and acceptability of technology-based interventions, compared to studies focused on demonstrating the short-term and long-term behavioral outcomes [63]. Indeed, findings from this review suggest that there remains a need for the development and testing of technology-based interventions for populations most affected by HIV/STIs, including African American and Hispanic youth. This review will help advance the state of the science on technology-based youth HIV/STI and drug abuse preventive interventions.

Conflicts of Interest

None declared.

References

1. Centers for Disease Control and Prevention. HIV Surveillance Report. Atlanta, GA: Centers for Disease Control and Prevention; 2016. Diagnoses of HIV infection the United States and Dependent Areas, 2015 URL: <https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-2015-vol-27.pdf> [accessed 2017-10-01] [WebCite Cache ID 6u8QkMCiV]
2. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance. Atlanta, GA: U.S. Department of Health and Human Services; 2014. URL: <https://www.cdc.gov/std/stats14/surv-2014-print.pdf> [accessed 2016-12-06] [WebCite Cache ID 6mYXdJff2]

3. Centers for Disease Control and Prevention. HIV Among Youth. Atlanta, GA: Centers for Disease Control; 2017. URL: <https://www.cdc.gov/hiv/pdf/group/age/youth/cdc-hiv-youth.pdf> [accessed 2016-12-01] [WebCite Cache ID 6u8ROiRoR]
4. Johnston L, O'Malley P, Miech R, Bachman J, Schulenberg J. Monitoring the Future Occasional Paper Series. Ann Arbor, MI: The University of Michigan; 1975. Demographic subgroup trends among adolescents in the use of various licit and illicit drugs URL: <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ88.pdf> [accessed 2017-10-11] [WebCite Cache ID 6u8YRzZZz]
5. Kann L, McManus T, Harris W, Shanklin S, Flint K, Hawkins J, et al. Youth Risk Behavior Surveillance - United States. Atlanta, GA: Centers for Disease Control and Prevention; 2016. Morbidity and Mortality Weekly Report URL: https://www.cdc.gov/healthyyouth/data/yrbs/pdf/2015/ss6506_updated.pdf [accessed 2017-01-01] [WebCite Cache ID 6u8Si8370]
6. Córdova D, Heinze JE, Mistry R, Salas-Wright CP, Zimmerman MA. Ecodevelopmental trajectories of family functioning: links with HIV/STI risk behaviors and STI among Black adolescents. *Dev Psychol* 2016 Jul;52(7):1115-1127 [FREE Full text] [doi: [10.1037/dev0000136](https://doi.org/10.1037/dev0000136)] [Medline: [27253262](https://pubmed.ncbi.nlm.nih.gov/27253262/)]
7. National Institutes of Health. NIH HIV/AIDS research priorities and guidelines for determining AIDS funding. Bethesda: Office of AIDS Research; 2015 Aug 12. URL: <https://grants.nih.gov/grants/guide/notice-files/NOT-OD-15-137.html> [accessed 2016-12-01] [WebCite Cache ID 6mYWN9WK8]
8. Rasberry CN, Liddon N, Adkins SH, Lesesne CA, Hebert A, Kroupa E, et al. The importance of school staff referrals and follow-up in connecting high school students to HIV and STD testing. *J Sch Nurs* 2017 Apr;33(2):143-153. [doi: [10.1177/1059840516658695](https://doi.org/10.1177/1059840516658695)] [Medline: [27418443](https://pubmed.ncbi.nlm.nih.gov/27418443/)]
9. Chávez NR, Shearer LS, Rosenthal SL. Use of digital media technology for primary prevention of STIs/HIV in youth. *J Pediatr Adolesc Gynecol* 2014 Oct;27(5):244-257. [doi: [10.1016/j.jpag.2013.07.008](https://doi.org/10.1016/j.jpag.2013.07.008)] [Medline: [24332613](https://pubmed.ncbi.nlm.nih.gov/24332613/)]
10. Kumar D, Arya M. mHealth is an innovative approach to address health literacy and improve patient-physician communication - an HIV testing exemplar. *J Mob Technol Med* 2015 Jan 01;4(1):25-30 [FREE Full text] [doi: [10.7309/jmtm.4.1.6](https://doi.org/10.7309/jmtm.4.1.6)] [Medline: [25729441](https://pubmed.ncbi.nlm.nih.gov/25729441/)]
11. Cordova D, Huang S, Arzon M, Freitas D, Malcolm S, Prado G. The role of attitudes, family, peer and school on alcohol use, rule breaking and aggressive behavior in hispanic delinquent adolescents. *Open Fam Stud J* 2011;4(Suppl 1-M4):38-45 [FREE Full text] [Medline: [22473467](https://pubmed.ncbi.nlm.nih.gov/22473467/)]
12. Johnston LD, O'Malley PM, Miech RA, Bachman J. Monitoring the future national survey results on drug use, 1975-2016. Ann Arbor, MI: University of Michigan; 2016. Overview, key findings on adolescent drug use URL: <http://www.monitoringthefuture.org/pubs/monographs/mtf-overview2016.pdf> [WebCite Cache ID 6u8VFpWE2]
13. Salas-Wright CP, Vaughn MG, Todic J, Córdova D, Perron BE. Trends in the disapproval and use of marijuana among adolescents and young adults in the United States: 2002-2013. *Am J Drug Alcohol Abuse* 2015;41(5):392-404 [FREE Full text] [doi: [10.3109/00952990.2015.1049493](https://doi.org/10.3109/00952990.2015.1049493)] [Medline: [26156683](https://pubmed.ncbi.nlm.nih.gov/26156683/)]
14. Centers for Disease Control and Prevention. HIV and substance use. Atlanta: Centers for Disease Control; 2016. URL: <https://www.cdc.gov/hiv/pdf/risk/cdc-hiv-substanceuse.pdf> [WebCite Cache ID 6u8Zyg26v]
15. Centers for Disease Control and Prevention. Estimated HIV incidence in the United States. Atlanta, GA: Centers for Disease Control and Prevention; 2007. URL: https://www.cdc.gov/hiv/pdf/statistics_hssr_vol_17_no_4.pdf [accessed 2016-01-12] [WebCite Cache ID 6u8cXulZ8]
16. Centers for Disease Control and Prevention. Vital signs: HIV infection, testing, and risk behaviors among youths - United States 2012. Atlanta, GA: Centers for Disease Control and Prevention; 2012. URL: <https://www.cdc.gov/mmwr/pdf/wk/mm6147.pdf> [WebCite Cache ID 6u8d4Ai0Y]
17. Agency for Healthcare Research Quality. Guide to Clinical Preventive Services. 2014. URL: <https://www.ahrq.gov/sites/default/files/wysiwyg/professionals/clinicians-providers/guidelines-recommendations/guide/cpsguide.pdf> [WebCite Cache ID 6u8dYJffD]
18. United States Preventive Services Task Force. Screening for HIV in Adolescents and Adults: Evidence Summary. 2012. Human Immunodeficiency Virus (HIV) Infection: Screening URL: <https://www.uspreventiveservicestaskforce.org/Page/Document/screening-for-hiv-in-adolescents-and-adults-evidence-summary/human-immunodeficiency-virus-hiv-infection-screening> [accessed 2017-11-21] [WebCite Cache ID 6v9SyKJyf]
19. Córdova D, Lua FM, Ovardje L, Fessler K, Bauermeister JA, Salas-Wright CP, et al. Adolescent experiences of clinician-patient HIV/STI communication in primary care. *Health Commun* 2017 Jul 07:1-7. [doi: [10.1080/10410236.2017.1339379](https://doi.org/10.1080/10410236.2017.1339379)] [Medline: [28686489](https://pubmed.ncbi.nlm.nih.gov/28686489/)]
20. Alexander SC, Fortenberry JD, Pollak KI, Bravender T, Davis JK, Ostbye T, et al. Sexuality talk during adolescent health maintenance visits. *JAMA Pediatr* 2014 Feb;168(2):163-169 [FREE Full text] [doi: [10.1001/jamapediatrics.2013.4338](https://doi.org/10.1001/jamapediatrics.2013.4338)] [Medline: [24378686](https://pubmed.ncbi.nlm.nih.gov/24378686/)]
21. Shindel AW, Ando KA, Nelson CJ, Breyer BN, Lue TF, Smith JF. Medical student sexuality: how sexual experience and sexuality training impact U.S. and Canadian medical students' comfort in dealing with patients' sexuality in clinical practice. *Acad Med* 2010 Aug;85(8):1321-1330 [FREE Full text] [doi: [10.1097/ACM.0b013e3181e6c4a0](https://doi.org/10.1097/ACM.0b013e3181e6c4a0)] [Medline: [20671459](https://pubmed.ncbi.nlm.nih.gov/20671459/)]
22. Lenhart A. Pew Research Center. 2015. Teen, social media and technology overview 2015 URL: http://assets.pewresearch.org/wp-content/uploads/sites/14/2015/04/PI_TeensandTech_Update2015_0409151.pdf [accessed 2017-10-11] [WebCite Cache ID 6u8fzRikE]

23. Danaher BG, Brendryen H, Seeley JR, Tyler MS, Woolley T. From black box to toolbox: outlining device functionality, engagement activities, and the pervasive information architecture of mHealth interventions. *Internet Interv* 2015 Mar 1;2(1):91-101. [doi: [10.1016/j.invent.2015.01.002](https://doi.org/10.1016/j.invent.2015.01.002)] [Medline: [25750862](https://pubmed.ncbi.nlm.nih.gov/25750862/)]
24. Kalomara Information. 2014. mHealth markets world wide URL: <http://www.kaloramainformation.com/mhealth-Worldwide-9138421> [accessed 2016-12-06] [WebCite Cache ID [6mYX7ABXS](https://www.webcitation.org/6mYX7ABXS)]
25. IMS Institute for Healthcare Informatics. Patient Adoption of mHealth. Parsippany, NJ: IMS Institute for Healthcare Informatics; 2015. URL: https://www.iqvia.com/-/media/iqvia/pdfs/institute-reports/patient-adoption-of-mhealth.pdf?la=en&hash=01603C9DB399D3C96EA306056C06966C573123C9&_=1511310731481 [accessed 2016-01-12] [WebCite Cache ID [6v9TWOKzz](https://www.webcitation.org/6v9TWOKzz)]
26. Bhuyan SS, Lu N, Chandak A, Kim H, Wyant D, Bhatt J, et al. Use of mobile health applications for health-seeking behavior among US adults. *J Med Syst* 2016 Jun;40(6):153. [doi: [10.1007/s10916-016-0492-7](https://doi.org/10.1007/s10916-016-0492-7)] [Medline: [27147516](https://pubmed.ncbi.nlm.nih.gov/27147516/)]
27. Cordova D, Bauermeister JA, Fessler K, Delva J, Nelson A, Nurenberg R, et al. A community-engaged approach to developing an mHealth HIV/STI and drug abuse preventive intervention for primary care: a qualitative study. *JMIR Mhealth Uhealth* 2015;3(4):e106 [FREE Full text] [doi: [10.2196/mhealth.4620](https://doi.org/10.2196/mhealth.4620)] [Medline: [26685288](https://pubmed.ncbi.nlm.nih.gov/26685288/)]
28. Lightfoot M. HIV prevention for adolescents: where do we go from here? *Am Psychol* 2012 Nov;67(8):661-671. [doi: [10.1037/a0029831](https://doi.org/10.1037/a0029831)] [Medline: [23163452](https://pubmed.ncbi.nlm.nih.gov/23163452/)]
29. Bull S, Pratte K, Whitesell N, Rietmeijer C, McFarlane M. Effects of an Internet-based intervention for HIV prevention: the Youthnet trials. *AIDS Behav* 2009 Jun;13(3):474-487 [FREE Full text] [doi: [10.1007/s10461-008-9487-9](https://doi.org/10.1007/s10461-008-9487-9)] [Medline: [19037719](https://pubmed.ncbi.nlm.nih.gov/19037719/)]
30. Szapocznik J, Prado G, Burlew AK, Williams RA, Santisteban DA. Drug abuse in African American and Hispanic adolescents: culture, development, and behavior. *Annu Rev Clin Psychol* 2007;3:77-105. [doi: [10.1146/annurev.clinpsy.3.022806.091408](https://doi.org/10.1146/annurev.clinpsy.3.022806.091408)] [Medline: [17716049](https://pubmed.ncbi.nlm.nih.gov/17716049/)]
31. Unger JB. Cultural influences on substance use among Hispanic adolescents and young adults: findings from Project RED. *Child Dev Perspect* 2014 Mar 01;8(1):48-53 [FREE Full text] [doi: [10.1111/cdep.12060](https://doi.org/10.1111/cdep.12060)] [Medline: [24729791](https://pubmed.ncbi.nlm.nih.gov/24729791/)]
32. Castro F, Barrera M, Martinez C. The cultural adaptation of prevention interventions: resolving tensions between fidelity and fit. *Prev Sci* 2004 Mar;5(1):41-45. [Medline: [15058911](https://pubmed.ncbi.nlm.nih.gov/15058911/)]
33. Prado G, Cordova D, Cano N, Arzon M, Pantin H, Brown CH. Drug abuse preventive interventions for Hispanic youth: state of the science and implications for future research. In: *Drug Use Trajectories Among Minority Youth*. Dordrecht: Springer; 2016.
34. Brody GH, Murry VM, Gerrard M, Gibbons FX, McNair L, Brown AC, et al. The strong African American families program: prevention of youths' high-risk behavior and a test of a model of change. *J Fam Psychol* 2006 Mar;20(1):1-11. [doi: [10.1037/0893-3200.20.1.1](https://doi.org/10.1037/0893-3200.20.1.1)] [Medline: [16569084](https://pubmed.ncbi.nlm.nih.gov/16569084/)]
35. Jemmott LS, Jemmott JB, Lanier Y, Thompson C, Baker JL. Development of a barbershop-based HIV/STI risk reduction intervention for young heterosexual African American men. *Health Promot Pract* 2016 Aug 12. [doi: [10.1177/1524839916662601](https://doi.org/10.1177/1524839916662601)] [Medline: [27519261](https://pubmed.ncbi.nlm.nih.gov/27519261/)]
36. Marsiglia FF, Ayers SL, Baldwin-White A, Booth J. Changing Latino adolescents' substance use norms and behaviors: the effects of synchronized youth and parent drug use prevention interventions. *Prev Sci* 2016 Jan;17(1):1-12 [FREE Full text] [doi: [10.1007/s11121-015-0574-7](https://doi.org/10.1007/s11121-015-0574-7)] [Medline: [26103920](https://pubmed.ncbi.nlm.nih.gov/26103920/)]
37. Szapocznik J, Lopez B, Prado G, Schwartz SJ, Pantin H. Outpatient drug abuse treatment for Hispanic adolescents. *Drug Alcohol Depend* 2006 Sep;84 Suppl 1:S54-S63. [doi: [10.1016/j.drugalcdep.2006.05.007](https://doi.org/10.1016/j.drugalcdep.2006.05.007)] [Medline: [16765535](https://pubmed.ncbi.nlm.nih.gov/16765535/)]
38. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. 2011. URL: <http://www.cochrane-handbook.org/> [WebCite Cache ID [6v9UTsePP](https://www.webcitation.org/6v9UTsePP)]
39. The Steering Group of the Campbell Collaboration. Campbell policies guidelines series. *Campbell systematic reviews* 2016 Nov. [doi: [10.4073/cpg.2016.1](https://doi.org/10.4073/cpg.2016.1)]
40. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700 [FREE Full text] [Medline: [19622552](https://pubmed.ncbi.nlm.nih.gov/19622552/)]
41. Mustanski B, Garofalo R, Monahan C, Gratzner B, Andrews R. Feasibility, acceptability, and preliminary efficacy of an online HIV prevention program for diverse young men who have sex with men: the keep it up! intervention. *AIDS Behav* 2013 Nov;17(9):2999-3012 [FREE Full text] [doi: [10.1007/s10461-013-0507-z](https://doi.org/10.1007/s10461-013-0507-z)] [Medline: [23673793](https://pubmed.ncbi.nlm.nih.gov/23673793/)]
42. Mason MJ, Campbell L, Way T, Keyser-Marcus L, Benotsch E, Mennis J, et al. Development and outcomes of a text messaging tobacco cessation intervention with urban adolescents. *Subst Abus* 2015;36(4):500-506. [doi: [10.1080/08897077.2014.987946](https://doi.org/10.1080/08897077.2014.987946)] [Medline: [25551337](https://pubmed.ncbi.nlm.nih.gov/25551337/)]
43. Marsch LA, Guarino H, Grabinski MJ, Syckes C, Dillingham ET, Xie H, et al. Comparative effectiveness of Web-based vs. educator-delivered HIV prevention for adolescent substance users: a randomized, controlled trial. *J Subst Abuse Treat* 2015 Dec;59:30-37 [FREE Full text] [doi: [10.1016/j.jsat.2015.07.003](https://doi.org/10.1016/j.jsat.2015.07.003)] [Medline: [26293644](https://pubmed.ncbi.nlm.nih.gov/26293644/)]
44. Miller W, Meyers R, Hiller-Sturmhöfel S. The community-reinforcement approach. *Alcohol Res Health* 1999;23(2):116-121. [doi: [10.4324/9780203503508](https://doi.org/10.4324/9780203503508)]

45. National Institute on Drug Abuse. Principles of HIV prevention in drug-using populations. 2012. URL: <https://archives.drugabuse.gov/pdf/POHP.pdf> [accessed 2017-10-11] [WebCite Cache ID 6u8xY8287]
46. Grossman J, Mackenzie FJ. The randomized controlled trial: gold standard, or merely standard? *Perspect Biol Med* 2005;48(4):516-534. [doi: [10.1353/pbm.2005.0092](https://doi.org/10.1353/pbm.2005.0092)] [Medline: [16227664](https://pubmed.ncbi.nlm.nih.gov/16227664/)]
47. Ybarra ML, Bull SS, Prescott TL, Korchmaros JD, Bangsberg DR, Kiwanuka JP. Adolescent abstinence and unprotected sex in CyberSenga, an Internet-based HIV prevention program: randomized clinical trial of efficacy. *PLoS One* 2013;8(8):e70083 [FREE Full text] [doi: [10.1371/journal.pone.0070083](https://doi.org/10.1371/journal.pone.0070083)] [Medline: [23967069](https://pubmed.ncbi.nlm.nih.gov/23967069/)]
48. Spijkerman R, Roek MA, Vermulst A, Lemmers L, Huiberts A, Engels RC. Effectiveness of a Web-based brief alcohol intervention and added value of normative feedback in reducing underage drinking: a randomized controlled trial. *J Med Internet Res* 2010 Dec 19;12(5):e65 [FREE Full text] [doi: [10.2196/jmir.1465](https://doi.org/10.2196/jmir.1465)] [Medline: [21169172](https://pubmed.ncbi.nlm.nih.gov/21169172/)]
49. Jander A, Crutzen R, Mercken L, Candel M, de Vries H. Effects of a Web-based computer-tailored game to reduce binge drinking among Dutch adolescents: a cluster randomized controlled trial. *J Med Internet Res* 2016 Feb 03;18(2):e29 [FREE Full text] [doi: [10.2196/jmir.4708](https://doi.org/10.2196/jmir.4708)] [Medline: [26842694](https://pubmed.ncbi.nlm.nih.gov/26842694/)]
50. Jander A, Crutzen R, Mercken L, de Vries H. A Web-based computer-tailored game to reduce binge drinking among 16 to 18 year old Dutch adolescents: development and study protocol. *BMC Public Health* 2014 Oct 09;14:1054 [FREE Full text] [doi: [10.1186/1471-2458-14-1054](https://doi.org/10.1186/1471-2458-14-1054)] [Medline: [25301695](https://pubmed.ncbi.nlm.nih.gov/25301695/)]
51. Bannink R, Broeren S, Joosten-van ZE, van As E, van de Looij-Jansen P, Raat H. Effectiveness of a Web-based tailored intervention (E-health4Uth) and consultation to promote adolescents' health: randomized controlled trial. *J Med Internet Res* 2014 May 30;16(5):e143 [FREE Full text] [doi: [10.2196/jmir.3163](https://doi.org/10.2196/jmir.3163)] [Medline: [24878521](https://pubmed.ncbi.nlm.nih.gov/24878521/)]
52. Scott-Sheldon LA, Carey KB, Cunningham K, Johnson BT, Carey MP, MASH Research Team. Alcohol use predicts sexual decision-making: a systematic review and meta-analysis of the experimental literature. *AIDS Behav* 2016 Jan;20 Suppl 1:S19-S39 [FREE Full text] [doi: [10.1007/s10461-015-1108-9](https://doi.org/10.1007/s10461-015-1108-9)] [Medline: [26080689](https://pubmed.ncbi.nlm.nih.gov/26080689/)]
53. Lansky A, Finlayson T, Johnson C, Holtzman D, Wejnert C, Mitsch A, et al. Estimating the number of persons who inject drugs in the united states by meta-analysis to calculate national rates of HIV and hepatitis C virus infections. *PLoS One* 2014;9(5):e97596 [FREE Full text] [doi: [10.1371/journal.pone.0097596](https://doi.org/10.1371/journal.pone.0097596)] [Medline: [24840662](https://pubmed.ncbi.nlm.nih.gov/24840662/)]
54. Strathdee SA, Beyrer C. Threading the needle--how to stop the HIV outbreak in rural Indiana. *N Engl J Med* 2015 Jul 30;373(5):397-399. [doi: [10.1056/NEJMp1507252](https://doi.org/10.1056/NEJMp1507252)] [Medline: [26106947](https://pubmed.ncbi.nlm.nih.gov/26106947/)]
55. Cordova D, Huang S, Lally M, Estrada Y, Prado G. Do parent-adolescent discrepancies in family functioning increase the risk of Hispanic adolescent HIV risk behaviors? *Fam Process* 2014 Jun;53(2):348-363 [FREE Full text] [doi: [10.1111/famp.12067](https://doi.org/10.1111/famp.12067)] [Medline: [24617745](https://pubmed.ncbi.nlm.nih.gov/24617745/)]
56. Prado G, Huang S, Cordova D, Malcolm S, Estrada Y, Cano N, et al. Ecodevelopmental and intrapersonal moderators of a family based preventive intervention for Hispanic youth: a latent profile analysis. *Prev Sci* 2013 Jun;14(3):290-299 [FREE Full text] [doi: [10.1007/s11121-012-0326-x](https://doi.org/10.1007/s11121-012-0326-x)] [Medline: [23408280](https://pubmed.ncbi.nlm.nih.gov/23408280/)]
57. Patrick ME, Lee CM, Neighbors C. Web-based intervention to change perceived norms of college student alcohol use and sexual behavior on spring break. *Addict Behav* 2014 Mar;39(3):600-606 [FREE Full text] [doi: [10.1016/j.addbeh.2013.11.014](https://doi.org/10.1016/j.addbeh.2013.11.014)] [Medline: [24333038](https://pubmed.ncbi.nlm.nih.gov/24333038/)]
58. Starling R, Helme D, Nodulman JA, Bryan AD, Buller DB, Donohew RL, et al. Testing a risky sex behavior intervention pilot website for adolescents. *Calif J Health Promot* 2014 Dec;12(3):24-34 [FREE Full text] [Medline: [26167134](https://pubmed.ncbi.nlm.nih.gov/26167134/)]
59. Cosner BS, Singer J, Chan C. Practice innovation through technology in the digital age: a grand challenge for social work. 2013. Grand challenges for social work initiative working paper no 12 URL: <http://aaswsw.org/wp-content/uploads/2013/10/Practice-Innovation-through-Technology-in-the-Digital-Age-A-Grand-Challenge-for-Social-Work-GC-Working-Paper-No-12.pdf> [accessed 2016-12-08] [WebCite Cache ID 6mbuM2D4T]
60. Cordova D, Alers-Rojas F, Lua FM, Bauermeister J, Nurenberg R, Ovadje L, et al. The usability and acceptability of an adolescent mHealth HIV/STI and drug abuse preventive intervention in primary care. *Behav Med* 2016 May 25:1-12. [doi: [10.1080/08964289.2016.1189396](https://doi.org/10.1080/08964289.2016.1189396)] [Medline: [27223646](https://pubmed.ncbi.nlm.nih.gov/27223646/)]
61. Garg S, Garg D, Turin TC, Chowdhury MFU. Web-based interventions for chronic back pain: a systematic review. *J Med Internet Res* 2016 Jul 26;18(7):e139 [FREE Full text] [doi: [10.2196/jmir.4932](https://doi.org/10.2196/jmir.4932)] [Medline: [27460413](https://pubmed.ncbi.nlm.nih.gov/27460413/)]
62. Nicholl BI, Sandal LF, Stochkendahl MJ, McCallum M, Suresh N, Vasseljen O, et al. Digital support interventions for the self-management of low back pain: a systematic review. *J Med Internet Res* 2017 May 21;19(5):e179 [FREE Full text] [doi: [10.2196/jmir.7290](https://doi.org/10.2196/jmir.7290)] [Medline: [28550009](https://pubmed.ncbi.nlm.nih.gov/28550009/)]
63. Bull S. Beyond acceptability and feasibility: moving mHealth into impact. *Mhealth* 2016;2:45 [FREE Full text] [doi: [10.21037/mhealth.2016.12.02](https://doi.org/10.21037/mhealth.2016.12.02)] [Medline: [28293615](https://pubmed.ncbi.nlm.nih.gov/28293615/)]

Abbreviations

IDU: injection drug use

mHealth: mobile health

RCT: randomized controlled trial

STI: sexually transmitted infection

YMSM: young men who have sex with men

Edited by G Eysenbach; submitted 08.12.16; peer-reviewed by S Bull, B Mustanski, J Zhang, J Buttolph, YC Ng; comments to author 03.01.17; revised version received 12.10.17; accepted 30.10.17; published 13.12.17

Please cite as:

Córdova D, Mendoza Lua F, Ovadje L, Hong E, Castillo B, Salas-Wright CP

Randomized Controlled Trials of Technology-Based HIV/STI and Drug Abuse Preventive Interventions for African American and Hispanic Youth: Systematic Review

JMIR Public Health Surveill 2017;3(4):e96

URL: <http://publichealth.jmir.org/2017/4/e96/>

doi: [10.2196/publichealth.7129](https://doi.org/10.2196/publichealth.7129)

PMID: [29237577](https://pubmed.ncbi.nlm.nih.gov/29237577/)

©David Córdova, Frania Mendoza Lua, Lauretta Ovadje, Ethan Hong, Berenice Castillo, Christopher P Salas-Wright. Originally published in JMIR Public Health and Surveillance (<http://publichealth.jmir.org>), 13.12.2017. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.