

Viewpoint

Programmatic Mapping: Providing Evidence for High Impact HIV Prevention Programs for Female Sex Workers

Faran Emmanuel¹, MBBS, MSc (Epidemiology); Navindra Persaud², MBBS, MPH, PhD; Sharon S Weir³, PhD; Parinita Bhattacharjee⁴, MA; Shajy Isac¹, PhD

¹Centre for Global Public Health, Winnipeg, MB, Canada

²Family Health International 360, Washington DC, WA, United States

³University of North Carolina, Chapel Hill, NC, United States

⁴Partners for Health and Development in Africa, Nairobi, Kenya

Corresponding Author:

Faran Emmanuel, MBBS, MSc (Epidemiology)

Centre for Global Public Health

R070 Med Rehab Bldg, 771 McDermot Avenue

Winnipeg, MB, R3E 0T6

Canada

Phone: 1 204 975 7784

Email: faran.emmanuel@umanitoba.ca

Abstract

Programmatic mapping (PM) is a rapid and efficient mechanism to develop size estimates of key populations including female sex workers (FSWs) and geolocate them at physical locations in a systematic and scientific manner. At the macro level, this information forms the basis for allocating program resources, setting performance targets, and assess coverage. At a micro level, PM data provide specific information on hot spots, estimates of FSWs at those spots, and hot spot typology and days and times of operation, all of which provides targeted service delivery strategies. This information can provide a reliable platform to plan HIV prevention and treatment services to considerable scale and intensity. Above all, the entire PM process requires deep involvement of FSWs, which increases community ownership of the data and can lead to an increased uptake of services. Despite a few limitations, the approach is versatile and can be used in varied country contexts to generate important information about sex work and its dynamics. In this paper, we describe experiences and lessons learned from using evidence generated from PM of FSWs in multiple countries to develop HIV prevention programs at scale.

(*JMIR Public Health Surveill* 2019;5(2):e12636) doi:[10.2196/12636](https://doi.org/10.2196/12636)

KEYWORDS

programmatic mapping; PLACE; size estimation; female sex workers; HIV prevention; microplanning; key populations

Introduction

Female sex workers (FSWs) are disproportionately affected by the HIV epidemic [1]. Globally, 15% of the HIV infections among women may be attributed to sex work [2]. Primary prevention remains the key to epidemic control [3] as available evidence shows that targeted interventions implemented at scale among FSW can halt HIV transmission and reverse the epidemic [4-7]. Programs most likely to reduce new infections are community-owned, grounded in evidence and use a mix of biomedical, behavioral, and structural interventions [8,9].

Acknowledging local epidemic diversity, the Joint United Nations Programme on HIV/AIDS (UNAIDS) Practical Guidelines for Intensifying HIV Prevention recommend the use

of local data to inform programs [10,11]. This includes *knowing the epidemic* to develop a clear understanding of the epidemic and tailoring “the response” based on a combination prevention strategy. Moreover, HIV prevention programs tailored to FSWs should include active outreach, engagement, mobilization, and empowerment of FSWs to reduce their vulnerability [12,13]. Thus, for programs to be effective and accountable, they need to prioritize FSWs that can be directly reached through peer outreach and geographically located services [14]. The recently launched HIV prevention 2020 roadmap recommends a “location-population approach” that addresses the heterogeneity of the HIV epidemic and ensures effective and efficient planning and delivery of HIV prevention services [15].

More recently, The Global Fund, the World Bank, World Health Organization (WHO) and UNAIDS, as well as various national

programs have successfully used programmatic mapping (PM) approaches to estimate size, geo-distribution, operational typologies, and structural determinants of various key populations (KPs) [16]. Although various size estimation methods provide an absolute number of the key vulnerable population [17,18], PM produces estimates at the level of each hot spot and also assesses the amount of overlap between various spots. Thus, population size estimates derived from PM are adjusted for potential duplication and also provide a consequential distribution of KP members at different spots. In addition to identifying key locations and quantifying risk populations, mapping results can be used to identify existing programs and quickly institute services where they were unavailable. Thus, it provides site-level data in a timely manner to inform program design, monitor the effect of programs, and guide programs to meet the changing needs of FSWs.

The knowledge derived from PM has been used abundantly to provide a reliable platform to plan preventive and treatment services for FSWs. In this paper, we describe experiences and lessons learned from using evidence generated from PM of female sex workers in multiple countries to develop HIV prevention programs for effective coverage at scale.

What is Programmatic Mapping?

PM [16] is a systematic and scientific approach that identifies locations (often called sites, venues, or hot spots) where KPs engage in risk and provides the local and national population size estimation data necessary to plan prevention and treatment services at appropriate scale and intensity. Programmatic experience in diverse settings show that although a few FSWs operate through hidden networks, a larger proportion congregate and/or meet clients in definable geographic settings and targeting these spots for program delivery is an efficient prevention strategy [19]. To identify where to reach FSWs, PM engages the FSW community as coimplementers to estimate the number of FSWs, describe their geo-distribution and operational typologies, and inform the design and placement of HIV prevention programs to maximize coverage [20,21]. PM includes profiling locations where FSWs can be reached and assesses service availability in those locations or nearby. The term “programmatic” underscores PM’s focus on generating information useful to program design and improvement.


How Programmatic Mapping Works

During PM fieldwork, the study area is segmented into zones, and an exhaustive list of locations where FSWs congregate is systematically compiled within each zone. Subsequently, these spots are visited with the assistance of local community members (FSWs) to confirm onsite risk activities and the number of FSWs at each location. Population size is estimated at the site level, which is then adjusted to account for double counting and/or mobility across sites, to finally reach an estimated number of FSWs in the study area. Planning for and

calculating coverage requires an understanding of where sex work happens and an estimate of various subtypes in each of these locations and their characteristics. One of the key strengths of this approach lies not only in its development of size estimates but also in providing a distribution of FSWs at different hot spots. For planning services and subsequently monitoring coverage, this approach has been proven valuable in ensuring high level coverage [22].

PM includes 3 main approaches: basic geographic mapping (GM), the Priorities for Local AIDS Control Efforts (PLACE) method, and progressive mapping. Selecting the best approach depends on the context and the availability of time and resources. However, all adhere to the principles of generating location and population size information to design and implement programs, and community ownership and leadership. Although basic GM has been used primarily for mapping and estimating size of KPs, PLACE has been used pragmatically to identify locations where persons meet new partners, and thus, networks where HIV transmission is most likely to occur. Both GM and PLACE have been used where mapping and size estimates are not available and/or more rigorous baseline or follow-up estimates of program coverage are required. Progressive mapping approach is implemented as a component of a more mature program, where it is used quickly by program teams to generate information to scale up programs. A practical difference between the methods is that GM and PLACE include systematic interviews with key informants across all study zones to develop an exhaustive list of sites, whereas progressive mapping uses a crude list of sites from previous mapping as the starting point, available program data, and discussions with peer educators and outreach workers. Each method includes visits to identified spots to verify they are operational, describe their characteristics, and collect information on estimates, typology, and other operational characteristics of the hot spot. This is done by interviewing a key informant, preferably an FSW who operates from the identified spot, or someone knowledgeable about the spot (eg, a bar manager). Both GM and PLACE have extended the value of basic PM by using the venue list as a sampling frame for venue-based biobehavioral surveillance. Examples have been provided in subsequent sections on how countries with large FSW populations such as India, Kenya, and Haiti have used these approaches to gather sufficient information in a timely manner with enough geographic specificity and at low cost to plan KP program scale-up [23-25]. Other critical information such as condom availability or service availability for FSWs in the specific spots can be also collected in all the methods if such information is needed for effective program planning.

Development partners (eg, The Global Fund, the World Bank, the Bill and Melinda Gates Foundation, WHO/UNAIDS, and PEPFAR) have supported various PM approaches in a number of countries. [Figure 1](#) shows countries where various forms of PM have been implemented.

Figure 1. Countries where various forms of programmatic mapping have been implemented.

Using Programmatic Mapping for Development of Prevention Programs

There are a number of ways PM data can be used to develop strategic plans for targeted HIV prevention among KPs including FSWs and rapidly establish appropriate programs and basic services. In addition, PM has been used extensively to estimate the size of FSW populations, identify prevention coverage gaps, and quickly institute services where they are nonexistent. The following section presents some examples of how PM data have been used in various countries to strategize, develop, and implement FSW programs.

Use of Programmatic Mapping Data at a Macro Level to Develop National and Subnational Program Plans

Conceptually, high impact programs aim to reach the highest number of FSWs with quality services. At the macro level, national and subnational programming [23] focuses on establishing HIV prevention and care services, upscaling programs where needed, and launching new programs where they are nonexistent. Effective program coverage can only be measured when a realistic estimate of the number of FSWs is available.

PM data in Kenya estimated 103,298 FSWs in 7 provinces [26], which was used by National AIDS and STIs Control Program as a denominator to set national coverage targets for the minimum package of services defined in the national guidelines [27]. PM results guided decisions about where to prioritize services to improve program coverage. Thus, Nairobi (with 27% of the FSW population) was the top priority for prevention efforts, followed by Coast, Rift valley, Nyanza, and Western province. Within Nairobi, the subcounty of Starehe, with 25% of the Nairobi FSW population, was prioritized over counties

with lesser concentrations of FSWs [26]. PM data also inform progressive coverage based on resources and targets and provide insights into the heterogeneity of site typology and overall sex work so that intervention packages can be tailored to the local context [25,28]. For example, PM in Nigeria identified diverse sex work typologies based on the operational dynamics of sites and sex workers [29]. Thus, as 80% of the FSWs were venue-based in Abuja, the program focused on robust outreach and service delivery efforts at bars and night clubs and implemented less intensive programming with FSWs at other type of spots. On the basis of the number of FSWs (Lagos: 46,691 vs Anambra: 4846), 10-fold more resources were allocated for the FSW prevention program in Lagos. The geographical distribution of FSWs in Mauritius highlighted that expanding HIV prevention services in only 04 of the 09 districts would provide coverage to more than 80% of the FSW population in Mauritius, targeting specifically the high-intensity sites, thus, improving outreach and coverage within available resources [30].

Use of Programmatic Mapping Data at a Micro Level to Enhance Outreach and Services

Micro planning decentralizes outreach management and planning to a grass-root level of outreach workers (ORW) and peer educators (PEs) through a process that collects and uses data at an individual level to empower a community to make healthy decisions for themselves [31]. Although macro plans are made at the national or subnational levels, micro level planning takes place at the site level and becomes the cornerstone of peer-led programs [32].

Through PM, a list of all locations or hot spots where FSWs operate is developed, assigning a range of estimates for FSWs at each spot by subtypology (eg, street, venue, and public place). Furthermore, PM data characterize the spots by providing

information on timing of sex work activity at the spot (busiest time of the day or busiest day of the week), which is used as the basis for micro planning. Steps in micro planning include “hot spot validation” identified by PM, followed by “spot load mapping” where FSWs’ peers validate and record peak times and days of operation along with other characteristics of the spot. This is done to build ownership of the PM data by the peers, update spot lists, and estimate the number of PEs required. “Contact mapping” is then done to understand FSW networks and select PEs and determine their allocation across the target area, taking spot clustering in account. “Peer plans” are developed by PEs for each allocated FSW based on her risk and vulnerability, prioritizing those with higher risk and vulnerability. Finally, “Peer calendars” are used to document services provided to each FSW based on individual peer plans and track them at each visit outreach and micro planning [31,32]. The entire process of micro planning was conducted in Haiti using the PM data generated using the PLACE approach. These tools assisted the program to recognize each sex worker as an individual, assess her HIV risk (client volume, age, duration in sex work) and vulnerability (violence), and help plan a prevention pathway for her [33].

Likewise, in Mombasa, Kenya, PM data supported successful scale-up of HIV programs for FSWs. PM identified 9208 FSWs with the highest concentration (5809) in Kisauni and Mvita [26]. Hot spot validation and spot load mapping informed that 60% of FSWs operated through venues such as bars and clubs and another 30% through street spots. The 233 validated spots were plotted on a map, highlighting the number of FSW congregating at each spot. Location, spot size, and typology were used to allocate spots to PEs within their area of operation. On the basis of a ratio of 1 PE for 60 to 75 FSWs, 68 PEs were recruited, and 12 ORWs were hired to supervise them. Within areas of high FSW concentrations, 2 drops in centers and clinics were established, and linkage with government and NGO clinics were established for FSWs in dispersed or far away spots. Through peer plans and peer calendars, cumulative enrollment increased to 80% of the estimated FSW population within the first 10 months. An average of 35 to 40 condoms per month were distributed to each FSW during outreach as per the client volume calculated at registration [24]. As shown in the example, sex worker programs benefit tremendously from PM data in its finest details to estimate the specific number of condoms, lubes, outreach testing supplies, and other materials needed. Not only does this improve the efficiency and effectiveness of programs, it has proven its value in strengthening community networks through a community-based approach, which adds to the ownership of such programs [34].

Use of Programmatic Mapping Data to Estimate Population Size

PM has been used extensively in a number of countries to produce size estimates of sex worker populations [25,26,29,30]. Mapping produces estimates of FSWs at each spot, assesses the amount of overlap between various spots, and then adjusts the

total number for potential duplication to calculate a range of estimated number of FSWs in a focused geography. Although PM provides valid population size estimates in a timely manner to inform program design and supports monitoring and evaluation of ongoing interventions, a few limitations of the approach need to be discussed. As the inherent approach focusses on geolocation-based sex work, there is a probability to miss out FSWs who do not visit physical venues and solicit online or via social media. Thus, care needs to be taken while relying on size estimates derived from PM approaches alone. Data from PM are best used as a starting point for programs, and it is anticipated that as programs implement, they will validate and supplement the information and thereby periodically update the estimates [23,24,33]. Despite this limitation, PM has been used extensively across countries for estimating size of the KPs successfully as part of a National size estimation exercise [25,26,30].

Use of Programmatic Mapping Data for Monitoring and Research

Finally, PM data provides service delivery programs with denominators, which are crucial to be able to set goals and establish benchmarks for key outcomes indicators. Key indicators related to program coverage and utilization of programs and services by target populations serve as markers for program success. Coverage gaps at spot level can be evaluated through mapping of these spots on a continuous basis. PM data provide denominators for monitoring 90-90-90 targets and service uptake indicators across the HIV prevention, care, and treatment cascade among FSWs. In Kenya, the national program routinely uses PM data as a denominator to measure the KP program performance every quarter [35]. Under the LINKAGES project, the list of spots generated through PM contributed to the development of a sampling frame for conducting integrated bio-behavioral surveys (IBBS) of KPs in Haiti, Malawi, Angola, and Mozambique, which helped development of 90-90-90 indicators, including proportion of FSW on treatment with suppressed viral loads [25,36-38].

Likewise, Pakistan has extensively used PM data to sample KPs for IBBS. In each city, the IBBS sample was distributed based on weights derived from the respective number of KPs in each zone. Spots were randomly selected from spot lists generated by PM within each zone, and later, respondents were selected based on sampling weights assigned to each spot based on spot size, making the sample representative of the overall population [39,40]. A similar approach was employed across India, where PM data were used to select a random sample of hot spots and select respondents from the selected hot spots. As PM listed all hot spots in a selected geography, the list of spots served as a sampling frame, and it was possible to select a representative sample of KPs using a Time Location Cluster Sampling technique [41]. Thus, PM provides a sampling frame from which a representative sample of FSWs can be selected for survey. Table 1 provides the information available through mapping and its use for service delivery planning.

Table 1. Information available through programmatic mapping and its use for service delivery planning.

Information	How this information is used
National size estimates	<p>Develop national strategic plans for HIV prevention, care, and support for key populations</p> <p>Decide on national budgetary and resource allocations and costing exercises</p> <p>Set denominators for coverage targets for HIV prevention and treatment programs</p> <p>Evaluate prevention response at a national level and identify coverage gaps</p>
Subnational size estimates	<p>Help the country to decide where to prioritize programs to improve meaningful coverage</p> <p>Decide on resource allocations at a regional or subnational level</p> <p>Compare prevention response among regions and identify prevention gaps</p> <p>Draw representative samples for key population research by assigning sampling weights based on national distribution of study population</p>
Geo-distribution of spots	<p>Target locations/spots for localizing interventions</p> <p>Develop local maps and set a plan for coverage</p> <p>Prepare spot clusters and allocate spots to peer educators based on geo-proximity of spots</p> <p>Establish locations for health clinics, drop-in centers, HIV testing centers etc, as well as condom and lubricant distribution channels</p>
Number and size of spots	<p>Target locations/spots for localizing interventions</p> <p>Decide which spots to prioritize and focus to match coverage targets</p>
Operational dynamics of a spot; peak days and peak times of operations	<p>Determine human resource needs, that is, how many peer educators and outreach workers are needed to adequately cover the population. For example, 1 peer educator could work with 50-60 FSWs^a, whereas 1 outreach worker can manage working with 5-6 peer educators.</p> <p>Size estimates provided at the spot level can be used to estimate the specific number of condoms, lube, outreach testing supplies, and other materials needed</p> <p>The human resource and commodity plan needs to be based on peak estimates so that no one remains uncovered</p> <p>Peak times and peak days might be utilized to determine time of outreach</p>
Operational typology of spots/FSWs	<p>Inform intervention design based on the subtypology of spots and FSWs (eg, brothel, street, bar, night club, massage parlor based, and home-based FSWs).</p>
Use in research	<p>Mapping data is used as a sampling frame for national level surveys, including integrated bio-behavioral surveys.</p>

^aFSWs: female sex workers.

Conclusions

PM is a low-cost and reliable approach designed to generate the most critical information required for planning and implementing effective HIV prevention, care, and treatment programs. PM data have been used at national, provincial, district, subdistrict, and spot levels. At a national level, PM informs funding requirements and resource allocation, program scale-up, target setting, and coverage assessment. At a subnational level, PM prioritizes geographies where interventions should be scaled up to ensure the highest coverage possible within the available resources. Saturating coverage of KPs in high concentration geographies is preferable to spreading services thinly across a wider area. At the district or town level, PM assists programs to assess resources required for outreach and clinical staffs and for commodity requirements such as

number of condoms/lube and HIV testing kits. At a spot level, PM enhances outreach to ensure the right number of peer ORWs are available to saturate coverage as well as allocate spots to the most ideal PE. PM further improves program design by carefully evaluating typologies and functional timings of the spots and establishes denominator to monitor access to services and coverage at each spot.

Despite a few limitations, PM provides essential information about the size, distribution, and characteristics of FSWs in a systematic and scientific manner. Above all, the process relies substantially on the strength and involvement of civil society and community organizations that represent and are engaged with KPs. This practically increases the appropriateness and ownership of program design and implementation and results in increased uptake and more efficient and effective programs based on the needs and priorities of the FSW community.

Acknowledgments

The authors would like to acknowledge research teams in all countries where PM was conducted. They would like to thank the Ministries of Health in several countries where PM has been conducted in partnership. They thank international donors, that is, The World Bank, Bill and Melinda Gates Foundation, and GFATM for their funding support to conduct PM in various countries. This publication was made possible by the generous support of the American people through the United States Agency for International Development and the United States President's Emergency Plan for AIDS Relief (PEPFAR) through the LINKAGES project, cooperative agreement number AID-OAA-A-14-00045. The contents of this publication are the sole responsibility of LINKAGES and do not necessarily reflect the views of USAID, PEPFAR, or the US Government.

Authors' Contributions

FE conceptualized the whole paper, developed the first draft, and finalized the manuscript. SSW, NP, and PB wrote programmatic examples and provided additions in the discussion section. SI wrote up a large part of the conclusion section and added examples in the text.

Conflicts of Interest

None declared.

References

1. Baral S, Beyrer C, Muessig K, Poteat T, Wirtz AL, Decker MR, et al. Burden of HIV among female sex workers in low-income and middle-income countries: a systematic review and meta-analysis. *Lancet Infect Dis* 2012 Jul;12(7):538-549. [doi: [10.1016/S1473-3099\(12\)70066-X](https://doi.org/10.1016/S1473-3099(12)70066-X)] [Medline: [22424777](https://pubmed.ncbi.nlm.nih.gov/22424777/)]
2. Prüss-Ustün A, Wolf J, Driscoll T, Degenhardt L, Neira M, Calleja JM. HIV due to female sex work: regional and global estimates. *PLoS One* 2013;8(5):e63476 [FREE Full text] [doi: [10.1371/journal.pone.0063476](https://doi.org/10.1371/journal.pone.0063476)] [Medline: [23717432](https://pubmed.ncbi.nlm.nih.gov/23717432/)]
3. Ainsworth M, Teokul W. Breaking the silence: setting realistic priorities for AIDS control in less-developed countries. *Lancet* 2000 Jul 1;356(9223):55-60. [doi: [10.1016/S0140-6736\(00\)02440-5](https://doi.org/10.1016/S0140-6736(00)02440-5)] [Medline: [10892777](https://pubmed.ncbi.nlm.nih.gov/10892777/)]
4. Steen R, Wheeler T, Gorgens M, Mziray E, Dallabetta G. Feasible, efficient and necessary, without exception - Working with sex workers interrupts HIV/STI transmission and brings treatment to many in need. *PLoS One* 2015;10(10):e0121145 [FREE Full text] [doi: [10.1371/journal.pone.0121145](https://doi.org/10.1371/journal.pone.0121145)] [Medline: [26488796](https://pubmed.ncbi.nlm.nih.gov/26488796/)]
5. Steen R, Zhao P, Wi TE, PUNCHIHEWA N, Abeyewickreme I, Lo Y. Halting and reversing HIV epidemics in Asia by interrupting transmission in sex work: experience and outcomes from ten countries. *Expert Rev Anti Infect Ther* 2013 Oct;11(10):999-1015. [doi: [10.1586/14787210.2013.824717](https://doi.org/10.1586/14787210.2013.824717)] [Medline: [24124797](https://pubmed.ncbi.nlm.nih.gov/24124797/)]
6. UNAIDS. Female sex worker HIV prevention projects?: Lessons learnt from Papua New Guinea, India and Bangladesh URL: http://www.unaids.org/sites/default/files/media_asset/jc438-femsexwork_en_0.pdf [accessed 2019-04-23] [WebCite Cache ID 77rGasXlu]
7. Boily M, Pickles M, Lowndes CM, Ramesh BM, Washington R, Moses S, et al. Positive impact of a large-scale HIV prevention programme among female sex workers and clients in South India. *AIDS* 2013 Jun 1;27(9):1449-1460 [FREE Full text] [doi: [10.1097/QAD.0b013e32835fba81](https://doi.org/10.1097/QAD.0b013e32835fba81)] [Medline: [23462221](https://pubmed.ncbi.nlm.nih.gov/23462221/)]
8. UNAIDS. Combination HIV Prevention: tailoring and coordinating biomedical, behavioural and structural strategies to reduce new HIV infections URL: http://www.unaids.org/sites/default/files/media_asset/JC2007_Combination_Prevention_paper_en_0.pdf [accessed 2019-04-23] [WebCite Cache ID 77rGdgz2Z]
9. Mishra S, Steen R, Gerbase A, Lo Y, Boily M. Impact of high-risk sex and focused interventions in heterosexual HIV epidemics: a systematic review of mathematical models. *PLoS One* 2012;7(11):e50691 [FREE Full text] [doi: [10.1371/journal.pone.0050691](https://doi.org/10.1371/journal.pone.0050691)] [Medline: [23226357](https://pubmed.ncbi.nlm.nih.gov/23226357/)]
10. UNAIDS. Practical Guidelines for Intensifying HIV Prevention URL: http://data.unaids.org/pub/manual/2007/20070306_prevention_guidelines_towards_universal_access_en.pdf [accessed 2019-04-23] [WebCite Cache ID 77rGgA1O5]
11. Jones A, Cremin I, Abdullah F, Idoko J, Cherutich P, Kilonzo N, et al. Transformation of HIV from pandemic to low-endemic levels: a public health approach to combination prevention. *Lancet* 2014 Jul 19;384(9939):272-279. [doi: [10.1016/S0140-6736\(13\)62230-8](https://doi.org/10.1016/S0140-6736(13)62230-8)] [Medline: [24740087](https://pubmed.ncbi.nlm.nih.gov/24740087/)]
12. Wilson D, Halperin DT. Know your epidemic, know your response: a useful approach, if we get it right. *Lancet* 2008 Aug 9;372(9637):423-426. [doi: [10.1016/S0140-6736\(08\)60883-1](https://doi.org/10.1016/S0140-6736(08)60883-1)] [Medline: [18687462](https://pubmed.ncbi.nlm.nih.gov/18687462/)]
13. Vassall A, Chandrashekar S, Pickles M, Beattie TS, Shetty G, Bhattacharjee P, CHARME India Group, et al. Community mobilisation and empowerment interventions as part of HIV prevention for female sex workers in Southern India: a cost-effectiveness analysis. *PLoS One* 2014;9(10):e110562 [FREE Full text] [doi: [10.1371/journal.pone.0110562](https://doi.org/10.1371/journal.pone.0110562)] [Medline: [25333501](https://pubmed.ncbi.nlm.nih.gov/25333501/)]
14. World Health Organization. Implementing Comprehensive HIV / STI Programmes with Sex Workers URL: <https://www.who.int/sorry/> [accessed 2019-04-23] [WebCite Cache ID 77rGs6fwZ]
15. UNAIDS. HIV Prevention 2020 Road Map URL: http://www.unaids.org/sites/default/files/media_asset/hiv-prevention-2020-road-map_en.pdf [accessed 2019-04-23] [WebCite Cache ID 77rGxVklX]

16. GFATM. Using Programmatic Mapping to Improve Program Access and Coverage for Key Populations: Guidelines for Countries URL: https://www.theglobalfund.org/media/1291/publication_genderequalitykeypopulationsreview_report_en.pdf?u=636727910290000000 [accessed 2019-04-23] [WebCite Cache ID 77rH28ujr]
17. Sabin K, Zhao J, Garcia CJ, Sheng Y, Arias GS, Reinisch A, et al. Availability and quality of size estimations of female sex workers, men who have sex with men, people who inject drugs and transgender women in low- and middle-income countries. *PLoS One* 2016;11(5):e0155150 [FREE Full text] [doi: [10.1371/journal.pone.0155150](https://doi.org/10.1371/journal.pone.0155150)] [Medline: [27163256](https://pubmed.ncbi.nlm.nih.gov/27163256/)]
18. Abdul-Quader AS, Baughman AL, Hladik W. Estimating the size of key populations: current status and future possibilities. *Curr Opin HIV AIDS* 2014 Mar;9(2):107-114. [doi: [10.1097/COH.0000000000000041](https://doi.org/10.1097/COH.0000000000000041)] [Medline: [24393694](https://pubmed.ncbi.nlm.nih.gov/24393694/)]
19. Emmanuel F, Isac S, Blanchard JF. Using geographical mapping of key vulnerable populations to control the spread of HIV epidemics. *Expert Rev Anti Infect Ther* 2013 May;11(5):451-453. [doi: [10.1586/eri.13.33](https://doi.org/10.1586/eri.13.33)] [Medline: [23627850](https://pubmed.ncbi.nlm.nih.gov/23627850/)]
20. GFATM. The Global Fund position on size estimation and collection of spatial data, including programmatic mapping, of key populations URL: https://www.theglobalfund.org/media/1291/publication_genderequalitykeypopulationsreview_report_en.pdf?u=636727910290000000 [accessed 2019-04-23] [WebCite Cache ID 77rH8P4tx]
21. UNAIDS. 2015. Focus on Location and Populations URL: http://www.unaids.org/sites/default/files/media_asset/WAD2015_report_en_part01.pdf [accessed 2019-04-29] [WebCite Cache ID 77z7V3tXA]
22. Laga M, Vuylsteke B. Evaluating AVAHAN's design, implementation and impact: lessons learned for the HIV Prevention Community. *BMC Public Health* 2011 Dec 29;11 Suppl 6:S16 [FREE Full text] [doi: [10.1186/1471-2458-11-S6-S16](https://doi.org/10.1186/1471-2458-11-S6-S16)] [Medline: [22376320](https://pubmed.ncbi.nlm.nih.gov/22376320/)]
23. Blanchard JF, Bhattacharjee P, Kumaran S, Ramesh BM, Kumar NS, Washington RG, et al. Concepts and strategies for scaling up focused prevention for sex workers in India. *Sex Transm Infect* 2008 Oct;84(Suppl 2):ii19-ii23. [doi: [10.1136/sti.2008.033134](https://doi.org/10.1136/sti.2008.033134)] [Medline: [18799487](https://pubmed.ncbi.nlm.nih.gov/18799487/)]
24. McClarty LM, Bhattacharjee P, Isac S, Emmanuel F, Kioko J, Njiraini M, et al. Key Programme Science lessons from an HIV prevention 'Learning Site' for sex workers in Mombasa, Kenya. *Sex Transm Infect* 2018 Aug;94(5):346-352 [FREE Full text] [doi: [10.1136/sextrans-2017-053228](https://doi.org/10.1136/sextrans-2017-053228)] [Medline: [29242195](https://pubmed.ncbi.nlm.nih.gov/29242195/)]
25. FHI 360. 2018. Programmatic Mapping and Size Estimation of Key Populations in Haiti FINAL URL: <https://www.fhi360.org/sites/default/files/media/documents/resource-linkages-haiti-size-key-populations-april-2017.pdf> [accessed 2019-04-23] [WebCite Cache ID 77rHNCJB9]
26. Odek WO, Githuka GN, Avery L, Njoroge PK, Kasonde L, Gorgens M, et al. Estimating the size of the female sex worker population in Kenya to inform HIV prevention programming. *PLoS One* 2014;9(3):e89180 [FREE Full text] [doi: [10.1371/journal.pone.0089180](https://doi.org/10.1371/journal.pone.0089180)] [Medline: [24595029](https://pubmed.ncbi.nlm.nih.gov/24595029/)]
27. NASCOP. National AIDS/STD Control Programme of Kenya. Standards for Peer-Education and Outreach Programs for Sex Workers URL: http://guidelines.health.go.ke:8000/media/Sex_Worker_Guidelines.pdf [accessed 2019-04-23] [WebCite Cache ID 77rHSr4qS]
28. Buzdugan R, Copas A, Moses S, Blanchard J, Isac S, Ramesh BM, et al. Devising a female sex work typology using data from Karnataka, India. *Int J Epidemiol* 2010 Apr;39(2):439-448. [doi: [10.1093/ije/dyp351](https://doi.org/10.1093/ije/dyp351)] [Medline: [19995861](https://pubmed.ncbi.nlm.nih.gov/19995861/)]
29. Ikpeazu A, Momah-Haruna A, Mari BM, Thompson LH, Ogungbemi K, Daniel U, et al. An appraisal of female sex work in Nigeria--implications for designing and scaling up HIV prevention programmes. *PLoS One* 2014;9(8):e103619 [FREE Full text] [doi: [10.1371/journal.pone.0103619](https://doi.org/10.1371/journal.pone.0103619)] [Medline: [25118691](https://pubmed.ncbi.nlm.nih.gov/25118691/)]
30. Pathack A, Sauntally A, Soobhany S, Comins CA, Kinoo SA, Emmanuel F. Programmatic mapping to determine the size and dynamics of sex work and injecting drug use in Mauritius. *Afr J AIDS Res* 2018 Jul;17(2):129-136. [doi: [10.2989/16085906.2018.1462216](https://doi.org/10.2989/16085906.2018.1462216)] [Medline: [29745291](https://pubmed.ncbi.nlm.nih.gov/29745291/)]
31. AIDSFree. Micro-Planning in Peer Led Outreach Programs: A Handbook Based on the Experiences of the Avahan India AIDS Initiative URL: <https://aidsfree.usaid.gov/resources/prevention-update/editions/september-2013/micro-planning-peer-led-outreach-programs> [accessed 2019-04-29] [WebCite Cache ID 77z8w0kiA]
32. Partners for Health and Development in Africa. Outreach and Micro-planning URL: <http://www.phdaf.org/docs/Learning%20Site%20Case%20Study,%20Outreach%20and%20Microplanning.pdf> [accessed 2019-04-23] [WebCite Cache ID 77rHdNlmm]
33. President's Emergency Plan For AIDS Relief. PEPFAR Haiti Country Operational Plan 2017 (COP 2017) Strategic Direction Summary URL: <https://www.pepfar.gov/documents/organization/272014.pdf> [accessed 2019-04-23] [WebCite Cache ID 77rHi1mgc]
34. KHPT. World Bank. A Systematic Approach to the Design and Scale-up of Targeted Interventions for HIV Prevention among Urban Female Sex Workers URL: <http://documents.worldbank.org/curated/en/336971468151769158/pdf/NonAsciiFileName0.pdf> [accessed 2019-04-23] [WebCite Cache ID 77rHmdmGw]
35. NASCOP. Programme Quarterly Data, December 2017 URL: <https://www.nascop.or.ke/?p=1884> [accessed 2019-04-23] [WebCite Cache ID 77rHqrSNU]
36. LINKAGES. FHI 360. 2018. PLACE report Malawi URL: <https://www.fhi360.org/sites/default/files/media/documents/resource-linkages-malawi-place-report.pdf> [accessed 2019-04-23] [WebCite Cache ID 77rIAwJcN]

37. Herce M, Miller W, Bula A, Edwards J, Sapalalo P, Lancaster K, et al. Achieving the first 90 for key populations in sub-Saharan Africa through venue-based outreach: challenges and opportunities for HIV prevention based on PLACE study findings from Malawi and Angola. *J Int AIDS Soc* 2018 Jul;21(Suppl 5):e25132 [FREE Full text] [doi: [10.1002/jia2.25132](https://doi.org/10.1002/jia2.25132)] [Medline: [30033589](https://pubmed.ncbi.nlm.nih.gov/30033589/)]
38. Miller W, Manuel J, Mbofana F, Herce M, Edwards J, Rocuts F, et al. HIV testing and treatment cascade among key populations in Mozambique 22nd International AIDS Conference. 2018 Presented at: Oral abstracts of the 22nd International AIDS Conference; July 23-27, 2018; Amsterdam, the Netherlands.
39. Emmanuel F, Salim M, Akhtar N, Arshad S, Reza TE. Second-generation surveillance for HIV/AIDS in Pakistan: results from the 4th round of Integrated Behavior and Biological Survey 2011-2012. *Sex Transm Infect* 2013 Nov;89(Suppl 3):iii23-iii28 [FREE Full text] [doi: [10.1136/sextans-2013-051161](https://doi.org/10.1136/sextans-2013-051161)] [Medline: [23912818](https://pubmed.ncbi.nlm.nih.gov/23912818/)]
40. Emmanuel F, Adrien A, Athar U, Imran M, Reza T, Blanchard J. Using surveillance data for action: lessons learnt from the second generation HIV/AIDS surveillance project in Pakistan. *East Mediterr Health J* 2011 Aug;17(8):712-718. [Medline: [21977576](https://pubmed.ncbi.nlm.nih.gov/21977576/)]
41. FHI 360. Integrated Behavioural and Biological Assessment: State Summary Report for Karnataka, Bangalore URL: <https://www.fhi360.org/sites/default/files/media/documents/Integrated> [accessed 2019-04-23] [WebCite Cache ID 77rIQ2BQa]

Abbreviations

FSWs: female sex workers
GM: geographic mapping
IBBS: integrated bio-behavioral survey
KPs: key populations
ORW: outreach worker
PE: peer educator
PLACE: Priorities for Local AIDS Control Efforts
PM: programmatic mapping
UNAIDS: Joint United Nations Programme on HIV/AIDS
WHO: World Health Organization

Edited by T Sanchez; submitted 29.10.18; peer-reviewed by A Altaf, T Saidel, M Parakh; comments to author 18.01.19; revised version received 08.03.19; accepted 22.03.19; published 09.05.19

Please cite as:

Emmanuel F, Persaud N, Weir SS, Bhattacharjee P, Isac S
Programmatic Mapping: Providing Evidence for High Impact HIV Prevention Programs for Female Sex Workers
JMIR Public Health Surveill 2019;5(2):e12636
URL: <http://publichealth.jmir.org/2019/2/e12636/>
doi: [10.2196/12636](https://doi.org/10.2196/12636)
PMID:

©Faran Emmanuel, Navindra Persaud, Sharon S Weir, Parinita Bhattacharjee, Shajy Isac. Originally published in *JMIR Public Health and Surveillance* (<http://publichealth.jmir.org>), 09.05.2019. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Public Health and Surveillance*, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.