

Original Paper

# Estimation of Asthma Symptom Onset Using Internet Search Queries: Lag-Time Series Analysis

Yulin Hswen<sup>1,2</sup>, ScD, MPH; Amanda Zhang<sup>3</sup>, AB; Bruno Ventelou<sup>2</sup>, PhD

<sup>1</sup>Department of Epidemiology and Biostatistics, Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, CA, United States

<sup>2</sup>Aix Marseille University, CNRS, AMSE, Marseille, France

<sup>3</sup>Mathematics Department, Harvard University, Cambridge, MA, United States

**Corresponding Author:**

Yulin Hswen, ScD, MPH

Department of Epidemiology and Biostatistics

Bakar Computational Health Sciences Institute

University of California San Francisco

490 Illinois St

San Francisco, CA, 94158

United States

Phone: 1 415 476 1000

Email: [yulin.hswen@ucsf.edu](mailto:yulin.hswen@ucsf.edu)

## Abstract

**Background:** Asthma affects over 330 million people worldwide. Timing of an asthma event is extremely important and lack of identification of asthma increases the risk of death. A major challenge for health systems is the length of time between symptom onset and care seeking, which could result in delayed treatment initiation and worsening of symptoms.

**Objective:** This study evaluates the utility of the internet search query data for the identification of the onset of asthma symptoms.

**Methods:** Pearson correlation coefficients between the time series of hospital admissions and Google searches were computed at lag times from 4 weeks before hospital admission to 4 weeks after hospital admission. An autoregressive integrated moving average (ARIMAX) model with an autoregressive process at lags of 1 and 2 and Google searches at weeks -1 and -2 as exogenous variables were conducted to validate our correlation results.

**Results:** Google search volume for asthma had the highest correlation at 2 weeks before hospital admission. The ARIMAX model using an autoregressive process showed that the relative searches from Google about asthma were significant at lags 1 ( $P < .001$ ) and 2 ( $P = .04$ ).

**Conclusions:** Our findings demonstrate that internet search queries may provide a real-time signal for asthma events and may be useful to measure the timing of symptom onset.

(*JMIR Public Health Surveill* 2021;7(5):e18593) doi: [10.2196/18593](https://doi.org/10.2196/18593)

**KEYWORDS**

digital epidemiology; Google queries; asthma; symptoms; health information seeking

## Introduction

Asthma is a significant contributor to disease burden globally [1]. It kills around 1000 people every day and affects over 330 million people worldwide, a number that continues to rise [2]. A major challenge for health systems is the length of time between symptom onset and care seeking, which could result in delayed treatment initiation and worsening of symptoms [3,4]. Consequently, the World Health Organization has prioritized reducing asthma burden, as over 330 million people


have asthma, and it is also the most common chronic disease of childhood. Avoidable asthma deaths still occur due to lack of early identification and inappropriate management [5]. Date of hospitalization has been used in the majority of time-series studies because it is the only available administrative data on the timing of the asthma event. However, evidence has emerged that assessment exposure based on hospitalization data may generate measurement bias and lead to misclassification of time of event onset [4]. The true onset of the symptomatic event may have occurred days prior to hospital admission, leading to

underestimation of the strength of association between environmental exposures such as ambient air pollution and acute clinical asthma events [3,6,7]. This is of particular concern for asthma because of its acute event onset and because it is sensitive to short-term ambient air pollution fluctuations.

Web searching has become integral for finding health-related information. Existing evidence shows that individuals use search engines to understand their health symptoms, especially at earlier stages of their illness, before making a medical visit or use the web to decide whether to admit themselves to a health care center [8,9]. Some individuals even use information gathered

from the internet to make decisions on how to treat their illness as opposed to visiting a provider [10]. Based on these information-searching behaviors, researchers have utilized internet search queries for early identification of disease onset, which has shown to be effective for the detection of infectious disease epidemics including influenza and Ebola [11-13]. However, research has yet to evaluate the potential utility of search queries to identify the onset of asthma symptoms and minimize measurement bias (Figure 1). This study examines whether internet search queries could reveal the lag time between onset of asthma symptoms and hospital admissions due to asthma events.

**Figure 1.** Potential difference in lag time between true time of onset and hospital admission. Diagrammatic representation of exposure measurement error. Date of hospital admission is not necessarily the date of symptom onset and may lead to misclassification, as exposure measurement does not fall within the period of hospital admission. Internet search queries may identify onset of symptoms in real time and earlier than administrative data and reduce measurement error.



## Methods

### Overview

To investigate the ability of internet search queries to detect the time of onset, we analyzed the lag time between internet search activity and asthma-related hospital visits in the Provence-Alpes-Côte d'Azur (PACA) region in France. France's health care system gives the rare opportunity to make a complete account of all admissions in a given territory (in both public and private hospitals) [14]. PACA, France, was chosen to be the focus for this study because of its high national emissions of air pollution [15,16]. The number of asthma-related hospitalizations (International Classification of Diseases [ICD]-10 codes J45 and J46) was collected from diagnosis-related group (DRG)-based Program for Medicalization of Information Systems (PMSI) from all the hospitals in the PACA region and aggregated at the weekly level [17].

### Google Relative Search Volumes

Time series of weekly Google relative search volumes (RSVs) for the term topic "asthme" (asthma) restricted to the PACA region were collected from January 1, 2017, to December 31, 2017, from Google Trends [18]. Google computes RSVs by dividing the total search volume for a query in a given geographical location by the total number of queries in that region at a given point in time [19]. Therefore, these data are normalized by the population density and search volume in a given geographical area and account for temporal fluctuations. This means that when we look at the search interest for the topic of asthma, it will be proportional to all searches on all topics on Google at that time and location. This function allowed us to measure the overall interest in the topic related to asthma in this study.

Because we were specifically interested in asthma hospital admissions in the PACA region in France, other related terms, such as "difficulty breathing," were not used, as they are not specific to asthma and overlap with other respiratory conditions. Search queries related to the topic term "saignement" (bleeding)

were collected as a control, as bleeding has no direct medical connection to asthma. Pearson correlation coefficients between the time series of hospital admissions and Google searches were computed at lag times from 4 weeks before hospital admission to 4 weeks after hospital admission. We further tested the Pearson correlation results with an autoregressive analysis using explanatory variables: autoregressive integrated moving average (ARIMAX) with an autoregressive process at lags of 1 and 2 and Google searches at weeks  $-1$  and  $-2$  weeks as exogenous variables (Table 1). This allowed us to assess how Google

searches were associated with hospital admissions while accounting for autocorrelation of the hospital admissions time series. The data that support the findings of this study were obtained from Google Trends that are available from [18] and from the DRG-based PMSI under a license for this study and are not publicly available; however, these can be obtained from the authors upon reasonable request and with permission of the DRG-based PMSI. All analyses were conducted using the statsmodels package in Python.

**Table 1.** ARIMAX<sup>a</sup> regression.

Variable	Coefficient	Standard error	<i>P</i> value	95% CI
AR <sup>b</sup> -1 (hospital admissions 1 week ago)	0.83	1.15	<.001	0.41 to 1.26
AR-2 (hospital admissions 2 weeks ago)	-0.31	1.05	.04	-0.59 to -0.02
“asthme” Google searches 1 week ago	3.67	0.22	.001	1.42 to 5.92
“asthme” Google searches 2 weeks ago	3.59	0.15	.001	1.52 to 5.65
Variance on error term	461.84	81.83	<.001	301.46 to 622.22

<sup>a</sup>ARIMAX: autoregressive integrated moving average.

<sup>b</sup>AR: autoregression.

## Ethics and Consent


Public use data sets used in this study are in aggregate format and not individually identifiable such that their analysis is deemed nonhuman subject research.

## Results

Google RSVs for asthma had the highest correlation at 2 weeks before admission with a correlation of 0.491 ( $P<.001$ ; Figure

2, Table 2). Searches for “saignement” (bleeding) did not exhibit significant positive correlations with asthma-related hospital admissions at any lag time (Table 2). Our results of the Pearson correlation were further validated with our ARIMAX model, whereby the relative Google searches about asthma were significant at 1 ( $P<.001$ ) and 2 ( $P=.004$ ) weeks’ lags before hospital admissions, which were consistent with our correlation results.

**Figure 2.** Lag correlation between searches and admissions for “asthme,” 2017.



**Table 2.** Pearson correlations between Google search term and hospital admissions for asthma.

Lag time	“Asthme” (asthma) searches × asthma admissions		“Saignement” (bleeding) searches × asthma admissions	
	Correlation	<i>P</i> value	Correlation	<i>P</i> value
4 weeks before	0.327	.02	0.139	.34
3 weeks before	0.452	.001	0.123	.39
2 weeks before	0.491	<.001	0.046	.74
1 week before	0.483	<.001	−0.046	.74
Same week	0.418	.002	−0.034	.80
1 week after	0.307	.02	−0.082	.56
2 weeks after	0.013	.93	−0.220	.12
3 weeks after	−0.261	.07	−0.248	.08
4 weeks after	−0.426	.003	−0.268	.06

## Discussion

### Principal Findings

Results from our study suggest that internet search queries detect asthma symptom onset earlier than hospital admissions. Delay between time of symptom onset and time of hospital presentation for acute clinical events has been shown to result in considerable underestimation of the effects of ambient air pollution [4]. Our results show the greatest correlation at a lag of 2 weeks between Google RSVs for asthma and asthma hospital visits. In comparison to a recent study that tested the lag time between Google RSVs for terms related to COVID-19 and COVID-19 cases in Taiwan, significant time-lag correlations ranged from 0.33 to 0.72 [20]. Results from our time-series correlation were in between the range of these correlations at 0.49 ( $P < .001$ ) at 2 weeks. This effect size of 0.49 indicates a moderate relationship as correlations over 0.3 are considered to indicate an underlying relationship between 2 variables of interest [21,22]. The 1- and 2-week lag we found between asthma searches and asthma hospitalizations is consistent with previous studies that have on average a 2-week time lag between internet-based and traditional surveillance systems for disease surveillance [23]. In a study that looked at internet searches on dengue fever and local dengue occurrences, a lag time of 1 week was reported [24]. In a more recent study, the relationship between chest pain search volume on Google and new COVID-19 cases saw a lag time of 18 days (2-3 weeks) [25]. This consistent time lag of around 2 weeks may indicate the amount of time that elapsed between users developing symptoms and seeking in-person medical care.

Our results highlight that online internet search queries about symptoms may offer a novel approach to (1) identify the timing and the magnitude of future admissions, to prepare and manage resources efficiently at the hospitals (as suggested for opioids [10]) and (2) correct for measurement bias and misclassification of time of asthma onset. This finding is important, as short-term fluctuations in ambient air pollution can have significant effects on acute symptoms; therefore, hour-by-hour estimates are important to understand the impact of these environmental exposures on symptomatic changes in order to identify the onset of larger more severe health events. Epidemiological studies

measuring exposure and response should investigate the lag time between search queries and hospitalization to uncover insights about the actual timing of onset. Recent evidence has also indicated that the COVID-19 pandemic has led patients to use internet search on Google to seek out medical information and treatment in replacement of professional medical attention [25]. For instance, compared with previous years, there have been significant reductions in hospital presentations for acute myocardial infarctions and concurrent increases in out-of-hospital cardiac arrests during the COVID-19 pandemic and a marked spike in search volume for chest pain [26,27]. Therefore, internet search queries related to respiratory symptoms may offer insight into the true incidence of respiratory illnesses during COVID-19, as fear of contracting COVID-19 may prevent patients from seeking hospital care. Future studies should use internet search queries to estimate the incidence of disease, as hospital admissions may not be able to provide accurate measurements in the time of COVID-19.

### Limitations

We recognize that the population on the search engine Google may not be entirely representative of the French population. However, statistics show that Google holds the largest market share of all search engines in France (92% as of September 2020) [28]. The population of internet users in France is 82.0% and skewed toward younger age and higher education level [29]. In this study, we validated that search strategies were effective at identifying the onset of future emergency hospital use. Despite the limitation that searches on Google might not be generalizable to the entire French population, our results still suggest that this methodology may be applicable to other chronic diseases as well. However, we acknowledge that this method may not be applicable to all types of symptoms or hospital uses, especially when the disease to be treated is very specific to a subpopulation.

We also recognize that recent developments in time series such as time-series forecasting were not used for our analysis. However, in this study we sought to model trends in searches as they related to external factors such as emergency visits, whereas time-series forecasting seeks to forecast future values of that series such as using historical hospital visits to predict future hospital visits. We also chose to use the most standard

and frequently used time-series model for consistency in the research area related to environmental respiratory disease in order to identify the time lag between symptom onset and hospital admission [30-33].

### Future Directions

Based on our study findings, we believe that earlier identification of potential cases of asthma exacerbation through internet searches could help improve the efficiency of resource allocation within hospitals such as staff, beds, and respiratory assistance. Future studies should test the ability of Google searches in the hospital setting to predict cases and reduce the burden on hospitals. In addition, since the COVID-19 pandemic, it has been postulated that many patients are not seeking care for their arising symptoms because of fears of COVID-19 transmission [27]. Therefore, the use of internet search could help identify real-time and accurate onset of asthma during the

time of the COVID-19 pandemic. This information can be used to provide timely and correct patient education including informing the public about the appropriate course of action. Public health efforts should consider the utility of internet searches for respiratory conditions such as asthma to measure care-seeking behaviors and prevent severe long-term consequences.

### Conclusions

Asthma is one of the most significant noncommunicable diseases globally [34]. Improving surveillance is crucial for the control of asthma and the prevention of avoidable deaths due to this disease. The use of online digital surveillance offers the ability to capture the onset of asthma more accurately and rapidly and has the potential to reduce the burden and deaths caused by asthma.

### Acknowledgments

We thank Lisa Fressard, ORS PACA, for her assistance in retrieving the PMSI data for this study. This work was supported by the French National Research Agency Grant (No. ANR-17-EURE-0020). YH was funded by the Chateaubriand Fellowship Program, Make Our Planet Great Again, The Embassy of France.

### Conflicts of Interest

None declared.

### References

1. Guarnieri M, Balmes JR. Outdoor air pollution and asthma. *The Lancet* 2014 May;383(9928):1581-1592. [doi: [10.1016/s0140-6736\(14\)60617-6](https://doi.org/10.1016/s0140-6736(14)60617-6)]
2. Loftus PA, Wise SK. Epidemiology and economic burden of asthma. *Int Forum Allergy Rhinol* 2015 Sep 23;5 Suppl 1(S1):S7-10. [doi: [10.1002/alr.21547](https://doi.org/10.1002/alr.21547)] [Medline: [26010063](https://pubmed.ncbi.nlm.nih.gov/26010063/)]
3. Charlton I, Jones K, Bain J. Delay in diagnosis of childhood asthma and its influence on respiratory consultation rates. *Arch Dis Child* 1991 May 01;66(5):633-635 [FREE Full text] [doi: [10.1136/adc.66.5.633](https://doi.org/10.1136/adc.66.5.633)] [Medline: [2039256](https://pubmed.ncbi.nlm.nih.gov/2039256/)]
4. Lokken R, Wellenius G, Coull B, Burger MR, Schlaug G, Suh HH, et al. Air pollution and risk of stroke: underestimation of effect due to misclassification of time of event onset. *Epidemiology* 2009 Jan;20(1):137-142. [doi: [10.1097/ede.0b013e31818ef34a](https://doi.org/10.1097/ede.0b013e31818ef34a)] [Medline: [19244659](https://pubmed.ncbi.nlm.nih.gov/19244659/)]
5. Landrigan PJ, Fuller R, Acosta NJR, Adeyi O, Arnold R, Basu NN, et al. The Lancet Commission on pollution and health. *Lancet* 2018 Feb 03;391(10119):462-512. [doi: [10.1016/S0140-6736\(17\)32345-0](https://doi.org/10.1016/S0140-6736(17)32345-0)] [Medline: [29056410](https://pubmed.ncbi.nlm.nih.gov/29056410/)]
6. Lynch BA, Van Norman CA, Jacobson RM, Weaver AL, Juhn YJ. Impact of delay in asthma diagnosis on health care service use. *Allergy Asthma Proc* 2010 Jul 01;31(4):e48-e52 [FREE Full text] [doi: [10.2500/aap.2010.31.3358](https://doi.org/10.2500/aap.2010.31.3358)] [Medline: [20819315](https://pubmed.ncbi.nlm.nih.gov/20819315/)]
7. Delfino RJ, Becklake MR, Hanley JA. Reliability of hospital data for population-based studies of air pollution. *Arch Environ Health* 1993 Jun;48(3):140-146. [doi: [10.1080/00039896.1993.9940812](https://doi.org/10.1080/00039896.1993.9940812)] [Medline: [8333783](https://pubmed.ncbi.nlm.nih.gov/8333783/)]
8. The online health care revolution: How the Web helps Americans take better care of themselves. *Pew Internet & American Life: Online life report*. URL: [https://www.pewinternet.org/wp-content/uploads/sites/9/media/Files/Reports/2000/PIP\\_Health\\_Report.pdf.pdf](https://www.pewinternet.org/wp-content/uploads/sites/9/media/Files/Reports/2000/PIP_Health_Report.pdf.pdf) [accessed 2021-04-16]
9. Fox S, Rainie L, Horrigan J, Lenhart A, Spooner T, Burke M. The online health care revolution: How the Web helps Americans take better care of themselves. In: *Online*; 2000. URL: [https://www.pewinternet.org/wp-content/uploads/sites/9/media/Files/Reports/2000/PIP\\_Health\\_Report.pdf.pdf](https://www.pewinternet.org/wp-content/uploads/sites/9/media/Files/Reports/2000/PIP_Health_Report.pdf.pdf) [accessed 2021-04-15]
10. Forkner-Dunn J. Internet-based patient self-care: the next generation of health care delivery. *J Med Internet Res* 2003 May 15;5(2):e8 [FREE Full text] [doi: [10.2196/jmir.5.2.e8](https://doi.org/10.2196/jmir.5.2.e8)] [Medline: [12857664](https://pubmed.ncbi.nlm.nih.gov/12857664/)]
11. Alicino C, Bragazzi NL, Faccio V, Amicizia D, Panatto D, Gasparini R, et al. Assessing Ebola-related web search behaviour: insights and implications from an analytical study of Google Trends-based query volumes. *Infect Dis Poverty* 2015 Dec 10;4(1):54 [FREE Full text] [doi: [10.1186/s40249-015-0090-9](https://doi.org/10.1186/s40249-015-0090-9)] [Medline: [26654247](https://pubmed.ncbi.nlm.nih.gov/26654247/)]
12. Althouse BM, Ng YY, Cummings DAT. Prediction of dengue incidence using search query surveillance. *PLoS Negl Trop Dis* 2011 Aug 2;5(8):e1258 [FREE Full text] [doi: [10.1371/journal.pntd.0001258](https://doi.org/10.1371/journal.pntd.0001258)] [Medline: [21829744](https://pubmed.ncbi.nlm.nih.gov/21829744/)]
13. Yang S, Santillana M, Kou SC. Accurate estimation of influenza epidemics using Google search data via ARGO. *Proc Natl Acad Sci U S A* 2015 Nov 24;112(47):14473-14478 [FREE Full text] [doi: [10.1073/pnas.1515373112](https://doi.org/10.1073/pnas.1515373112)] [Medline: [26553980](https://pubmed.ncbi.nlm.nih.gov/26553980/)]

14. Weeks WB, Jardin M, Dufour J, Paraponaris A, Ventelou B. Geographic variation in admissions for knee replacement, hip replacement, and hip fracture in France: evidence of supplier-induced demand in for-profit and not-for-profit hospitals. *Med Care* 2014 Oct;52(10):909-917. [doi: [10.1097/MLR.0000000000000211](https://doi.org/10.1097/MLR.0000000000000211)] [Medline: [25215648](https://pubmed.ncbi.nlm.nih.gov/25215648/)]
15. Mazenq J, Dubus J, Gaudart J, Charpin D, Viudes G, Noel G. City housing atmospheric pollutant impact on emergency visit for asthma: A classification and regression tree approach. *Respir Med* 2017 Nov;132:1-8 [FREE Full text] [doi: [10.1016/j.rmed.2017.09.004](https://doi.org/10.1016/j.rmed.2017.09.004)] [Medline: [29229079](https://pubmed.ncbi.nlm.nih.gov/29229079/)]
16. Air PACA. Quelles mesures pour améliorer la qualité de l'air en Région Provence-Alpes-Côte-D'azur. Paris, France: Air PACA; 2020.
17. Moulis G, Lapeyre-Mestre M, Palmaro A, Pugnet G, Montastruc J, Sailler L. French health insurance databases: What interest for medical research? *Rev Med Interne* 2015 Jun;36(6):411-417. [doi: [10.1016/j.revmed.2014.11.009](https://doi.org/10.1016/j.revmed.2014.11.009)] [Medline: [25547954](https://pubmed.ncbi.nlm.nih.gov/25547954/)]
18. Google Trends. URL: <https://trends.google.com> [accessed 2021-04-15]
19. Wu L, Brynjolfsson E. The Future of Prediction: How Google Searches Foreshadow Housing Prices and Sales. In: Goldfarb A, Greenstein SM, Tucker CE, editors. *Economic Analysis of the Digital Economy*. Chicago, IL: University of Chicago Press; Apr 2015.
20. Husnayain A, Fuad A, Su EC. Applications of Google Search Trends for risk communication in infectious disease management: A case study of the COVID-19 outbreak in Taiwan. *Int J Infect Dis* 2020 Jun;95:221-223 [FREE Full text] [doi: [10.1016/j.ijid.2020.03.021](https://doi.org/10.1016/j.ijid.2020.03.021)] [Medline: [32173572](https://pubmed.ncbi.nlm.nih.gov/32173572/)]
21. Ratner B. The correlation coefficient: Definition. URL: <http://dmstat1.com/res/TheCorrelationCoefficientDefined.html> [accessed 2021-04-14]
22. Mukaka MM. Statistics corner: A guide to appropriate use of correlation coefficient in medical research. *Malawi Med J* 2012 Sep;24(3):69-71 [FREE Full text] [Medline: [23638278](https://pubmed.ncbi.nlm.nih.gov/23638278/)]
23. Chen Y, Zhang Y, Xu Z, Wang X, Lu J, Hu W. Avian Influenza A (H7N9) and related Internet search query data in China. *Sci Rep* 2019 Jul 18;9(1):10434 [FREE Full text] [doi: [10.1038/s41598-019-46898-y](https://doi.org/10.1038/s41598-019-46898-y)] [Medline: [31320681](https://pubmed.ncbi.nlm.nih.gov/31320681/)]
24. Li Z, Liu T, Zhu G, Lin H, Zhang Y, He J, et al. Dengue Baidu Search Index data can improve the prediction of local dengue epidemic: A case study in Guangzhou, China. *PLoS Negl Trop Dis* 2017 Mar 6;11(3):e0005354 [FREE Full text] [doi: [10.1371/journal.pntd.0005354](https://doi.org/10.1371/journal.pntd.0005354)] [Medline: [28263988](https://pubmed.ncbi.nlm.nih.gov/28263988/)]
25. Ciofani JL, Han D, Allahwala UK, Asrress KN, Bhindi R. Internet search volume for chest pain during the COVID-19 pandemic. *Am Heart J* 2021 Jan;231:157-159 [FREE Full text] [doi: [10.1016/j.ahj.2020.09.005](https://doi.org/10.1016/j.ahj.2020.09.005)] [Medline: [33010246](https://pubmed.ncbi.nlm.nih.gov/33010246/)]
26. Yom-Tov E. *Crowdsourced health: How what you do on the Internet will improve medicine*. Cambridge, MA: MIT Press; 2016.
27. Mavraganis G, Aivalioti E, Chatzidou S, Patras R, Paraskevaidis I, Kanakakis I, et al. Cardiac arrest and drug-related cardiac toxicity in the Covid-19 era. *Epidemiology, pathophysiology and management. Food Chem Toxicol* 2020 Nov;145:111742 [FREE Full text] [doi: [10.1016/j.fct.2020.111742](https://doi.org/10.1016/j.fct.2020.111742)] [Medline: [32916218](https://pubmed.ncbi.nlm.nih.gov/32916218/)]
28. Search Engine Market Share. URL: <https://gs.statcounter.com/search-engine-market-share/all/france> [accessed 2021-04-14]
29. Internet usage penetration in France from 2010 to 2019, by education level. URL: <https://www.statista.com/statistics/426201/internet-usage-penetration-study-level-france/> [accessed 2021-04-14]
30. Shokouhi M. Detecting seasonal queries by time-series analysis. New York, NY: Association for Computing Machinery; 2011 Presented at: SIGIR '11: Proceedings of the 34th international ACM SIGIR conference on Research and development in Information Retrieval; July 2011; Beijing, China.
31. Zhang Y, Milinovich G, Xu Z, Bambrick H, Mengersen K, Tong S, et al. Monitoring Pertussis Infections Using Internet Search Queries. *Sci Rep* 2017 Sep 05;7(1):10437 [FREE Full text] [doi: [10.1038/s41598-017-11195-z](https://doi.org/10.1038/s41598-017-11195-z)] [Medline: [28874880](https://pubmed.ncbi.nlm.nih.gov/28874880/)]
32. Nhung NTT, Amini H, Schindler C, Kutlar Joss M, Dien TM, Probst-Hensch N, et al. Short-term association between ambient air pollution and pneumonia in children: A systematic review and meta-analysis of time-series and case-crossover studies. *Environ Pollut* 2017 Nov;230:1000-1008. [doi: [10.1016/j.envpol.2017.07.063](https://doi.org/10.1016/j.envpol.2017.07.063)] [Medline: [28763933](https://pubmed.ncbi.nlm.nih.gov/28763933/)]
33. Ren M, Li N, Wang Z, Liu Y, Chen X, Chu Y, et al. The short-term effects of air pollutants on respiratory disease mortality in Wuhan, China: comparison of time-series and case-crossover analyses. *Sci Rep* 2017 Jan 13;7(1):40482 [FREE Full text] [doi: [10.1038/srep40482](https://doi.org/10.1038/srep40482)] [Medline: [28084399](https://pubmed.ncbi.nlm.nih.gov/28084399/)]
34. World Health Organization. Asthma (Fact Sheet). Geneva, Switzerland: World Health Organization URL: <https://www.who.int/news-room/fact-sheets/detail/asthma> [accessed 2021-05-05]

## Abbreviations

- ARIMAX:** autoregressive integrated moving average
- DRG:** diagnosis-related group
- ICD:** International Classification of Diseases
- PACA:** Provence-Alpes-Côte d'Azur
- PMSI:** Program for Medicalization of Information Systems
- RSV:** relative search volumes

*Edited by G Eysenbach; submitted 06.03.20; peer-reviewed by YH Liu, Y Cheng; comments to author 29.06.20; revised version received 26.11.20; accepted 11.03.21; published 10.05.21*

*Please cite as:*

Hswen Y, Zhang A, Ventelou B

*Estimation of Asthma Symptom Onset Using Internet Search Queries: Lag-Time Series Analysis*

*JMIR Public Health Surveill* 2021;7(5):e18593

URL: <https://publichealth.jmir.org/2021/5/e18593>

doi: [10.2196/18593](https://doi.org/10.2196/18593)

PMID:

©Yulin Hswen, Amanda Zhang, Bruno Ventelou. Originally published in JMIR Public Health and Surveillance (<https://publichealth.jmir.org>), 10.05.2021. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <https://publichealth.jmir.org>, as well as this copyright and license information must be included.