

Rapid Surveillance Report

The Annual American Men's Internet Survey of Behaviors of Men Who have Sex with Men in the United States: 2014 Key Indicators Report

Travis Sanchez, MPH, DVM; Maria Zlotorzynska, MPH, PhD; Craig Sineath, MPH; Erin Kahle, MPH, PhD; Patrick Sullivan, PhD, DVM

Emory University, Rollins School of Public Health, Atlanta, GA, United States

Corresponding Author:

Travis Sanchez, MPH, DVM

Emory University

Rollins School of Public Health

1518 Clifton Rd NE

Atlanta, GA, 30322

United States

Phone: 1 404 727 8403

Fax: 1 404 727 8737

Email: Travis.Sanchez@emory.edu

Abstract

The American Men's Internet Survey (AMIS) is an annual Web-based behavioral survey of men who have sex with men (MSM) who live in the United States. The purpose of this Rapid Surveillance Report is to report on the second cycle of data collection (November 2014 through April 2015; AMIS-2014) on the same key indicators previously reported for AMIS (December 2013 through May 2014; AMIS-2013). The AMIS survey methodology has not substantively changed since AMIS-2013. MSM were recruited from a variety of websites using banner advertisements or email blasts. Adult men currently residing in the United States were eligible to participate if they had ever had sex with a man. We examined demographic and recruitment characteristics using multivariable regression modeling ($P < .05$) stratified by the participants' self-reported human immunodeficiency virus (HIV) status. The AMIS-2014 round of data collection resulted in 9248 completed surveys from MSM representing every US state. Participants were mainly white, 40 years or older, living in the US South, living in urban/suburban areas, and recruited from a general social networking website. Self-reported HIV prevalence was 11.34% (1049/9248). Compared with HIV-negative/unknown status participants, HIV-positive participants were more likely to have had anal sex without a condom with any male partner in the past 12 months (76.55% vs 67.17%; $P < .001$) and more likely to have had anal sex without a condom with their last male sex partner who was discordant/unknown HIV status (39.66% vs 18.77%; $P < .001$). Marijuana and other illicit substance use in the past 12 months was more likely to be reported by HIV-positive participants than HIV-negative/unknown status participants (26.02% vs 21.27%, and 27.26% vs 17.60%, respectively; both $P < .001$). The vast majority (86.90%, 7127/8199) of HIV-negative/unknown status participants had been previously HIV tested, and 58.23% (4799/8199) had been tested in the past 12 months. Sexually transmitted infection (STI) testing and diagnosis was also more likely to be reported by HIV-positive participants than HIV-negative/unknown status participants (71.02% vs 37.34%, and 20.59% vs 7.54%, respectively; both $P < .001$). HIV-negative/unknown status participants <40 years of age were more likely than those 40 years or older to have had anal sex without a condom, were more likely to report substance use, were less likely to have been HIV tested, but were more likely to be tested for and diagnosed with an STI. Compared with those from general social networking, HIV-negative/unknown status participants from a geospatial social networking website were more likely to have reported all risk behaviors but were more likely to have been HIV tested, STI tested, and diagnosed with an STI.

(*JMIR Public Health Surveill* 2016;2(1):e23) doi: [10.2196/publichealth.5476](https://doi.org/10.2196/publichealth.5476)

KEYWORDS

MSM; gay; homosexual; bisexual; HIV; STD; Internet; survey; surveillance

Notice to the reader: Rapid Surveillance Reports are brief reports, which primarily report new data in table format from an existing well-described surveillance system, making a methods (and sometimes an introduction) section redundant.

The idea of this new article type is to allow rapid publication of emerging trends, or continuous publication in regular intervals of public health relevant data. If a method or system description has been published previously in JMIR Public Health Surveill or JMIR Res Protoc, the report does not have to be peer-reviewed again (although in many cases they still are).

Introduction

The American Men's Internet Survey (AMIS) is an annual Web-based behavioral survey of men who have sex with men who live in the United States. The methods have been previously published [1]. Methods in AMIS-2014 are unchanged from the previously published manuscript unless otherwise noted below.

Recruitment and Enrollment

As in the prior year, AMIS participants were recruited through convenience sampling from a variety of websites using banner advertisements or email blasts to website members (hereafter referred to generically as "ads"). Men who clicked on the ads were taken directly to the survey website hosted on a secure server administered by SurveyGizmo. To be eligible for the survey, participants had to be 15 years of age or older, consider themselves to be male, reside in the United States, and report that they had oral or anal sex with a man at least once in the past (hereafter referred to as MSM). Persons who reported being <15 years of age or refused to provide their age were not asked any other screening questions. MSM who met the eligibility criteria and consented to participate in the study started the Web-based survey immediately. The full questionnaire for AMIS-2014 is presented in [Multimedia Appendix 1](#). AMIS-2014 ran from November 2014 through April 2015, and resulted in 77,611 persons clicking on the ads and landing on the study's recruitment page ([Table 1](#)). Most were from a general social networking website (59,670/77,611, 76.88%). Nearly half (35,462/77,611 46.89%) of those who landed on the study's page started the screening process and 60.75% (47,149/77,611) were eligible. The most common reason for ineligibility was not ever having male-male sex. Nearly three-quarters (57,176/77,611, 73.67%) of those who were eligible consented to participate in the survey. There were 6.81% (1109/77,611) of the surveys determined to likely be from duplicate participants. Among unduplicated surveys, more than two-thirds (52,979/77,611, 68.25%) were considered successful. Success was defined using an examination of completed survey sections [1]. Most successful surveys were among men who reported having sex with another man in the past 12 months (9248/10,359, 89.28%).

Measures and Analyses

For AMIS-2014 analyses, we categorized participants by recruitment website and based on target audience and purpose: gay social networking (n=2), gay general interest (n=4), general social networking (n=1), and geospatial social networking (n=2). We do not provide the names of the websites to preserve operator/client privacy, particularly where a website category has only one operator. The participants who were eligible, consented, unduplicated, successful, and reported male-male

sex in the past 12 months were included in analyses of participant characteristics and behavior.

The following behavioral measures differed in AMIS-2014 from those previously published: both sexual behaviors (any condomless anal sex and condomless anal sex with a discordant/unknown status partner) were assessed for the past 12 months, binge alcohol drinking was not included, and substance using behaviors were recategorized. Human immunodeficiency virus (HIV) serostatus concordance was based on the participant's HIV status and the status of their sex partner. Discordant/unknown status was defined as either the participant or partner having unknown status or when one was HIV-negative and the other was HIV-positive. For substance-using behaviors in the past 12 months, we separated marijuana use from other illicit substance use. For AMIS-2014 all participants received questions on sexually transmitted infection (STI; chlamydia, gonorrhea, and syphilis) testing and diagnoses in the past 12 months. Participants could have been tested for an STI but not diagnosed with an STI. Persons who were diagnosed with an STI in the past 12 months all were considered to have been tested for an STI in the past 12 months.

The analysis methods for AMIS-2014 did not substantively differ from those previously published but are repeated in this report for clarity [1]. Overall chi-square tests were used to identify whether participant characteristics significantly differed between recruitment website types. Multivariable logistic regression modeling was used to determine significant differences in behaviors based on self-reported HIV status while controlling for race/ethnicity, age group, National HIV Behavioral Surveillance (NHBS) city residency, and recruitment website type. HIV-testing behaviors were only examined among those who did not report that they were HIV-positive and were also presented by participant characteristics. Multivariable logistic regression results are presented as Wald chi-square *P*-values to denote an independently significant difference in the behavior for each subgroup compared with a referent group. Statistical significance was determined at *P*<.05.

Results

Summary for AMIS-2014

Three-quarters (6819/9248, 73.73%) of participants included in this report were white, non-Hispanic, half (4676/9248, 50.6%) were ≥40 years of age, and their most common region of residence was the South followed by the West ([Table 2](#)). AMIS-2014 had participants from all US states and at least 100 participants from each of 27 states ([Figure 1](#)). Overall, 11.34% (1049/9248) of participants reported being HIV positive and 88.66% (8199/9248) reported being HIV negative or having an unknown HIV serostatus. There were significant differences in all participant characteristics based on where they were recruited ([Table 2](#)). Most of those differences were observed among participants recruited from geospatial social networking websites, who were less likely be white, less likely be 40 years or older, less likely to live in an NHBS city, more likely to live in the West, more likely to live in urban areas, and more likely to report being HIV positive.

Most participants had anal sex without a condom with another man in the past 12 months (Table 3). Compared with HIV-negative/unknown status participants, those who were HIV-positive were significantly more likely to report anal intercourse without a condom, including with male partners who were discordant/unknown status. Anal intercourse without a condom significantly differed by age group (HIV-positive and -negative/unknown status participants), recruitment website type (HIV-positive and -negative/unknown status participants), and race/ethnicity (HIV-negative/unknown status participants only).

More than one-quarter (273/1049, 26.02%) of HIV-positive participants reported using marijuana or other illicit substances in the past 12 months (Table 4). Compared with HIV-negative/unknown status participants, those who were HIV-positive were significantly more likely to report use of marijuana and other substances in the past 12 months. Marijuana or other illicit substance use significantly differed by age group (HIV-positive and -negative/unknown status participants), residence in an NHBS city (HIV-negative/unknown status participants only), and recruitment website type (HIV-negative/unknown status participants only).

HIV testing behaviors were only examined among those who did not report being HIV-positive. Most of those participants (7125/8199, 86.90%) had ever been previously tested for HIV infection, and just over half (4799/8199, 58.53%) reported being tested in the past 12 months (Table 5). HIV testing significantly differed by age group (ever tested), race/ethnicity (ever tested), residence in an NHBS city (past 12 months tested), and recruitment website type (past 12 months tested).

Compared with HIV-negative/unknown status participants, those who were HIV-positive were more likely to report being tested for and diagnosed with an STI in the past 12 months (Table 6). The most common STI diagnoses were syphilis (132/1049, 12.58%) and chlamydia (88/1049, 8.39%) among HIV-positive participants. STI testing significantly differed by age group, residence in an NHBS city and recruitment website type only for participants who were HIV-negative/unknown status. STI diagnosis significantly differed by age group (HIV-positive and HIV-negative/unknown status participants), race/ethnicity (HIV-negative/unknown status participants only), residence in an NHBS city (HIV-negative/unknown status participants only), and recruitment website type (HIV-negative/unknown status participants only).

Figure 1. Number of MSM participants in the American Men's Internet Survey by state, 2014.




Table 1. Recruitment outcomes with different recruitment website types for the American Men's Internet Survey, United States, 2014.

Recruitment Outcomes	Recruitment Website Type									
	Total		Gay social networking (n=2)		General gay interest (n=4)		General social networking (n=1)		Geospatial social networking (n=2)	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
Clicked ad	77,611		1988		8372		59,670		7581	
Screened ^a	36,392	(46.89)	944	(47.48)	1293	(15.44)	26,576	(44.54)	7579	(99.97)
Ineligible ^b	14,285	(39.25)	171	(18.11)	523	(40.45)	11,356	(42.73)	2235	(29.49)
Not 15+ years of age ^c	10,219	(71.54)	118	(69.01)	340	(65.01)	8008	(70.52)	1753	(78.43)
Not male ^c	10,942	(76.60)	138	(80.70)	380	(72.66)	8494	(74.80)	1930	(86.35)
Not ever MSM ^{c,d}	13,776	(96.44)	167	(97.66)	510	(97.51)	11,073	(97.51)	2026	(90.65)
Not a US resident ^c	3733	(26.13)	22	(12.87)	161	(30.78)	2460	(21.66)	1090	(48.77)
Eligible ^b	22,107	(60.75)	773	(81.89)	770	(59.55)	15,220	(57.27)	5344	(70.51)
Consented ^e	16,286	(73.67)	574	(74.26)	595	(77.27)	10,821	(71.10)	4296	(80.39)
Unduplicated ^f	15,177	(93.19)	557	(97.04)	564	(94.79)	9960	(92.04)	4096	(95.34)
Success ^g	10,359	(68.25)	414	(74.33)	410	(72.70)	6913	(69.41)	2622	(64.01)
MSM past 12 months ^h	9248	(89.28)	377	(91.06)	369	(90.00)	5987	(86.60)	2515	(95.92)

^aProportion is of total who clicked ad. Includes those who started the screening questionnaire.

^bProportion is among total screened. Ineligible includes those who did not complete the screening questionnaire.

^cProportion is among total ineligible. Includes those who may not have responded to the question.

^dMSM: men who have sex with men.

^eProportion is among eligible.

^fProportion is among consented. Unduplicated removes participants who were marked as duplicates using an Internet protocol address and demographic data matching.

^gProportion is among unduplicated. Success removes participants who did not pass the test for survey completeness.

^hProportion is among successes.

Table 2. Characteristics of MSM participants in the American Men's Internet Survey by recruitment website type, United States, 2014.

Participant characteristics	Recruitment website type										<i>P</i> -value ^a
	Total		Gay social networking (n=2)		General gay interest (n=4)		General social networking (n=1)		Geospatial social networking (n=2)		
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	
Race/ethnicity											<.001
Black, non-Hispanic	415	(4.49)	11	(2.92)	21	(5.69)	225	(3.76)	158	(6.28)	
Hispanic ^b	1308	(14.14)	14	(3.71)	37	(10.03)	713	(11.91)	544	(21.63)	
White, non-Hispanic	6819	(73.73)	327	(86.74)	286	(77.51)	4643	(77.55)	1563	(62.15)	
Other or multiple races	706	(7.63)	25	(6.63)	25	(6.78)	406	(6.78)	250	(9.94)	
Age (years)											<.001
15-24	1389	(15.02)	21	(5.57)	47	(12.74)	857	(14.31)	464	(18.45)	
25-29	1221	(13.20)	24	(6.37)	48	(13.01)	612	(10.22)	537	(21.35)	
30-39	1962	(21.22)	40	(10.61)	86	(23.31)	1164	(19.44)	672	(26.72)	
40 or older	4676	(50.56)	292	(77.45)	188	(50.95)	3354	(56.02)	842	(33.48)	
Region											<.001
Midwest	1560	(16.87)	68	(18.04)	64	(17.34)	985	(16.45)	443	(17.61)	
Northeast	1933	(20.90)	117	(31.03)	55	(14.91)	1306	(21.81)	455	(18.09)	
South	3634	(39.29)	122	(32.36)	180	(48.78)	2369	(39.57)	963	(38.29)	
West	2110	(22.82)	69	(18.30)	65	(17.62)	1323	(22.10)	653	(25.96)	
US dependent areas	11	(0.12)	1	(0.27)	5	(1.36)	4	(0.07)	1	(0.04)	
NHBS city resident ^c											<.001
Yes	3553	(38.42)	137	(36.34)	206	(55.83)	2177	(36.36)	1033	(41.07)	
No	5695	(61.58)	240	(63.66)	163	(44.17)	3810	(63.64)	1482	(58.93)	
Population density ^d											<.001
Rural	2774	(30.00)	133	(35.28)	80	(21.68)	1902	(31.77)	659	(26.20)	
Urban/suburban	6300	(68.12)	241	(63.93)	285	(77.24)	4001	(66.83)	1773	(70.50)	
Self-reported HIV status											<.001
Positive	1049	(11.34)	23	(6.10)	32	(8.67)	554	(9.25)	440	(17.50)	
Negative	6992	(75.61)	277	(73.47)	303	(82.11)	4588	(76.63)	1824	(72.52)	
Unknown	1207	(13.05)	77	(20.42)	34	(9.21)	845	(14.11)	251	(9.98)	
Total	9248		377		369		5987		2515		

^aChi-square testing difference in characteristics between website type.^bHispanic persons could have been of any race, including other or multiple.^cNHBS: National HIV Behavioral Surveillance System.^dThere were 71 participants missing information needed to determine the population density of the area where they lived.

Table 3. Sexual behaviors with male partners of MSM participants in the American Men's Internet Survey, United States, 2014.

Participant characteristics	N in sample	Sexual Behaviors with male partners in the past 12 months					
		Anal intercourse without a condom			Anal intercourse without a condom with a partner of discordant or unknown HIV status		
		n	(%)	P-value ^a	n	(%)	P-value ^a
HIV positive overall	1049	803	(76.55)	<.001^b	416	(39.66)	<.001^b
Race/ethnicity							
Black, non-Hispanic	92	69	(75.00)	.459	34	(36.96)	.253
Hispanic	172	134	(77.91)	.636	73	(42.44)	.893
White, non-Hispanic	716	546	(76.26)	REF	281	(39.25)	REF
Other or multiple races	69	54	(78.26)	.878	28	(40.58)	.794
Age (years)							
15-24	68	62	(91.18)	.038	38	(55.88)	.090
25-29	110	96	(87.27)	.226	65	(59.09)	.002
30-39	251	199	(79.28)	.086	111	(44.22)	.278
40 or older	620	446	(71.94)	REF	202	(32.58)	REF
NHBS city resident ^c							
Yes	464	361	(77.80)	.217	173	(37.28)	.291
No	585	442	(75.56)	REF	243	(41.54)	REF
Recruitment website type							
Gay social networking	23	18	(78.26)	.620	16	(69.57)	.007
General gay interest	32	24	(75.00)	.768	17	(53.13)	.478
General social networking	554	403	(72.74)	REF	192	(34.66)	REF
Geospatial social networking	440	358	(81.36)	.642	191	(43.41)	.002
HIV negative or unknown overall	8199	5507	(67.17)	REF	1539	(18.77)	REF
Race/ethnicity							
Black, non-Hispanic	323	202	(62.54)	.043	80	(24.77)	.121
Hispanic	1136	777	(68.40)	.377	276	(24.30)	.088
White, non-Hispanic	6103	4109	(67.33)	REF	1053	(17.25)	REF
Other or multiple races	637	419	(65.78)	.808	130	(20.41)	.323
Age (years)							
15-24	1321	870	(65.86)	<.001	334	(25.28)	.359
25-29	1111	820	(73.81)	<.001	193	(17.37)	.954
30-39	1711	1281	(74.87)	<.001	241	(14.09)	.572
40 or older	4056	2536	(62.52)	REF	505	(12.45)	REF
NHBS city resident ^c							
Yes	3089	2053	(66.46)	.300	607	(19.65)	.442
No	5110	3454	(67.59)	REF	932	(18.24)	REF
Recruitment website type							
Gay social networking	354	187	(52.82)	<.001	76	(21.47)	.318
General gay interest	337	219	(64.99)	.909	68	(20.18)	.680
General social networking	5433	3592	(66.11)	REF	825	(15.18)	REF
Geospatial social networking	2075	1509	(72.72)	<.001	570	(27.47)	<.001

^aWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared to a referent (REF) group.

^bWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among HIV-positive participants compared to HIV-negative or unknown serostatus participants. Model controlled for race/ethnicity, age, NHBS residency, and website type.

^cNHBS: National HIV Behavioral Surveillance System.

Table 4. Substance using behaviors of MSM participants in the American Men's Internet Survey, United States, 2014.

Participant characteristics	N in sample	Substance use behaviors in the past 12 months					
		Used marijuana			Used other substance(s)		
		n	(%)	P -value ^a	N	(%)	P -value ^a
HIV positive overall	1049	273	(26.02)	<.001^b	286	(27.26)	<.001^b
Race/ethnicity							
Black non-Hispanic	92	24	(26.09)	.549	19	(20.65)	.110
Hispanic	172	46	(26.74)	.468	44	(25.58)	.571
White non-Hispanic	716	181	(25.28)	REF	202	(28.21)	REF
Other or multiple races	69	22	(31.88)	.359	21	(30.43)	.400
Age (years)							
15-24	68	21	(30.88)	.988	17	(25.00)	.360
25-29	110	40	(36.36)	.082	34	(30.91)	.441
30-39	251	76	(30.28)	.771	91	(36.25)	.011
40 or older	620	136	(21.94)	REF	144	(23.23)	REF
NHBS city resident ^c							
Yes	464	123	(26.51)	.781	130	(28.02)	.573
No	585	150	(25.64)	REF	156	(26.67)	REF
Recruitment website type							
Gay social networking	23	7	(30.43)	.588	4	(17.39)	.316
General gay interest	32	10	(31.25)	.610	11	(34.38)	.255
General social networking	554	119	(21.48)	REF	134	(24.19)	REF
Geospatial social networking	440	137	(31.14)	.971	137	(31.14)	.353
HIV negative or unknown overall	8199	1744	(21.27)	REF	1443	(17.60)	REF
Race/Ethnicity							
Black non-Hispanic	323	60	(18.58)	.051	50	(15.48)	.060
Hispanic	1136	256	(22.54)	.586	212	(18.66)	.596
White non-Hispanic	6103	1274	(20.87)	REF	1051	(17.22)	REF
Other or multiple races	637	154	(24.18)	.123	130	(20.41)	.102
Age (years)							
15-24	1321	424	(32.10)	<.001	269	(20.36)	.160
25-29	1111	303	(27.27)	.005	259	(23.31)	.002
30-39	1711	411	(24.02)	.912	382	(22.33)	.002
40 or older	4056	606	(14.94)	REF	533	(13.14)	REF
NHBS city resident ^c							
Yes	3089	717	(23.21)	<.001	656	(21.24)	<.001
No	5110	1027	(20.10)	REF	787	(15.40)	REF
Recruitment website type							
Gay social networking	354	48	(13.56)	.032	39	(11.02)	.022
General gay interest	337	69	(20.47)	.985	55	(16.32)	.535
General social networking	5433	1095	(20.15)	REF	861	(15.85)	REF
Geospatial social networking	2075	532	(25.64)	<.001	488	(23.52)	<.001

^aWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared with a

referent (REF) group.

^bWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among HIV-positive participants compared to HIV-negative or unknown serostatus participants. Model controlled for race/ethnicity, age, NHBS residency, and website type.

^cNHBS: National HIV Behavioral Surveillance System.

Table 5. HIV testing behaviors of HIV-negative or unknown status MSM participants in the American Men's Internet Survey, United States, 2014.

Participant characteristics	N in sample	HIV testing behaviors					
		HIV tested ever			HIV tested past 12 months		
		n	(%)	<i>P</i> -value ^a	N	(%)	<i>P</i> -value ^a
Race/ethnicity							
Black non-Hispanic	323	287	(88.85)	.503	213	(65.94)	.221
Hispanic	1136	981	(86.36)	.544	706	(62.15)	.290
White non-Hispanic	6103	5299	(86.83)	REF	3454	(56.60)	REF
Other or multiple races	637	558	(87.60)	.469	406	(63.74)	.308
Age (years)							
15-24	1321	892	(67.52)	<.001	699	(52.91)	<.001
25-29	1111	997	(89.74)	.006	742	(66.79)	<.001
30-39	1711	1547	(90.41)	<.001	1037	(60.61)	.498
40 or older	4056	3689	(90.95)	REF	2301	(56.73)	REF
NHBS city resident ^b							
Yes	3089	2787	(90.22)	<.001	1992	(64.49)	<.001
No	5110	4338	(84.89)	REF	2787	(54.54)	REF
Recruitment website type							
Gay social networking	354	280	(79.10)	<.001	172	(48.59)	<.001
General gay interest	337	304	(90.21)	.054	188	(55.79)	.068
General social networking	5433	4662	(85.81)	REF	2881	(53.03)	REF
Geospatial social networking	2075	1879	(90.55)	<.001	1538	(74.12)	<.001
TOTAL	8199	7125	(86.90)		4799	(58.53)	

^aWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared with a referent (REF) group.

^bNHBS: National HIV Behavioral Surveillance System.

Table 6. Sexually transmitted infection testing and diagnosis of MSM participants in the American Men's Internet Survey, United States, 2014.

Participant characteristics	N in sample	STI History in the past 12 months					
		Tested for any STI ^a			Diagnosed with any STI ^a		
		n	(%)	P-value ^b	N	(%)	P-value ^b
HIV positive overall	1049	747	(71.21)	<.001^c	216	(20.59)	<.001^c
Race/ethnicity							
Black non-Hispanic	92	72	(78.26)	.508	25	(27.17)	.905
Hispanic	172	129	(75.00)	.354	47	(27.33)	.687
White non-Hispanic	716	493	(68.85)	REF	120	(16.76)	REF
Other or multiple races	69	53	(76.81)	.867	24	(34.78)	.065
Age (years)							
15-24	68	57	(83.82)	.365	20	(29.41)	.968
25-29	110	94	(85.45)	.086	44	(40.00)	<.001
30-39	251	203	(80.88)	.583	67	(26.69)	.846
40 or older	620	393	(63.39)	REF	85	(13.71)	REF
NHBS city resident ^d							
Yes	464	344	(74.14)	.073	105	(22.63)	.326
No	585	403	(68.89)	REF	111	(18.97)	REF
Recruitment website type							
Gay social networking	23	13	(56.52)	.282	2	(8.70)	.302
General gay interest	32	24	(75.00)	.416	8	(25.00)	.154
General social networking	554	364	(65.70)	REF	81	(14.62)	REF
Geospatial social networking	440	346	(78.64)	.214	125	(28.41)	.246
HIV negative or unknown overall	8199	3086	(37.64)	REF	618	(7.54)	REF
Race/Ethnicity							
Black non-Hispanic	323	156	(48.30)	.272	35	(10.84)	.559
Hispanic	1136	553	(48.68)	.173	147	(12.94)	.007
White non-Hispanic	6103	2086	(34.18)	REF	378	(6.19)	REF
Other or multiple races	637	291	(45.68)	.470	58	(9.11)	.454
Age (years)							
15-24	1321	512	(38.76)	.033	109	(8.25)	.612
25-29	1111	579	(52.12)	<.001	136	(12.24)	.001
30-39	1711	754	(44.07)	.040	174	(10.17)	.021
40 or older	4056	1241	(30.60)	REF	199	(4.91)	REF
NHBS city resident ^d							
Yes	3089	1442	(46.68)	<.001	300	(9.71)	<.001
No	5110	1644	(32.17)	REF	318	(6.22)	REF
Recruitment website type							
Gay social networking	354	84	(23.73)	<.001	14	(3.95)	.173
General gay interest	337	109	(32.34)	.013	17	(5.04)	.107
General social networking	5433	1740	(32.03)	REF	284	(5.23)	REF
Geospatial social networking	2075	1153	(55.57)	<.001	303	(14.60)	<.001

^aSTI: sexually transmitted infection; includes chlamydia, gonorrhea and syphilis.

^bWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared with a referent (REF) group.

^cWald chi-square from multivariable logistic regression comparing behavior (yes versus no) among HIV-positive participants compared with HIV-negative or unknown serostatus participants. Model controlled for race/ethnicity, age, NHBS residency, and website type.

^dNHBS: National HIV Behavioral Surveillance System.

Acknowledgments

The study was funded by a grant from the MAC AIDS Fund and by the National Institutes of Health [P30AI050409] – the Emory Center for AIDS Research.

Conflicts of Interest

Authors Sanchez and Sullivan are members of the Editorial Board of JMIR Public Health and Surveillance. However, they had no involvement in the editorial decision for this manuscript. It was reviewed and handled by an independent editor.

Multimedia Appendix 1

American Men's Internet Survey, 2014.

[[PDF File \(Adobe PDF File\), 141KB-Multimedia Appendix 1](#)]

Reference

1. Sanchez TH, Sineath RC, Kahle EM, Tregear SJ, Sullivan PS. The Annual American Men's Internet survey of behaviors of men who have sex with men in the United States: Protocol and Key Indicators Report 2013. JMIR Public Health Surveill. 2015;1:e3. [doi: [10.2196/publichealth.4314](https://doi.org/10.2196/publichealth.4314)]

Abbreviations

AMIS: American Men's Internet Survey

HIV: human immunodeficiency virus

MSM: men who have sex with men

NHBS: national HIV behavioral surveillance system

STI: sexually transmitted infection

Edited by G Eysenbach; submitted 22.12.15; peer-reviewed by C Khosropour, N Lachowsky, M Gates, J Sewell; comments to author 22.01.16; revised version received 07.03.16; accepted 18.03.16; published 25.05.16

Please cite as:

Sanchez T, Zlotorzynska M, Sineath C, Kahle E, Sullivan P

The Annual American Men's Internet Survey of Behaviors of Men Who have Sex with Men in the United States: 2014 Key Indicators Report

JMIR Public Health Surveill 2016;2(1):e23

URL: <http://publichealth.jmir.org/2016/1/e23/>

doi: [10.2196/publichealth.5476](https://doi.org/10.2196/publichealth.5476)

PMID: [27244770](https://pubmed.ncbi.nlm.nih.gov/27244770/)

©Travis Sanchez, Maria Zlotorzynska, Craig Sineath, Erin Kahle, Patrick Sullivan. Originally published in JMIR Public Health and Surveillance (<http://publichealth.jmir.org>), 25.05.2016. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.