

Protocol

The Prevalence of Sexual Behavior Stigma Affecting Gay Men and Other Men Who Have Sex with Men Across Sub-Saharan Africa and in the United States

Shauna Stahlman¹, MPH, PhD; Travis Howard Sanchez², MPH, DVM; Patrick Sean Sullivan², PhD, DVM; Sosthenes Ketende¹, MSc; Carrie Lyons¹, MPH; Manhattan E Charurat³, PhD; Fatou Maria Drame^{4,5}, PhD; Daouda Diouf⁴, MSc; Rebecca Ezouatchi⁶, MA; Seni Kouanda^{7,8}; Simplicie Anato⁹, MA; Tampose Mothopeng¹⁰; Zandile Mnisi¹¹, BNSc; Stefan David Baral¹, MPH, MD

¹Center for Public Health and Human Rights, Department of Epidemiology, Johns Hopkins University, Baltimore, MD, United States

²Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, United States

³Division of Epidemiology and Prevention, Institute of Human Virology, University of Maryland, Baltimore, MD, United States

⁴Enda Santé, Dakar, Senegal

⁵Department of Geography, School of Social Sciences, Gaston Berger University, Saint-Louis, Senegal

⁶Enda Santé, Abidjan, Côte d'Ivoire

⁷Institut de Recherche en Sciences de la Santé-IRSS, Ouagadougou, Burkina Faso

⁸Institut Africain de Santé Publique, Ouagadougou, Burkina Faso

⁹Arc-en-ciel, Lomé, Togo

¹⁰Matrix Support Group, Maseru, Lesotho

¹¹Ministry of Health, Mbabane, Swaziland

Corresponding Author:

Shauna Stahlman, MPH, PhD
Center for Public Health and Human Rights
Department of Epidemiology
Johns Hopkins University
615 N. Wolfe Street
Baltimore, MD, 21205
United States
Phone: 1 443 287 2370
Fax: 1 410 614 8371
Email: sstahlm1@jhu.edu

Abstract

Background: There has been increased attention for the need to reduce stigma related to sexual behaviors among gay men and other men who have sex with men (MSM) as part of comprehensive human immunodeficiency virus (HIV) prevention and treatment programming. However, most studies focused on measuring and mitigating stigma have been in high-income settings, challenging the ability to characterize the transferability of these findings because of lack of consistent metrics across settings.

Objective: The objective of these analyses is to describe the prevalence of sexual behavior stigma in the United States, and to compare the prevalence of sexual behavior stigma between MSM in Southern and Western Africa and in the United States using consistent metrics.

Methods: The same 13 sexual behavior stigma items were administered in face-to-face interviews to 4285 MSM recruited in multiple studies from 2013 to 2016 from 7 Sub-Saharan African countries and to 2590 MSM from the 2015 American Men's Internet Survey (AMIS), an anonymous Web-based behavioral survey. We limited the study sample to men who reported anal sex with a man at least once in the past 12 months and men who were aged 18 years and older. Unadjusted and adjusted prevalence ratios were used to compare the prevalence of stigma between groups.

Results: Within the United States, prevalence of sexual behavior stigma did not vary substantially by race/ethnicity or geographic region except in a few instances. Feeling afraid to seek health care, avoiding health care, feeling like police refused to protect,

being blackmailed, and being raped were more commonly reported in rural versus urban settings in the United States ($P < .05$ for all). In the United States, West Africa, and Southern Africa, MSM reported verbal harassment as the most common form of stigma. Disclosure of same-sex practices to family members increased prevalence of reported stigma from family members within all geographic settings ($P < .001$ for all). After adjusting for potential confounders and nesting of participants within countries, AMIS-2015 participants reported a higher prevalence of family exclusion ($P = .02$) and poor health care treatment ($P = .009$) as compared with participants in West Africa. However, participants in both West Africa ($P < .001$) and Southern Africa ($P < .001$) reported a higher prevalence of blackmail. The prevalence of all other types of stigma was not found to be statistically significantly different across settings.

Conclusions: The prevalence of sexual behavior stigma among MSM in the United States appears to have a high absolute burden and similar pattern as the same forms of stigma reported by MSM in Sub-Saharan Africa, although results may be influenced by differences in sampling methodology across regions. The disproportionate burden of HIV is consistent among MSM across Sub-Saharan Africa and the United States, suggesting the need in all contexts for stigma mitigation interventions to optimize existing evidence-based and human-rights affirming HIV prevention and treatment interventions.

(*JMIR Public Health Surveill* 2016;2(2):e35) doi: [10.2196/publichealth.5824](https://doi.org/10.2196/publichealth.5824)

KEYWORDS

stigmatization; social stigma; HIV; male homosexuality; United States; Western Africa; Southern Africa; mental health

Introduction

The World Health Organization, UNAIDS, and the White House National HIV/AIDS Strategy have collectively called for the need to reduce stigma toward key populations to reduce the impact of human immunodeficiency virus (HIV) and to optimize HIV treatment outcomes [1-3]. In this context, gay men and other men who have sex with men (MSM) continue to be among the populations at highest risk for HIV infection worldwide [4,5]. The Centers for Disease Control and Prevention recently estimated that if current incidence rates continue, 1 in 6 MSM in the United States will be diagnosed with HIV in their lifetime, including 1 in 2 Black MSM, 1 in 4 Latino MSM, and 1 in 11 White MSM [6]. In low- and middle-income countries, MSM are estimated to have at least 19 times the odds of living with HIV compared with other reproductive-aged adults [7]. And in Sub-Saharan Africa, even in regions commonly characterized as generalized epidemics, the prevalence and incidence of HIV among MSM is consistently higher than that of age-matched men in all settings [8,9].

Sexual behavior stigma, defined here as stigma that is anticipated, perceived, or experienced as a result of one's sexual experience [10], has been linked with adverse HIV-related outcomes among MSM including reduced rates of HIV testing and increased sexual risk practices leading to enhanced risk for HIV infection [11-14]. Separately, sexual behavior stigma has been linked to adverse mental health outcomes such as depression, suicidal ideation, and substance use disorders [15-17]. This stigma can operate at community- or structural-levels to negatively influence health outcomes. At the community level, sexual behavior-related stigma can limit the provision and uptake of sexual health services [18,19]. For example, culturally insensitive health workers may result in MSM avoiding HIV prevention services; or even more problematically, MSM living with HIV may avoid HIV treatment. Reduced use of health and HIV services by MSM, due to enacted or perceived discrimination, may limit knowledge of the risks of unprotected anal intercourse and opportunities for access to prevention services [20,21]. At the structural level,

laws against same-sex practices can increase fear and avoidance of health care services, risk behaviors, stress, and promote violence, which can worsen health conditions for MSM and the broader communities [20,22-25].

Within the United States, HIV disproportionately affects young, Black and Hispanic MSM [26-32]. However, individual behavioral risk factors do not appear to explain the increased risk for HIV among racial/ethnic minorities [30,33]. Instead, the burden of stigma and mental health secondary to sexual and gender minority stress may contribute to the disparities observed around the world [34,35]. As a result, there is a growing recognition for the need to be able to measure and evaluate stigma toward MSM.

However, a consensus is lacking for how to quantify, measure, and evaluate sexual behavior stigma for key populations. Previous studies have identified high levels of sexual behavior stigma among MSM both in Sub-Saharan Africa and in the United States including physical assault and verbal harassment [23,36-39]. However, comparisons are imperfect because many studies do not use consistent metrics for measuring this stigma across settings [40]. Thus, the objectives of this paper are (1) to describe the prevalence of these sexual behavior stigma items in the United States and across age, racial/ethnic, population density, and regional subgroups, and (2) to make a comparison of sexual behavior stigma between MSM in 3 different global regions (United States, West Africa, and Southern Africa) using consistent metrics.

Methods

Study Population and Key Measures

American Men's Internet Survey (AMIS)

The American Men's Internet Survey (AMIS) is an annual cross-sectional behavioral survey of MSM in the United States [41,42]. AMIS-2015 recruited MSM from a variety of websites using banner advertisements or email blasts. Websites included general social networking sites (eg, Facebook), general gay interest sites (eg, sites that post news and articles relevant to a

gay audience), gay social networking sites, and mobile-only geospatial social networking applications that connect men with other men based on their proximity. Approximately one-half of the surveys were performed on desktop computers and one-half on mobile phones. We used an interim survey dataset from September 2015 to March 2016. To be eligible to participate, men had to be aged 15 years or older, identify as the male gender, and report that they had oral or anal sex with a man at least once in the past. For the purpose of making comparisons with the African data sets, we limited the study sample to men who reported anal sex with a man at least once in the past 12 months and men who were aged 18 years and older, because these were eligibility requirements in the African studies. We also limited the AMIS-2015 sample to men with a current US residency as measured using reported zip codes and Internet protocol addresses. No incentive was provided to the participants.

AMIS consisted of a core questionnaire that was administered to all participants and included questions about demographics and disclosure status, which refers to whether the participant disclosed his same sex practices to either health care workers or family members. It also consisted of 3 different subset questionnaires to which participants were randomized at the start of the survey. All AMIS-2015 participants were asked if they identified as Hispanic/Latino, American Indian or Alaskan Native, Asian, Black or African American, Native Hawaiian or Pacific Islander, or White, and could choose all groups that apply. For the purpose of these analyses, we created a single variable for race/ethnicity, which grouped participants into the following mutually exclusive categories: Hispanic, non-Hispanic Black (“Black”), non-Hispanic White (“White”), and non-Hispanic Other (“Other”), which included those who identified as multiracial, Asian, and Native American/Hawaiian. Participants who indicated that they preferred not to answer questions about race, or that they didn’t know/did not apply were treated as missing (39/2551, 1.5%). We used a combination

of county and zip code of residence to determine population density. Population density was classified as urban if participants were recruited from a county or zip code that had a population density of at least 1000 people per square mile, and rural if less than 1000 people per square mile. Sexual behavior stigma items were included in a subset questionnaire; therefore, one-third of participants were asked about sexual behavior stigma (2590/7853, 33.0%).

Sub-Saharan Africa

Data from Sub-Saharan Africa have been presented in previous studies [23,43-45]. In this secondary analysis, we created a combined data set that was limited to MSM who were aged 18 years or older, reported being assigned male sex at birth and also identified as male, and reported anal intercourse with a man in the past 12 months. Data were collected predominantly via respondent driven sampling (RDS) [46] and snowball sampling [47] from 4285 MSM from 7 Sub-Saharan African countries (Table 1). RDS seeds were recruited from local MSM-affiliated community-based organizations (CBOs) although not all seeds were CBO members. Seeds were selected to represent the diversity of the target population with respect to demographics, HIV status, partnering, and other factors. There is significant consistency of methods across settings with face-to-face administration of a structured survey instrument by MSM or MSM-friendly staff members at the local CBOs. In addition, all data sets contain measures of sociodemographics, disclosure status, and sexual behavior stigma. For these analyses, population density was similarly classified as urban if participants were recruited from a study site located in a city that had a population density of at least 1000 people per square mile, and rural if less than 1000 people per square mile. Participants received a modest reimbursement for their time, the cost of travel to the study site (US\$2-\$6), and for each eligible participant they recruited into the study (for RDS studies).

Table 1. Summary of Sub-Saharan Africa MSM data sets.

Region	Country	Laws pertaining to same sex practices [48-50]	Recruitment dates	Study site locations	Recruitment method	Sample size
West Africa	Burkina Faso	Not criminalized	January-August 2013	Bobo Dioulasso and Ouagadougou	RDS ^a	443
West Africa	Cote d'Ivoire	Not criminalized	March-October 2015	Abidjan, Gagnoa, Bouaké, and Yamoussoukro	RDS	794
Southern Africa	Lesotho	Sodomy prohibited as common-law offence	February- September 2014	Maseru and Maputsoe	RDS	487
West Africa	Nigeria	Imprisonment	March 2013- August 2015	Abuja and Lagos	RDS	1067
West Africa	Senegal	Imprisonment	February- November 2015	Dakar, Mbour, and Thies	RDS	522
Southern Africa	Swaziland	Unenforced penalties	October- December 2014	Multiple	Snowball	419
West Africa	Togo	Imprisonment	January-June 2013	Kara and Lome	RDS	553

^aAbb: respondent driven sampling.

Sexual Behavior Stigma

Sexual behavior stigma was measured using the same 13 items in AMIS-2015 and in each of the Sub-Saharan Africa studies (Textbox 1). These measures include perceived, anticipated, and experienced stigma, such as stigma from family and friends,

stigma from health care workers, and stigma from broader society. For questions pertaining to physical attacks and rape, we included responses as a “yes” only if the participant believed that these experiences were related to the fact that he has sex with men.

Textbox 1. Sexual behavior stigma items (response options: yes/no).

1. Have you ever felt excluded from family activities because you have sex with men?
2. Have you ever felt that family members have made discriminatory remarks or gossiped about you because you have sex with men?
3. Have you ever felt rejected by your friends because you have sex with men?
4. Have you ever felt afraid to go to health care services because you worry someone may learn you have sex with men?
5. Have you ever avoided going to health care services because you worry someone may learn you have sex with men?
6. Have you ever felt that you were not treated well in a health center because someone knew that you have sex with men?
7. Have you ever heard health care providers gossiping about you (talking about you) because you have sex with men?
8. Have you ever felt that the police refused to protect you because you have sex with men?
9. Have you ever felt scared to be in public places because you have sex with men?
10. Have you ever been verbally harassed and felt it was because you have sex with men?
11. Have you ever been blackmailed by someone because you have sex with men?
- 12a. Has someone ever physically hurt you (pushed, shoved, slapped, hit, kicked, choked or otherwise physically hurt you)?
- 12b. Do you believe any of these experiences of physical violence was/were related to the fact that you have sex with men?
- 13a. Have you ever been forced to have sex when you did not want to? (By forced, I mean physically forced, coerced to have sex, or penetrated with an object, when you did not want to).
- 13b. Do you believe any of these experiences of sexual violence were related to the fact that you have sex with men?

Ethics

AMIS-2015 data collection was approved by the human subjects research review board at Emory University. Sub-Saharan African studies were approved by respective in-country ethics committees: the Health Research Ethics Committee of Burkina Faso, the Health Research Ethics Committee of Côte d'Ivoire, the Lesotho National Health Research Ethics Committee, the Ethical Committee of Togo, the Senegalese National Health Research Ethics, the Swaziland Scientific Ethics Committee, and the institutional review board (IRB) at the Johns Hopkins Bloomberg School of Public Health. For Nigeria, approval was obtained by the Federal Capital Territory Health Research Ethics Committee, the University of Maryland Baltimore IRB, and the Walter Reed Army Institute of Research IRB. Informed consent was obtained from all individual participants included in these analyses.

Statistical Analysis

We used descriptive statistics (frequencies, percentages) to describe the distribution of sexual behavior stigma items between and within groups. In bivariate analyses, we used log-binomial regression models to generate crude prevalence ratios (PRs) including 95% confidence intervals, which were used to test variables of interest for associations with stigma. Specifically, we examined the association of race/ethnicity, US region, population density, and age group with each sexual behavior stigma item in AMIS-2015. Within US/Africa regions, we measured the association between disclosure and stigma from family members and health care workers.

Multivariable log-binomial models were used to test for associations between region (United States, West Africa, Southern Africa) and prevalence of each stigma item after adjusting for potential confounders (age, disclosure, population density, and education level), which were identified based on findings from previous studies [16,39,43,51,52]. In addition, we accounted for nesting of participants within countries using a random intercept in each of the multivariable models, which is a method that has been used in similar studies [20,53,54]. Listwise deletion was used to handle missing data. Significance was determined at $\alpha = .05$.

Results

Sample Demographics

In AMIS-2015, the median age of participants was 32 years (interquartile range (IQR)=24-50) (Table 2). The sample was roughly evenly divided across regions of the United States, with 19.9% (516/2589) from the Midwest, 18.3% (474/2589) from the Northeast, 37.1% (961/2589) from the South, and 24.6% (638/2589) from the West. Most lived in an urban (1749/2551, 68.6%) area. The distribution of race/ethnicity was as follows: 13.4% (342/2551) Hispanic, 7.4% (188/2551) non-Hispanic Black, 71.7% (1829/2551) non-Hispanic White, and 7.5% (192/2551) Other. The majority (1368/2569, 53.3%) had completed a college level education or higher, 35.0% (898/2569) completed some college, and 11.8% (303/2569) completed high school or lower. Most participants had disclosed their same-sex practices to either a family member (1982/2412, 82.2%) or health care worker (1735/2397, 72.4%).

In West Africa, participants were recruited from Nigeria (1067/3379, 31.6%), Cote d'Ivoire (794/3379, 23.5%), Togo (553/3379, 16.4%), Senegal (522/3379, 15.5%), and Burkina Faso (443/3379, 13.1%). The median age was 23 years (IQR=21-27). Most had completed at least a secondary school education (1898/3359, 56.5%), whereas 24.8% (834/3359) completed more than a secondary school education and 18.7% (627/3359) completed primary school or lower. Overall, 19.7% (665/3379) had disclosed their same-sex practices to a family member and 35.6% (1199/3379) had disclosed to a health care worker.

In Southern Africa, participants were recruited from Lesotho (487/906, 53.8%) and Swaziland (419/906, 46.3%). The median age was 24 years (IQR=21-28). Similar to West Africa, most (597/905, 66.0%) had completed a secondary school education, whereas 20.8% (188/905) completed more than a secondary school education, and 13.3% (120/905) completed primary school or less. Just less than one-third (277/901, 30.7%) disclosed same-sex practices to a family member and 12.2% (110/901) disclosed to a health care worker.

Table 2. Sample demographics by United States/Africa region.

	United States N (%)	Southern Africa N (%)	West Africa N (%)
Median age (IQR)	32 (24-50)	24 (21-28)	23 (21-27)
Education completed			
High school/secondary school or less	303 (11.8)	717 (79.2)	2525 (75.2)
More than secondary/high school	2266 (88.2)	188 (20.8)	834 (24.8)
Population density			
Rural	802 (31.4)	603 (66.6)	597 (17.7)
Urban	1749 (68.6)	303 (33.4)	2782 (82.3)
Disclosed same-sex practices to health care worker	1735 (72.4)	110 (12.2)	1199 (35.6)
Disclosed same-sex practices to family member	1982 (82.2)	277 (30.7)	665 (19.7)

Sexual Behavior Stigma in AMIS-2015

Overall prevalence of stigma items in AMIS-2015 are presented in [Table 3](#). Briefly, the most commonly reported items were verbal harassment (1423/2509, 56.7%), family gossip (1155/2312, 50.0%), and being scared to be in public (811/2551, 31.8%). Prevalence of sexual behavior stigma by race/ethnicity within the United States are shown in [Figure 1](#) and [Multimedia Appendix 1](#). In the bivariate analyses, Hispanic MSM were more likely to report being blackmailed as compared with White MSM ($P=.002$). Black MSM were less likely than White MSM to report family exclusion ($P=.04$), fear of being in public ($P<.001$), verbal harassment ($P<.001$), and physical attacks ($P=.002$).

There were no differences in sexual behavior stigma prevalence by United States region, except that MSM in the West were less likely to feel like police refused to protect them as compared with MSM in the South ($P=.04$), and MSM in the Midwest

($P<.001$), and Northeast ($P=.02$) were less likely to report blackmail as compared with MSM in the South ([Figure 2](#) and [Multimedia Appendix 2](#)). MSM who lived in rural as compared with urban areas were more likely to report being afraid to seek health care ($P<.001$), avoiding health care ($P=.003$), feeling like police refused to protect ($P=.001$), being blackmailed ($P=.004$), and being raped ($P=.04$) ([Figure 3](#) and [Multimedia Appendix 3](#)).

Finally, young MSM (aged 18-24 years) were more likely than older MSM to be excluded by family members ($P=.002$), gossiped about by family members ($P=.007$), afraid to seek health care ($P<.001$), avoid seeking health care ($P=.002$), scared to be seen in public ($P=.03$), and to be blackmailed ($P=.003$) ([Figure 4](#) and [Multimedia Appendix 4](#)). However, older MSM (aged 25 years and older) were more likely to have been treated poorly in a health care center ($P<.001$), and to feel like police refused to protect ($P=.03$).





Figure 1. Prevalence of sexual behavior stigma among MSM in AMIS-2015 by race/ethnicity.**Figure 2.** Prevalence of sexual behavior stigma among MSM in AMIS-2015 by United States region.

Figure 3. Prevalence of sexual behavior stigma among MSM in AMIS-2015 by population density.**Figure 4.** Prevalence of sexual behavior stigma among MSM in AMIS-2015 by age group.

Comparison of Stigma Prevalence Across United States and Sub-Saharan Africa

Overall, sexual behavior stigma reported in AMIS-2015 was similar to or higher than what was reported by participants in the Sub-Saharan African studies (Figure 5 and Table 3). Family exclusion, family gossip, friend rejection, being afraid to seek health care, being treated poorly by a health care worker, feeling like police refused to protect, being scared to be in public, verbal harassment, and being physically hurt were all more commonly reported by AMIS-2015 participants ($P<.001$ for all, except

$P=.005$ for comparing fear of seeking health care with Southern Africa). AMIS-2015 participants were also more likely than MSM in West Africa to report avoiding seeking health care ($P=.002$) and health care worker gossip ($P<.001$). Participants in West and Southern Africa more commonly reported blackmail ($P<.001$ for both) compared with those in the United States, and participants in West Africa more commonly reported rape as compared with MSM in the United States ($P<.001$). In all settings, MSM reported verbal harassment as the most common experience of stigma. Rape was the least common experience in United States and Southern Africa, whereas being treated

poorly in a health care center was least commonly reported in West Africa.

Table 3. Prevalence of sexual behavior stigma among MSM by United States/Africa region.

Stigma	Region	n/N (%)	Prevalence ratio (95% confidence interval)	P value
Family exclusion	United States	775/2489 (31.1)	Reference	--
	Southern Africa	126/905 (13.9)	0.45 (0.38-0.53)	<.001
	West Africa	298/3378 (8.8)	0.28 (0.25-0.32)	<.001
Family gossip	United States	1155/2312 (50.0)	Reference	--
	Southern Africa	181/903 (20.0)	0.40 (0.35-0.46)	<.001
	West Africa	727/3377 (21.5)	0.43 (0.40-0.47)	<.001
Friend rejection	United States	677/2400 (28.2)	Reference	--
	Southern Africa	173/902 (19.2)	0.68 (0.59-0.79)	<.001
	West Africa	604/3378 (17.9)	0.63 (0.58-0.70)	<.001
Afraid to seek health care	United States	665/2429 (27.4)	Reference	--
	Southern Africa	203/905 (22.4)	0.82 (0.71-0.94)	.005
	West Africa	756/3378 (22.4)	0.82 (0.75-0.89)	<.001
Poor healthcare treatment	United States	462/2374 (19.5)	Reference	--
	Southern Africa	62/905 (6.9)	0.35 (0.27-0.45)	<.001
	West Africa	102/3355 (3.0)	0.16 (0.13-0.19)	<.001
Avoided health care	United States	488/2430 (20.1)	Reference	--
	Southern Africa	180/905 (19.9)	0.99 (0.85-1.15)	.90
	West Africa	573/3377 (17.0)	0.84 (0.76-0.94)	.002
Health care worker gossip	United States	201/2397 (8.4)	Reference	--
	Southern Africa	74/905 (8.2)	0.98 (0.76-1.26)	.85
	West Africa	195/3350 (5.8)	0.69 (0.57-0.84)	<.001
Police refused to protect	United States	298/2367 (12.6)	Reference	--
	Southern Africa	65/902 (7.2)	0.57 (0.44-0.74)	<.001
	West Africa	244/3374 (7.2)	0.57 (0.49-0.67)	<.001
Scared to be in public	United States	811/2551 (31.8)	Reference	--
	Southern Africa	208/905 (23.0)	0.72 (0.63-0.83)	<.001
	West Africa	452/3376 (13.4)	0.42 (0.38-0.47)	<.001
Verbally harassed	United States	1423/2509 (56.7)	Reference	--
	Southern Africa	351/905 (38.8)	0.68 (0.63-0.75)	<.001
	West Africa	934/3377 (27.7)	0.49 (0.46-0.52)	<.001
Blackmailed	United States	252/2507 (10.1)	Reference	--
	Southern Africa	178/905 (19.7)	1.96 (1.64-2.33)	<.001
	West Africa	686/3378 (20.3)	2.02 (1.77-2.31)	<.001
Physically hurt	United States	473/2513 (18.8)	Reference	--
	Southern Africa	120/903 (13.3)	0.71 (0.59-0.85)	<.001
	West Africa	388/3356 (11.6)	0.61 (0.54-0.70)	<.001
Raped	United States	159/2377 (6.7)	Reference	--
	Southern Africa	59/902 (6.5)	0.98 (0.73-1.31)	.88
	West Africa	388/3365 (11.5)	1.72 (1.44-2.06)	<.001

Sexual behavior stigma from family and friends was higher among those who had disclosed their sexual practices to a family member in all regions ($P<.001$ for all); however, the prevalence of friend rejection among AMIS-2015 participants was similar between the two disclosure groups ($P=.75$) (Figure 6 and Multimedia Appendix 5).

In the United States, participants who had disclosed their sexual practices to a health care worker were less likely to be afraid to

seek health care ($P<.001$) or to avoid seeking health care ($P<.001$) as compared with those who had not disclosed; however, they were more likely than those who had not disclosed to report being treated poorly in a health care center ($P<.001$) (Figure 7 and Multimedia Appendix 6). In both West and Southern Africa, the prevalence of each health care-related sexual behavior stigma item increased among those who had disclosed versus not disclosed ($P<.001$ for all).

Figure 5. Prevalence of sexual behavior stigma among MSM by United States/Africa region.




Figure 6. Prevalence of sexual behavior stigma among MSM who disclosed same-sex behaviors to family vs not disclosed.





Figure 7. Prevalence of sexual behavior stigma among MSM who disclosed same-sex behaviors to health care worker vs not disclosed.

Multivariable Adjusted Associations of United States/Africa Region With Sexual Behavior Stigma Items

After adjusting for age, disclosure status with family members and health care workers, education level, and population density, we found that AMIS-2015 participants continued to report

higher levels of family exclusion ($P=.02$) and poor health care treatment ($P=.009$) as compared with West Africa (Table 4). However, participants in both West Africa ($P<.001$) and Southern Africa ($P<.001$) reported a higher prevalence of blackmail. The prevalence of all other types of stigma was not found to be statistically significantly different across settings.

Table 4. Adjusted associations of United States/Africa region with sexual behavior stigma.

Stigma	Region	Adjusted prevalence ratio ^a (95% CI)	P value
Family exclusion	United States	Reference	--
	Southern Africa	0.52 (0.19-1.44)	.21
	West Africa	0.33 (0.13-0.82)	.02
Family gossip	United States	Reference	--
	Southern Africa	0.52 (0.20-1.35)	.18
	West Africa	0.50 (0.22-1.17)	.11
Friend rejection	United States	Reference	--
	Southern Africa	0.91 (0.42-1.97)	.81
	West Africa	0.75 (0.37-1.49)	.41
Afraid to seek health care	United States	Reference	--
	Southern Africa	0.79 (0.20-3.18)	.74
	West Africa	0.77 (0.22-2.67)	.68
Poor health care treatment	United States	Reference	--
	Southern Africa	0.52 (0.14-1.95)	.33
	West Africa	0.21 (0.06-0.67)	.009
Avoided health care	United States	Reference	--
	Southern Africa	0.91 (0.20-4.17)	.09
	West Africa	0.85 (0.22-3.32)	.82
Health care worker gossip	United States	Reference	--
	Southern Africa	1.25 (0.32-4.96)	.75
	West Africa	0.97 (0.29-3.30)	.97
Police refused to protect	United States	Reference	--
	Southern Africa	0.84 (0.08-8.38)	.88
	West Africa	0.59 (0.08-4.59)	.61
Scared to be in public	United States	Reference	--
	Southern Africa	0.71 (0.15-3.28)	.66
	West Africa	0.37 (0.09-1.45)	.15
Verbally harassed	United States	Reference	--
	Southern Africa	0.84 (0.44-1.59)	.59
	West Africa	0.52 (0.30-0.93)	.03
Blackmailed	United States	Reference	--
	Southern Africa	2.94 (1.70-5.07)	<.001
	West Africa	2.86 (1.76-4.64)	<.001
Physically hurt	United States	Reference	--
	Southern Africa	0.83 (0.17-4.07)	.92
	West Africa	0.56 (0.14-2.31)	.43
Raped	United States	Reference	--
	Southern Africa	0.94 (0.17-5.15)	.94
	West Africa	1.68 (0.37-7.62)	.50

^aModels adjust for age, disclosure of same-sex behaviors to family or health care workers, education level, and population density. Models include a random intercept to account for nesting of participants within countries.

Discussion

Principal Results

In these analyses, we identified a high prevalence of experienced, anticipated, and perceived sexual behavior stigma for MSM in all settings. Based on [Figure 5](#), the overall pattern of responses appeared to be similar across regions, with MSM in all settings most commonly reporting verbal harassment and less commonly reporting rape or being treated poorly in a health care center. Surprisingly, in many cases US participants reported even greater levels of stigma than MSM in settings where same sex practices were criminalized. However, many of these differences were not significant after adjusting for potential confounders, and may also be the result of inherent differences in sampling methodology (eg, Web-based questionnaire vs face-to-face interviews).

Interpretation and Comparison with Prior Work

Traumatic experiences among MSM are problematic for the HIV response because they can lead to reduced uptake and use of health services [18,20,21,23,55]. Indeed, stigma has been associated with reduced rates of HIV testing, increased risk for HIV infection, increased fear and avoidance of health care, increased condomless anal sex, and reduced engagement in HIV treatment for those living with HIV [11-14,23]. Stigma has further been linked to adverse mental health outcomes such as reduced self-esteem, internalized homophobia, and depression, with potential mediation by resiliency and coping ability [56-59]. Thus, the findings of high prevalence of multiple forms of stigma suggest the potential for stigma mitigation interventions to improve existing mental health services and HIV prevention and treatment interventions for MSM.

Within AMIS-2015, there were minimal differences in report of sexual behavior stigma by race/ethnicity and by region, suggesting the pervasiveness of these experiences or perceptions among MSM in the United States. However, MSM in rural areas were more likely to report fear and avoidance of health care, feeling like police refused to protect, being blackmailed, and being raped. This is consistent with the findings from previous studies suggesting that stigma toward accessing mental health and HIV services is higher in rural regions of both the United States and Sub-Saharan Africa as compared with more urban regions [51,60,61]. Our finding that Black MSM experienced lower levels of certain types of sexual behavior stigma is in contrast to previous literature suggesting that stigma surrounding sexual orientation is more pervasive in the Black community [62,63]. However, much of this previous work assessed internalized homo-negativity among Black MSM, which we did not measure in this study [64]. In addition, there is potential selection bias in that Black MSM are underrepresented in AMIS-2015, and thus this finding should be further explored and confirmed using alternative sampling methods. There were some differences within AMIS-2015 between age groups, with young MSM (aged 18-24 years) being more likely than older MSM to report stigma from family, to be afraid to seek health care or avoid seeking health care, to be scared to be seen in public, and to be blackmailed. This is somewhat surprising given that we measured lifetime exposure to stigma and

suspected that older MSM would have had more opportunity for exposure, and thus higher levels of stigma. These high levels of anticipated stigma and particularly fear and avoidance of health care among young MSM are important to address, as they likely contribute to the sustained or growing incidence of HIV among young MSM worldwide [65-69].

When we stratified by disclosure status, the prevalence of stigma from family members increased for participants who had disclosed same-sex practices to family as compared with those who had not disclosed, and this occurred within each setting of the United States, West Africa, and Southern Africa. This is likely reflective of the fact that in all of these settings MSM who have disclosed their same sex practices are more easily identified as targets for stigma, discrimination, and harassment [43,70-72]. However, the patterns of stigma associated with disclosure status were somewhat different when we examined health care-related stigma. In the United States, those who had disclosed same-sex practices to a health care worker were less likely to fear or avoid seeking health care services, although they were more likely to report being treated poorly in a health care center. Based on these findings, it seems possible that MSM who have disclosed to their providers are more comfortable seeking health care, even though it can sometimes result in negative experiences. In Sub-Saharan Africa, health care stigma increased among those who had disclosed versus not disclosed, indicating the immediate need for structural interventions to improve access to culturally competent health care for MSM across all settings [1,22,73].

There have been successful efforts in Sub-Saharan Africa and other regions to increase clinical and cultural competency for health care workers who provide HIV and sexually transmitted infection prevention, treatment, and care to MSM patients [74-78]. Given our findings, these efforts should be intensified and expanded to cover all domains of health care, particularly because the proper training of health care professionals tends to be one of the more easily implementable intervention strategies to reduce or mitigate stigma. However, stigma toward MSM is not limited to health care settings and structural interventions may also be needed to reduce stigma in community settings including schools, workplaces, churches, and families. These interventions would need to be appropriately tailored to meet the needs of different cultures and communities. In the United States, for example, gay-straight alliances have been successful in reducing homophobia in schools, although this method may or may not be suitable for African settings [79]. Eventual acceptance of lesbian, gay, bisexual, and transgender individuals may be inevitable with slowly increasing social acceptance over time; however, these social changes are not happening at the pace required to make an immediate impact on reducing negative health outcomes.

Limitations

Although we restricted the study populations from each data set to be as similar as possible (eg, aged 18 years and older, anal sex with a man in the past 12 months, cis-gender male) and performed adjusted analyses, there are some inherent differences between the United States and Sub-Saharan Africa study populations that cannot be adjusted for, including sampling

strategies, mode of survey administration, and time period of study participation. Because AMIS-2015 was an anonymous Web-based survey, it is possible some participants were more comfortable to disclose this sensitive information in an anonymous Internet setting as compared to during a face-to-face interview [80,81]. This bias might be mitigated because interviewers were administered by highly trained MSM or MSM-friendly staff members. Moreover, participants in these studies reported high levels of another sensitive issue, condomless anal sex, which further suggests minimal bias due to social desirability (data not shown).

Another limitation is that the African studies were conducted with incentives for participation that were not included in AMIS, including in some cases HIV and biological testing, and therefore may have reached individuals who would have not otherwise participated. Participants in AMIS were highly educated and mostly of White, non-Hispanic race/ethnicity, suggesting limitations to generalizability to other MSM in the United States. It is also possible that AMIS participants, who were recruited from gay-related websites, were more actively involved in the gay community and more likely to be exposed to stigma. This is because participants in Sub-Saharan Africa were recruited using RDS (except in Swaziland), which has been shown to be capable of reaching MSM who are less engaged in the gay community and HIV testing [82]. Finally, the prevalence of missing values for AMIS-2015 (1.5%-10.7%

for each stigma item) were higher than those in Western and Southern Africa (0%-1%), likely because AMIS-2015 was a Web-based survey, whereas African data were collected during face-to-face interviews. However, this study is the first to our knowledge to make a direct comparison of sexual behavior stigma between MSM in the United States and Sub-Saharan Africa using consistent metrics.

Conclusions

Sexual behavior stigma felt or experienced by MSM appears to be pervasive across the United States and Sub-Saharan Africa, and this has implications for both sexual and mental health. Given the growing desire to measure, quantify, and evaluate stigma toward key populations as part of the HIV response, this paper is timely in its comparison of prevalence of sexual behavior stigma across 3 widely different regions (the United States and Western and Southern Sub-Saharan Africa). It is important to note that results may be influenced by differences in sampling methodology across regions. However, the consistently observed high burden of stigma points to the need for immediate structural interventions to address each of the domains of sexual behavior stigma presented here; that is, stigma from family and friends, health care workers, and from broader society. These interventions will be critical for making a positive impact on the mental health of these men, and also for reducing the global sustained burden of HIV and other adverse health outcomes in this key population.

Acknowledgments

AMIS-2015 was funded by a grant from the MAC AIDS Fund and by the National Institutes of Health (NIH) [P30AI050409] – the Emory Center for AIDS Research. Work in Togo and Burkina Faso was supported by Project SEARCH, which was funded by the US Agency for International Development under Contract GHH-I-00-07-00032-00 and by the President's Emergency Plan for AIDS Relief (PEPFAR). Work in Côte d'Ivoire was funded by the Global Fund to Fight AIDS, Tuberculosis and Malaria through the Government of Côte d'Ivoire National AIDS Control Program (PNPEC) contract to Enda Santé, an organization based in Senegal, and subcontracted for technical assistance to Johns Hopkins University. Work in Lesotho was funded by the US Agency for International Development (USAID, AID-674-A-00-00001), and implemented by Population Services International/Lesotho (PSI). Work in Nigeria was supported by a cooperative agreement (W81XWH-11-2-0174) between the Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc., and the US Department of Defense (DOD). The Nigeria study is also supported by funds from the US NIH under Award No. R01MH099001-01, the US Military HIV Research Program (Grant No. W81XWH-07-2-0067), Fogarty AITRP (D43TW01041), and PEPFAR through cooperative agreement U2G IPS000651 from the United States Department of Health and Human Services (HHS)/Centers for Disease Control and Prevention (CDC), Global AIDS Program with the Institute of Human Virology Nigeria (IHVN). Work in Senegal was funded by USAID (AID-OAA-A-13-00089) and work in Swaziland was funded by PEPFAR through the USAID Swaziland (GHH-I-00-07-00032-00). Finally, this publication was made possible with help from the Johns Hopkins University Center for AIDS Research, an NIH funded program (P30AI094189), which is supported by the following NIH Cofunding and Participating Institutes and Centers: National Institute of Allergy and Infectious Diseases (NIAID), National Cancer Institute (NCI), National Institute of Child Health and Human Development (NICHD), National Heart, Lung, and Blood Institute (NHLBI), National Institute on Drug Abuse (NIDA), National Institute of Mental Health (NIMH), National Institute on Aging (NIA), Fogarty International Center (FIC), National Institute of General Medical Sciences (NIGMS), National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), and the Office of AIDS Research (OAR). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The funding sources had no role in the study design; in the collection, analysis and interpretation of data; in the writing of the articles; or in the decision to submit the manuscript for publication.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Prevalence of sexual behavior stigma among MSM in AMIS-2015 by race/ethnicity.

[[PDF File \(Adobe PDF File\), 23KB-Multimedia Appendix 1](#)]

Multimedia Appendix 2

Prevalence of sexual behavior stigma among MSM in AMIS-2015 by United States region.

[[PDF File \(Adobe PDF File\), 23KB-Multimedia Appendix 2](#)]

Multimedia Appendix 3

Prevalence of sexual behavior stigma among MSM in AMIS-2015 by population density.

[[PDF File \(Adobe PDF File\), 22KB-Multimedia Appendix 3](#)]

Multimedia Appendix 4

Prevalence of sexual behavior stigma among MSM in AMIS-2015 by age group.

[[PDF File \(Adobe PDF File\), 22KB-Multimedia Appendix 4](#)]

Multimedia Appendix 5

Prevalence of sexual behavior stigma among MSM who disclosed same-sex behaviors to family vs. not disclosed, by United States/Africa region.

[[PDF File \(Adobe PDF File\), 21KB-Multimedia Appendix 5](#)]

Multimedia Appendix 6

Prevalence of sexual behavior stigma among MSM who disclosed same-sex behaviors to healthcare worker vs. not disclosed, by United States/Africa region.

[[PDF File \(Adobe PDF File\), 21KB-Multimedia Appendix 6](#)]

References

1. World Health Organization. Prevention and Treatment of HIV and Other Sexually Transmitted Infections Among Men Who Have Sex With Men and Transgender People: Recommendations for a Public Health Approach. 2011. URL: http://www.who.int/hiv/pub/guidelines/msm_guidelines2011/en/ [accessed 2016-07-11] [WebCite Cache ID 6ivSmgy4m]
2. Joint United Nations Programme on HIV/AIDS (UNAIDS). Action on Social Drivers to End AIDS and Extreme Poverty: An Expert Consultation on HIV/AIDS and the Global Health and Development Agendas. Washington, D.C. World Bank Main Complex; Jan 10, 2014. URL: https://www.health.ny.gov/diseases/aids/ending_the_epidemic/docs/key_resources/housing_and_supportive_services/social_drivers_to_end_aids.pdf [accessed 2016-07-11] [WebCite Cache ID 6ivT9FBxP]
3. White House Office of National AIDS Policy. National HIV/AIDS Strategy for the United States: Updated to 2020. 2015. URL: <https://www.aids.gov/federal-resources/national-hiv-aids-strategy/overview/> [accessed 2016-07-11] [WebCite Cache ID 6ivTNgZjT]
4. Beyrer C, Baral SD, van GF, Goodreau SM, Chariyalertsak S, Wirtz AL, et al. Global epidemiology of HIV infection in men who have sex with men. *Lancet*. 2012;380:367-377. [FREE Full text] [doi: [10.1016/S0140-6736\(12\)60821-6](https://doi.org/10.1016/S0140-6736(12)60821-6)] [Medline: [22819660](https://pubmed.ncbi.nlm.nih.gov/22819660/)]
5. Beyrer C, Sullivan P, Sanchez J, Baral SD, Collins C, Wirtz AL, et al. The increase in global HIV epidemics in MSM. *AIDS*. 2013;27:2665-2678. [doi: [10.1097/01.aids.0000432449.30239.fe](https://doi.org/10.1097/01.aids.0000432449.30239.fe)] [Medline: [23842129](https://pubmed.ncbi.nlm.nih.gov/23842129/)]
6. Hess K, Hu X, Lansky A, Mermin J, Hall H. Estimating the lifetime risk of a diagnosis of HIV infection in the United States. 2016. Presented at: Conference on Retroviruses and Opportunistic Infections (CROI); February 23, 2016; Boston, MA. URL: <http://www.croiconference.org/sessions/estimating-lifetime-risk-diagnosis-hiv-infection-united-states>
7. Baral S, Sifakis F, Cleghorn F, Beyrer C. Elevated risk for HIV infection among men who have sex with men in low- and middle-income countries 2000-2006: a systematic review. *PLoS Med*. 2007;4:e339. [FREE Full text] [doi: [10.1371/journal.pmed.0040339](https://doi.org/10.1371/journal.pmed.0040339)] [Medline: [18052602](https://pubmed.ncbi.nlm.nih.gov/18052602/)]
8. Baral S. HIV risks and vulnerabilities among men who have sex with men across Sub-Saharan Africa. 2015. Presented at: Conference on Retroviruses and Opportunistic Infections (CROI); 2015; Seattle, Washington.
9. van Griensven F, de Lind van Wijngaarden JW, Baral S, Grulich A. The global epidemic of HIV infection among men who have sex with men. *Curr Opin HIV AIDS*. 2009;4:300-307. [doi: [10.1097/COH.0b013e32832c3bb3](https://doi.org/10.1097/COH.0b013e32832c3bb3)] [Medline: [19532068](https://pubmed.ncbi.nlm.nih.gov/19532068/)]

10. Brown L, Macintyre K, Trujillo L. Interventions to reduce HIV/AIDS stigma: what have we learned? *AIDS Educ Prev*. 2003;15:49-69. [Medline: [12627743](#)]
11. Knox J, Sandfort T, Yi H, Reddy V, Maimane S. Social vulnerability and HIV testing among South African men who have sex with men. *Int J STD AIDS*. 2011;22:709-713. [FREE Full text] [doi: [10.1258/ijsa.2011.010350](#)] [Medline: [22174050](#)]
12. Hladik W, Barker J, Ssenkusu JM, Opio A, Tappero JW, Hakim A, et al. Crane Survey Group. HIV infection among men who have sex with men in Kampala, Uganda--a respondent driven sampling survey. *PLoS One*. 2012;7:e38143. [FREE Full text] [doi: [10.1371/journal.pone.0038143](#)] [Medline: [22693590](#)]
13. Hu Y, Lu H, Raymond HF, Sun Y, Sun J, Jia Y, et al. Measures of condom and safer sex social norms and stigma towards HIV/AIDS among Beijing MSM. *AIDS Behav*. 2014;18:1068-1074. [FREE Full text] [doi: [10.1007/s10461-013-0609-7](#)] [Medline: [24057931](#)]
14. Choi K, Hudes ES, Steward WT. Social discrimination, concurrent sexual partnerships, and HIV risk among men who have sex with men in Shanghai, China. *AIDS Behav*. 2008;12(Suppl):S71-S77. [doi: [10.1007/s10461-008-9394-0](#)] [Medline: [18427972](#)]
15. Cochran SD, Mays VM, Sullivan JG. Prevalence of mental disorders, psychological distress, and mental health services use among lesbian, gay, and bisexual adults in the United States. *J Consult Clin Psychol*. 2003;71:53-61. [FREE Full text] [Medline: [12602425](#)]
16. Stahlman S, Grosso A, Ketende S, Sweitzer S, Mothopeng T, Taruberekera N, et al. Depression and social stigma among MSM in Lesotho: implications for HIV and sexually transmitted infection prevention. *AIDS Behav*. 2015;19:1460-1469. [doi: [10.1007/s10461-015-1094-y](#)] [Medline: [25969182](#)]
17. Stoloff K, Joska JA, Feast D, De SG, Hugo J, Struthers H, et al. A description of common mental disorders in men who have sex with men (MSM) referred for assessment and intervention at an MSM clinic in Cape Town, South Africa. *AIDS Behav*. 2013;Suppl 1:S77-S81. [doi: [10.1007/s10461-013-0430-3](#)] [Medline: [23532397](#)]
18. Baral SD, Ketende S, Mnisi Z, Mabuza X, Grosso A, Sithole B, et al. A cross-sectional assessment of the burden of HIV and associated individual- and structural-level characteristics among men who have sex with men in Swaziland. *J Int AIDS Soc*. 2013;Suppl 3:18768. [FREE Full text] [Medline: [24321117](#)]
19. The Foundation for AIDS Research (amfar), Johns Hopkins Bloomberg School of Public Health. Achieving an AIDS-Free Generation for Gay Men and Other MSM: Financing Implementation of HIV Programs Targeting MSM. 2012. URL: [http://www.amfar.org/uploadedFiles/amfar.org/In The Community/Publications/MSM-GlobalRept2012.pdf](http://www.amfar.org/uploadedFiles/amfar.org/In%20The%20Community/Publications/MSM-GlobalRept2012.pdf) [accessed 2016-07-11] [WebCite Cache ID 6ivXfnfkI]
20. Pachankis JE, Hatzenbuehler ML, Hickson F, Weatherburn P, Berg RC, Marcus U, et al. Hidden from health: structural stigma, sexual orientation concealment, and HIV across 38 countries in the European MSM Internet Survey. *AIDS*. 2015;29:1239-1246. [doi: [10.1097/QAD.0000000000000724](#)] [Medline: [26035323](#)]
21. Ayala G, Makofane K, Santos G, Beck J, Do TD, Hebert P, et al. Access to basic HIV-related services and PrEP acceptability among men who have sex with men worldwide: barriers, facilitators, and implications for combination prevention. *J Sex Transm Dis*. 2013;2013:953123. [FREE Full text] [doi: [10.1155/2013/953123](#)] [Medline: [26316968](#)]
22. Baral S, Holland CE, Shannon K, Logie C, Semugoma P, Sithole B, et al. Enhancing benefits or increasing harms: community responses for HIV among men who have sex with men, transgender women, female sex workers, and people who inject drugs. *J Acquir Immune Defic Syndr*. 2014;Suppl 3:S319-S328. [doi: [10.1097/QAI.0000000000000233](#)] [Medline: [25007203](#)]
23. Schwartz S, Nowak R, Orazulike I, Keshinro B, Ake J, Kennedy S, et al. The immediate effect of the Same-Sex Marriage Prohibition Act on stigma, discrimination, and engagement on HIV prevention and treatment services in men who have sex with men in Nigeria: analysis of prospective data from the TRUST cohort. *Lancet HIV*. 2015;2:e299-e306. [FREE Full text] [doi: [10.1016/S2352-3018\(15\)00078-8](#)] [Medline: [26125047](#)]
24. Oldenburg CE, Perez-Brumer AG, Hatzenbuehler ML, Krakower D, Novak DS, Mimiaga MJ, et al. State-level structural sexual stigma and HIV prevention in a national online sample of HIV-uninfected MSM in the United States. *AIDS*. 2015;29:837-845. [FREE Full text] [doi: [10.1097/QAD.0000000000000622](#)] [Medline: [25730508](#)]
25. Hatzenbuehler ML, McLaughlin KA. Structural stigma and hypothalamic-pituitary-adrenocortical axis reactivity in lesbian, gay, and bisexual young adults. *Ann Behav Med*. 2014;47:39-47. [FREE Full text] [doi: [10.1007/s12160-013-9556-9](#)] [Medline: [24154988](#)]
26. Centers for Disease Control and Prevention (CDC). HIV infections attributed to male-to-male sexual contact - metropolitan statistical areas, United States and Puerto Rico, 2010. *MMWR Morb Mortal Wkly Rep*. 2012;61:962-966. [FREE Full text] [Medline: [23190569](#)]
27. Prejean J, Tang T, Hall HI. HIV diagnoses and prevalence in the southern region of the United States, 2007-2010. *J Community Health*. 2013;38:414-426. [doi: [10.1007/s10900-012-9633-1](#)] [Medline: [23179388](#)]
28. Stall R, Duran L, Wisniewski SR, Friedman MS, Marshal MP, McFarland W, et al. Running in place: implications of HIV incidence estimates among urban men who have sex with men in the United States and other industrialized countries. *AIDS Behav*. 2009;13:615-629. [FREE Full text] [doi: [10.1007/s10461-008-9509-7](#)] [Medline: [19205867](#)]
29. Millett GA, Peterson JL, Wolitski RJ, Stall R. Greater risk for HIV infection of black men who have sex with men: a critical literature review. *Am J Public Health*. 2006;96:1007-1019. [doi: [10.2105/AJPH.2005.066720](#)] [Medline: [16670223](#)]

30. Millett GA, Flores SA, Peterson JL, Bakeman R. Explaining disparities in HIV infection among black and white men who have sex with men: a meta-analysis of HIV risk behaviors. *AIDS*. 2007;21:2083-2091. [doi: [10.1097/QAD.0b013e3282e9a64b](https://doi.org/10.1097/QAD.0b013e3282e9a64b)] [Medline: [17885299](#)]
31. Centers for Disease Control and Prevention (CDC). HIV Incidence. 2010. URL: <http://www.cdc.gov/hiv/statistics/surveillance/incidence.html> [accessed 2016-01-20] [WebCite Cache ID 6ivYI0IOz]
32. Matthews DD, Herrick AL, Coulter RWS, Friedman MR, Mills TC, Eaton LA, et al. Running backwards: consequences of current HIV incidence rates for the next generation of Black MSM in the United States. *AIDS Behav*. 2016;20:7-16. [doi: [10.1007/s10461-015-1158-z](https://doi.org/10.1007/s10461-015-1158-z)] [Medline: [26267251](#)]
33. Millett GA, Peterson JL, Flores SA, Hart TA, Jeffries WL, Wilson PA, et al. Comparisons of disparities and risks of HIV infection in black and other men who have sex with men in Canada, UK, and USA: a meta-analysis. *Lancet*. 2012;380:341-348. [doi: [10.1016/S0140-6736\(12\)60899-X](https://doi.org/10.1016/S0140-6736(12)60899-X)] [Medline: [22819656](#)]
34. Poteat T, Reinsner SL, Radix A. HIV epidemics among transgender women. *Curr Opin HIV AIDS*. 2014;9:168-173. [doi: [10.1097/COH.000000000000030](https://doi.org/10.1097/COH.000000000000030)] [Medline: [24322537](#)]
35. De Santis JP. HIV infection risk factors among male-to-female transgender persons: a review of the literature. *J Assoc Nurses AIDS Care*. 2009;20:362-372. [doi: [10.1016/j.jana.2009.06.005](https://doi.org/10.1016/j.jana.2009.06.005)] [Medline: [19732695](#)]
36. Balaji A, Smith J, Bowles K, Paz-Bailey G, NHBS study group. Association between enacted stigma and HIV related risk behavior among MSM, National HIV Behavioral Surveillance System. 2011. Presented at: Conference on Retroviruses and Opportunistic Infections (CROI); February 23-26, 2015; Seattle, Washington. URL: <http://www.croiconference.org/sites/default/files/posters-2015/1058.pdf>
37. Cange CW, LeBreton M, Billong S, Saylors K, Tamoufe U, Papworth E, et al. Influence of stigma and homophobia on mental health and on the uptake of HIV/sexually transmissible infection services for Cameroonian men who have sex with men. *Sex Health*. 2015. [doi: [10.1071/SH15001](https://doi.org/10.1071/SH15001)] [Medline: [26117222](#)]
38. Fay H, Baral SD, Trapence G, Motimedi F, Umar E, Ipinge S, et al. Stigma, health care access, and HIV knowledge among men who have sex with men in Malawi, Namibia, and Botswana. *AIDS Behav*. 2011;15:1088-1097. [doi: [10.1007/s10461-010-9861-2](https://doi.org/10.1007/s10461-010-9861-2)] [Medline: [21153432](#)]
39. Risher K, Adams D, Sithole B, Ketende S, Kennedy C, Mnisi Z, et al. Sexual stigma and discrimination as barriers to seeking appropriate healthcare among men who have sex with men in Swaziland. *J Int AIDS Soc*. 2013;16(Suppl 2):18715. [FREE Full text] [Medline: [24242263](#)]
40. Fitzgerald-Husek A, Van Wert M, Rosman L, Bowes A, Grosso A, Baral S. Measuring stigma related to sex worker identity and MSM identity: a systematic review. 2014. Presented at: APHA 142nd Annual Meeting and Expo; November 15-19, 2014; New Orleans, LA. URL: <https://apha.confex.com/apha/142am/webprogram/Paper312252.html>
41. Sanchez T, Sineath R, Kahle E, Tregear S, Sullivan P. The annual american men's internet survey of behaviors of men who have sex with men in the United States: protocol and key indicators report 2013. *JMIR Public Health Surveill*. 2015;1:e3. [FREE Full text] [doi: [10.2196/publichealth.4314](https://doi.org/10.2196/publichealth.4314)] [Medline: [27227126](#)]
42. Emory University. American Men's Internet Survey (AMIS): Online HIV Behavioral Survey of Men Who Have Sex With Men. Atlanta, GA. PRISM Health, Emory University; 2015. URL: http://www.cfar.emory.edu/downloads/news/2016/AMIS-2014_National_Report_&_Survey.pdf [accessed 2016-07-11]
43. Stahlman S, Bechtold K, Sweitzer S, Mothopeng T, Taruberekera N, Nkonyana J, et al. Sexual identity stigma and social support among men who have sex with men in Lesotho: a qualitative analysis. *Reprod Health Matters*. 2015;23:127-135. [doi: [10.1016/j.rhm.2015.11.017](https://doi.org/10.1016/j.rhm.2015.11.017)] [Medline: [26719004](#)]
44. Stahlman S, Grosso A, Ketende S, Mothopeng T, Taruberekera N, Nkonyana J, et al. Characteristics of men who have sex with men in southern Africa who seek sex online: a cross-sectional study. *J Med Internet Res*. 2015;17:e129. [FREE Full text] [doi: [10.2196/jmir.4230](https://doi.org/10.2196/jmir.4230)] [Medline: [26006788](#)]
45. Stahlman S, Holland C, Ketende S, Lith L, Kochelani D, Mnisi Z, et al. Stigma and openness about sexual identity among men who have sex with men: a latent class analysis. 2016. Presented at: Conference on Retroviruses and Opportunistic Infections (CROI); February 22-25, 2016; Boston, MA. URL: <http://www.croiconference.org/sessions/stigma-and-openness-about-sexual-identity-among-msm-latent-class-analysis>
46. Heckathorn D. Respondent-driven sampling: a new approach to the study of hidden populations. *Social Problems*. 1997;44:174-199. [doi: [10.2307/3096941](https://doi.org/10.2307/3096941)]
47. Biernacki P, Waldorf D. Snowball sampling: problems and techniques of chain referral sampling. *Sociological Methods & Research*. 1981;10:141-163. [doi: [10.1177/004912418101000205](https://doi.org/10.1177/004912418101000205)]
48. International Lesbian Gay Bisexual Trans and Intersex Association. State Sponsored Homophobia Report. 2015. URL: <http://ilga.org/what-we-do/state-sponsored-homophobia-report/> [accessed 2016-07-12] [WebCite Cache ID 6iwgGWsjj]
49. Itaborahy L, Zhu J. A World Survey of Laws: Criminalisation, Protection and Recognition of Same-Sex Love. International Lesbian Gay Bisexual Trans and Intersex Association; 2013. URL: http://old.ilga.org/Statehomophobia/ILGA_State_Sponsored_Homophobia_2013.pdf [accessed 2016-07-12] [WebCite Cache ID 6iwgRTnTs]
50. Itaborahy LP, Zhu J. A World Survey of Laws: Criminalisation, Protection and Recognition of Same-Sex Love. International Lesbian Gay Bisexual Trans and Intersex Association; 2014. URL: http://old.ilga.org/Statehomophobia/ILGA_SSHR_2014_Eng.pdf [accessed 2016-07-12] [WebCite Cache ID 6iwgaEQZ6]

51. Yebei VN, Fortenberry JD, Ayuku DO. Felt stigma among people living with HIV/AIDS in rural and urban Kenya. *Afr Health Sci*. 2008;8:97-102. [FREE Full text] [Medline: [19357758](#)]
52. Halkitis PN, Perez-Figueroa RE, Carreiro T, Kingdon MJ, Kupprat SA, Eddy J. Psychosocial burdens negatively impact HIV antiretroviral adherence in gay, bisexual, and other men who have sex with men aged 50 and older. *AIDS Care*. 2014;26:1426-1434. [FREE Full text] [doi: [10.1080/09540121.2014.921276](#)] [Medline: [24865599](#)]
53. Lépine A, Terris-Prestholt F, Vickerman P. Determinants of HIV testing among Nigerian couples: a multilevel modelling approach. *Health Policy and Planning*. 2014;30:579-592. [doi: [10.1093/heapol/czu036](#)]
54. Kahana SY, Jenkins RA, Bruce D, Fernandez MI, Hightow-Weidman LB, Bauermeister JA, et al. Adolescent Medicine Trials Network for HIV/AIDS Interventions. Structural determinants of antiretroviral therapy use, HIV care attendance, and viral suppression among adolescents and young adults living with HIV. *PLoS One*. 2016;11:e0151106. [FREE Full text] [doi: [10.1371/journal.pone.0151106](#)] [Medline: [27035905](#)]
55. Wei C, Cheung DH, Yan H, Li J, Shi L, Raymond HF. The impact of homophobia and HIV stigma on HIV testing uptake among chinese men who have sex with men: a mediation analysis. *J Acquir Immune Defic Syndr*. 2016;71:87-93. [doi: [10.1097/QAI.0000000000000815](#)] [Medline: [26334742](#)]
56. Secor AM, Wahome E, Micheni M, Rao D, Simoni JM, Sanders EJ, et al. Depression, substance abuse and stigma among men who have sex with men in coastal Kenya. *AIDS*. 2015;Suppl 3:S251-S259. [FREE Full text] [doi: [10.1097/QAD.0000000000000846](#)] [Medline: [26562814](#)]
57. Choi K, Steward WT, Miège P, Hudes E, Gregorich SE. Sexual stigma, coping styles, and psychological distress: a longitudinal study of men who have sex with men in Beijing, China. *Arch Sex Behav*. 2015. [doi: [10.1007/s10508-015-0640-z](#)] [Medline: [26679303](#)]
58. Logie CH, Newman PA, Chakrapani V, Shunmugam M. Adapting the minority stress model: associations between gender non-conformity stigma, HIV-related stigma and depression among men who have sex with men in South India. *Soc Sci Med*. 2012;74:1261-1268. [doi: [10.1016/j.socscimed.2012.01.008](#)] [Medline: [22401646](#)]
59. Zervoulis K, Lyons E, Dinos S. Stigma and self-esteem across societies: avoiding blanket psychological responses to gay men experiencing homophobia. *BJPsych Bull*. 2015;39:167-173. [FREE Full text] [doi: [10.1192/pb.bp.114.048421](#)] [Medline: [26755948](#)]
60. Yannesss JF, Reece M, Basta TB. HIV provider perspectives: the impact of stigma on substance abusers living with HIV in a rural area of the United States. *AIDS Patient Care STDS*. 2008;22:669-675. [doi: [10.1089/apc.2007.0151](#)] [Medline: [18627281](#)]
61. Rost K, Smith GR, Taylor JL. Rural-urban differences in stigma and the use of care for depressive disorders. *J Rural Health*. 1993;9:57-62. [Medline: [10124199](#)]
62. Glick SN, Golden MR. Persistence of racial differences in attitudes toward homosexuality in the United States. *J Acquir Immune Defic Syndr*. 2010;55:516-523. [FREE Full text] [doi: [10.1097/QAI.0b013e3181f275e0](#)] [Medline: [20838226](#)]
63. Lewis GB. Black-White differences in attitudes toward homosexuality and gay rights. *Public Opinion Quarterly*. 2003;67:59-78. [doi: [10.1086/346009](#)]
64. Quinn K, Dickson-Gomez J, DiFranceisco W, Kelly JA, St Lawrence JS, Amirkhanian YA, et al. Correlates of internalized homonegativity among black men who have sex with men. *AIDS Educ Prev*. 2015;27:212-226. [FREE Full text] [doi: [10.1521/aeap.2015.27.3.212](#)] [Medline: [26010313](#)]
65. Garofalo R, Hottot AL, Kuhns LM, Gratzner B, Mustanski B. Incidence of HIV infection and sexually transmitted infections and related risk factors among very young men who have sex with men. *J Acquir Immune Defic Syndr*. 2016. [doi: [10.1097/QAI.0000000000000933](#)] [Medline: [26745827](#)]
66. Bedoya CA, Mimiaga MJ, Beauchamp G, Donnell D, Mayer KH, Safren SA. Predictors of HIV transmission risk behavior and seroconversion among Latino men who have sex with men in Project EXPLORE. *AIDS Behav*. 2012;16:608-617. [FREE Full text] [doi: [10.1007/s10461-011-9911-4](#)] [Medline: [21390540](#)]
67. Mustanski B, Newcomb M, Du Bois SN, Garcia S, Grov C. HIV in young men who have sex with men: a review of epidemiology, risk and protective factors, and interventions. *J Sex Res*. 2011;48:218-253. [FREE Full text] [doi: [10.1080/00224499.2011.558645](#)] [Medline: [21409715](#)]
68. Baral SD, Grosso A, Holland C, Papworth E. The epidemiology of HIV among men who have sex with men in countries with generalized HIV epidemics. *Curr Opin HIV AIDS*. 2014;9:156-167. [doi: [10.1097/COH.0000000000000037](#)] [Medline: [24445371](#)]
69. Sullivan PS, Rosenberg ES, Sanchez TH, Kelley CF, Luisi N, Cooper HL, et al. Explaining racial disparities in HIV incidence in black and white men who have sex with men in Atlanta, GA: a prospective observational cohort study. *Ann Epidemiol*. 2015;25:445-454. [doi: [10.1016/j.annepidem.2015.03.006](#)] [Medline: [25911980](#)]
70. McAdams-Mahmoud A, Stephenson R, Rentsch C, Cooper H, Arriola KJ, Jobson G, et al. Minority stress in the lives of men who have sex with men in Cape Town, South Africa. *J Homosex*. 2014;61:847-867. [doi: [10.1080/00918369.2014.870454](#)] [Medline: [24392722](#)]
71. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychol Bull*. 2003;129:674-697. [FREE Full text] [doi: [10.1037/0033-2909.129.5.674](#)] [Medline: [12956539](#)]

72. D'augelli A, Grossman A. Disclosure of sexual orientation, victimization, and mental health among lesbian, gay, and bisexual older adults. *Journal of Interpersonal Violence*. 2001;16:1008-1027. [doi: [10.1177/088626001016010003](https://doi.org/10.1177/088626001016010003)]
73. Cahill S, Valadéz R, Ibarrola S. Community-based HIV prevention interventions that combat anti-gay stigma for men who have sex with men and for transgender women. *J Public Health Policy*. 2013;34:69-81. [doi: [10.1057/jphp.2012.59](https://doi.org/10.1057/jphp.2012.59)] [Medline: [23151921](https://pubmed.ncbi.nlm.nih.gov/23151921/)]
74. Charurat ME, Emmanuel B, Akolo C, Keshinro B, Nowak RG, Kennedy S, et al. Uptake of treatment as prevention for HIV and continuum of care among HIV-positive men who have sex with men in Nigeria. *J Acquir Immune Defic Syndr*. 2015;Suppl 2:S114-S123. [FREE Full text] [doi: [10.1097/QAI.0000000000000439](https://doi.org/10.1097/QAI.0000000000000439)] [Medline: [25723975](https://pubmed.ncbi.nlm.nih.gov/25723975/)]
75. McNaghten A, Kearns R, Siegler AJ, Phaswana-Mafuya N, Bekker L, Stephenson R, et al. Methods for prevention packages program project protocol: pilot study of hiv prevention interventions for men who have sex with men in South Africa. *JMIR Res Protoc*. 2014;3:e55. [FREE Full text] [doi: [10.2196/resprot.3737](https://doi.org/10.2196/resprot.3737)] [Medline: [25325296](https://pubmed.ncbi.nlm.nih.gov/25325296/)]
76. van der Elst EM, Kombo B, Gichuru E, Omar A, Musyoki H, Graham SM, et al. The green shoots of a novel training programme: progress and identified key actions to providing services to MSM at Kenyan health facilities. *J Int AIDS Soc*. 2015;18:20226. [FREE Full text] [Medline: [26493863](https://pubmed.ncbi.nlm.nih.gov/26493863/)]
77. Colchero MA, Bautista-Arredondo S, Cortés-Ortiz MA, Romero-Martinez M, Salas J, Sosa-Rubí SG, et al. Impact and economic evaluations of a combination prevention programme for men who have sex with men in Mexico. *AIDS*. 2016;30:293-300. [doi: [10.1097/QAD.0000000000000925](https://doi.org/10.1097/QAD.0000000000000925)] [Medline: [26691549](https://pubmed.ncbi.nlm.nih.gov/26691549/)]
78. Henderson M. On the Ground: Programmes Serving the Needs of Key Populations. World Health Organization; 2014. URL: http://apps.who.int/iris/bitstream/10665/145002/1/WHO_HIV_2014.50_eng.pdf [accessed 2016-07-12] [WebCite Cache ID 6iwhyDjzo]
79. Toomey R, Ryan C, Diaz R, Russell S. High school gay-straight alliances (GSAs) and young adult well-being: an examination of GSA presence, participation, and perceived effectiveness. *Applied Developmental Science*. 2011;15:175-185. [doi: [10.1080/10888691.2011.607378](https://doi.org/10.1080/10888691.2011.607378)]
80. Adebajo S, Obianwu O, Eluwa G, Vu L, Oginni A, Tun W, et al. Comparison of audio computer assisted self-interview and face-to-face interview methods in eliciting HIV-related risks among men who have sex with men and men who inject drugs in Nigeria. *PLoS One*. 2014;9:e81981. [FREE Full text] [doi: [10.1371/journal.pone.0081981](https://doi.org/10.1371/journal.pone.0081981)] [Medline: [24416134](https://pubmed.ncbi.nlm.nih.gov/24416134/)]
81. Castro R. Inconsistent respondents and sensitive questions. *Field Methods*. 2012;25:283-298. [doi: [10.1177/1525822x12466988](https://doi.org/10.1177/1525822x12466988)]
82. Stahlman S, Johnston LG, Yah C, Ketende S, Maziya S, Trapence G, et al. Respondent-driven sampling as a recruitment method for men who have sex with men in southern sub-Saharan Africa: a cross-sectional analysis by wave. *Sex Transm Infect*. 2016;92:292-298. [doi: [10.1136/sextrans-2015-052184](https://doi.org/10.1136/sextrans-2015-052184)] [Medline: [26424712](https://pubmed.ncbi.nlm.nih.gov/26424712/)]

Abbreviations

CBOs: community-based organizations
CI: confidence interval
HIV: human immunodeficiency virus
IQR: interquartile range
IRB: institutional review board
MSM: men who have sex with men
PR: prevalence ratio
RDS: respondent driven sampling

Edited by G Eysenbach; submitted 31.03.16; peer-reviewed by S LeGrand, C Oldenburg; comments to author 03.05.16; revised version received 20.05.16; accepted 10.06.16; published 26.07.16

Please cite as:

Stahlman S, Sanchez TH, Sullivan PS, Ketende S, Lyons C, Charurat ME, Drame FM, Diouf D, Ezouatchi R, Kouanda S, Anato S, Mothopeng T, Mnisi Z, Baral SD

The Prevalence of Sexual Behavior Stigma Affecting Gay Men and Other Men Who Have Sex with Men Across Sub-Saharan Africa and in the United States

JMIR Public Health Surveill 2016;2(2):e35

URL: <http://publichealth.jmir.org/2016/2/e35/>

doi: [10.2196/publichealth.5824](https://doi.org/10.2196/publichealth.5824)

PMID: [27460627](https://pubmed.ncbi.nlm.nih.gov/27460627/)

©Shauna Stahlman, Travis Howard Sanchez, Patrick Sean Sullivan, Sosthenes Ketende, Carrie Lyons, Manhattan E Charurat, Fatou Maria Drame, Daouda Diouf, Rebecca Ezouatchi, Seni Kouanda, Simplicie Anato, Tampose Mothopeng, Zandile Mnisi, Stefan David Baral. Originally published in JMIR Public Health and Surveillance (<http://publichealth.jmir.org>), 26.07.2016. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.