

Original Paper

Whiplash Syndrome Reloaded: Digital Echoes of Whiplash Syndrome in the European Internet Search Engine Context

Michael Noll-Hussong, MD

Department of Psychosomatic Medicine and Psychotherapy, University of Ulm, Ulm, Germany

Corresponding Author:

Michael Noll-Hussong, MD

Department of Psychosomatic Medicine and Psychotherapy

University of Ulm

Albert-Einstein-Allee 23

Ulm, 89081

Germany

Phone: 49 731 500 ext 61833

Fax: 49 731 500 61802

Email: minohu@gmx.net

Abstract

Background: In many Western countries, after a motor vehicle collision, those involved seek health care for the assessment of injuries and for insurance documentation purposes. In contrast, in many less wealthy countries, there may be limited access to care and no insurance or compensation system.

Objective: The purpose of this infodemiology study was to investigate the global pattern of evolving Internet usage in countries with and without insurance and the corresponding compensation systems for whiplash injury.

Methods: We used the Internet search engine analytics via Google Trends to study the health information-seeking behavior concerning whiplash injury at national population levels in Europe.

Results: We found that the search for “whiplash” is strikingly and consistently often associated with the search for “compensation” in countries or cultures with a tort system. Frequent or traumatic painful injuries; diseases or disorders such as arthritis, headache, radius, and hip fracture; depressive disorders; and fibromyalgia were not associated similarly with searches on “compensation.”

Conclusions: In this study, we present evidence from the evolving viewpoint of naturalistic Internet search engine analytics that the expectations for receiving compensation may influence Internet search behavior in relation to whiplash injury.

(*JMIR Public Health Surveill* 2017;3(1):e15) doi: [10.2196/publichealth.7054](https://doi.org/10.2196/publichealth.7054)

KEYWORDS

search engine; whiplash injuries; legislation & jurisprudence; medicolegal aspects; compensation and redress; compensation; accidents, traffic; adult; female; humans; incidence; insurance claim reporting; male; neck pain; prognosis; search engine analytics; whiplash syndrome; Google Trends

Introduction

In many Western countries, after a motor vehicle collision, those involved seek health care for the assessment of injuries and for insurance documentation purposes. In contrast, in many less wealthy countries, there may be limited access to care and insurance may only be available to the wealthy. Against this background, the “whiplash syndrome” (ICD-10: S13.4) has been one special focus of continuous and controversial scientific research since the 1950s [1-5] (Figure 1) as the worldwide incidence of such injuries varies enormously 16-2000 per 100,000 population and the late whiplash syndrome in these

cases varies between 18% to 40% [6]. Whiplash injuries are estimated to cost European society up to 10 billion euro per year [7]. Recently, and after extensive evaluation of over 1600 publications about whiplash listed in *Pubmed* [8] since 1996, the nosology of the chronic whiplash syndrome has been still classified as “doubtful” [9].

Now, 2 decades after Schrader et al’s important work in *The Lancet* showing that late whiplash syndrome after a motor vehicle collision is rare or uncommon in Lithuania [10], and Cassidy et al’s conclusion in the *New England Journal of Medicine* that “the elimination of compensation for pain and suffering is associated with a decreased incidence and improved

prognosis of whiplash injury” [11], a new method of examining different types of “contagiosity” of diseases has evolved using the Internet search engine analytics [12-15].

According to Wikipedia, search analytics “is the use of search data to investigate particular interactions among Web searchers, the search engine, or the content during searching episodes. (...)

Search analytics includes search volume trends and analysis, reverse searching (entering websites to see their keywords), keyword monitoring, search result and advertisement history, advertisement spending statistics, website comparisons, affiliate marketing statistics, and multivariate ad testing” [16]. The Internet usage especially in some European countries is presented in Table 1 [17].

Table 1. Internet usage on November 30, 2015.

Internet usage	Internet users	Penetration rate (% population)
Worldwide	3,366,261,156	46.37
Europe	604,147,280	73.54
Switzerland (CH)	7,180,749	87.18
Germany (DE)	71,727,551	88.36
Finland (FI)	5,117,660	93.53
France (FR)	55,429,382	83.82
Greece (GR)	6,834,560	63.21
Lithuania (LT)	2,399,678	82.15
United Kingdom (UK)	59,333,154	91.61

The number of Europeans using the Internet to obtain health information is significantly growing in all age groups, but there is especially strong growth among young women [18,19]. Individuals presented with chronic pain associated with whiplash injury are more likely to be female [20]. Internet search queries are exhaustively cataloged for marketing purposes by search engine providers [21]. Thus, as a “side-effect,” an analysis of Internet search queries can also “detect” public interests in infectious (eg, influenza) [22] and noninfectious [23,24] diseases. In addition to gathering epidemiologic data on disease incidence and prevalence through traditional, labor-intensive processes involving large surveys, chart reviews, prospective studies, or extraction from previously created databases, Internet search trend analysis tools, since they provide self-reported information by consumers, represent a complementary source of information on a population level [25,26]. The subjectively perceived “anonymity” in using the most popular organic Internet search engine may be attractive to consumers because some diseases are burdened with a social stigma [27] or are controversial and linked to monetary [28] or secondary gain [29]. Thus, Internet search data may reduce selection bias in some aspects, even though it is equally challenging to confirm the source. At the very least, using an Internet tool, culture-related attributions can be mapped on a global population level [30].

Thus, this source has the potential to reveal epidemiologic trends and patterns in near real time and with minimal expense. The current leading Internet search engine provider is owned by Alphabet Inc (marketed as Alphabet), Google Inc, which is also the brand name of the most visited website worldwide [31]. This information is freely provided to the public through Google Trends. It is of note that globally there are at present more than 3.5 billion Google searches per day and 1.2 trillion searches per year worldwide [32].

In terms of the controversial whiplash syndrome, countries in which there is an established compensation system for whiplash injury might be expected to have more Internet traffic and volume regarding whiplash injury than in countries without an established compensation system. In other words, “diverse assessments and principles for approving a claim are reflected in the fact that the prevalence of chronic spine pain after whiplash injuries (late whiplash syndrome) varies between 16% and 71% in different countries, and the proportion of whiplash injuries involved in petitions for compensation differs greatly across Europe” [33] as France and Finland [34] have the lowest and Great Britain the highest incidence of minor cervical spine trauma (eg, United Kingdom 75%, Germany 47%, Switzerland 33%, Finland 8.5%, and France 3% of all personal injuries), whereas in Greece and Lithuania whiplash injury is reported to be an almost nonexistent condition [6].


Regarding the latter, Obelieniene et al state, “Lithuania is a country in which there is no or little awareness or experience among the general population of the notion that a whiplash injury may cause chronic pain and disability. Accident victims with acute symptoms from rear end collisions generally view this as a benign injury not requiring any medical attention” [35]. Thus, it has been “hypothesized that cultural [36] and psychosocial [37-39] factors may be important in explaining why accident victims in some other societies report chronic symptoms. Such factors may include expectation of disability, symptom amplification as a result of this expectation, the effects of inappropriate therapy, insurance [40], and attribution of symptoms from nonaccident related causes (spontaneous symptoms, occupational symptoms, symptoms before an accident being amplified after an accident)” [35].

As a pilot effort, this paper deals with Google-based Internet search engine statistics on the search for “whiplash syndrome”

in European countries to offer insight into both the method of Internet-based population epidemiology in whiplash-associated disorders, and the condition itself in the context of various pain cultures and national social insurance or compensation systems [41]. Specifically, the first purpose of this infodemiology or infoveillance study was to compare the Internet search patterns in Germany and the United Kingdom, countries with established compensation systems for whiplash injury, to those used in Greece and Lithuania, countries where a system for monetary compensation for motor vehicle collision injury has not yet been established. Second, in order to validate our data, we looked for the European countries for which the lowest incidence of minor cervical spine trauma has been described, that is, Finland and France [6]. Third, we compared the “googled” whiplash data with Internet search patterns concerning frequent or traumatic painful injuries; diseases and disorders like arthritis, headache, radius, and hip fracture; depressive disorders [42]; and fibromyalgia. Finally, we wanted to test if there were hints

that search engine usage may reflect national changes in the medicolegal compensation rules as has been shown for Saskatchewan, Canada, where the tort-compensation system for traffic injuries, that includes payments for health and suffering, was changed to a no-fault system in 1995, which did not include such payments, resulting in a decreased incidence and improved prognosis of whiplash injury [11]. As publicly available records of search engine analytics in Google start with 2005, we chose a similar event in Switzerland; the federal court abridged the possibilities for receiving a disability pension after whiplash injury in August 2010 [43] (modified again in 2015: DFR - BGer 9C_492/2014; 03.06.2015). In this context it is interesting to know that Switzerland has the highest expenditure per claim at an average cost of €35,000 compared with the European average of €9,000, and there are large differences between German-speaking and French- or Italian-speaking parts of Switzerland [6].

Figure 1. Number of publications on “whiplash” or “whiplash associated disorders” cited in Pubmed during the period from 1956 to 2010.



Methods

Internet Search Engine Analytics via Google Trends and the Whiplash Syndrome

In order to compare the pattern of the Internet usage surrounding whiplash injury and compensation in Germany, Finland, France, Greece, Lithuania, and the United Kingdom, we first used Google Trends to search for uncapitalized national lay terminologies related to whiplash syndrome, such as the English word “whiplash injury,” the Finnish equivalent “piiskansiima” or “piiskaniskuvammoilta vammoja,” the French equivalent “coup de fouet (cervicale)” or “coup du lapin,” the German equivalent “schleudertrauma,” the Greek “αυχενικού τραυματισμού,” and the Lithuanian “Bicz žalos” or “kaklo nyris” or “kaklo slankstelių trauma,” and, for a more lenient “threshold,” “kaklo skausmas” (neck pain). Then we searched for the term “injury compensation” (and the respective

appropriate translations and back-translations with native speakers) in Germany, Finland, France, Greece, Lithuania, and the United Kingdom.

Then, to obtain an impression of whether or not overall Internet usage in Europe for health information was different in Finland, France, Greece, and Lithuania in general, we repeated this language-adapted assessment of Google usage for arthritis (a condition known to be associated with chronic pain) in all mentioned countries, to determine how searches for this term may have differed among Germany, France, Greece, Lithuania, and the United Kingdom. In Northern Europe, the incidence of rheumatoid arthritis, the clinically most relevant subtype of arthritis, is estimated at 20-50 cases per 100,000 population and the prevalence at 0.5-1.1%, lower incidences and prevalence have been reported in Southern Europe [44]. Moreover, as more than 50% of community-dwelling adults in Europe indicate that they suffer from headache in general during the last year or less,

with most headaches more prevalent in women [45], we searched for “headache” (ICD-10: R51). Important traumatic events apart from whiplash injury are hip fracture (ICD-10: S72) and radius fracture (ICD-10: S52), which are a major public health problem in the elderly and the active younger adults, respectively [46-50]. Having said this, the incidence rates of hip fracture vary from northern to southern Europe, with the highest levels in Sweden and Norway and the lowest in France and Switzerland. The reported age-standardized annual incidence rate of hip fracture, for example, in Switzerland is 346/10,000 and 137.8/100,000 in women and men, respectively [51]. The incidence of distal radius fracture is in patients >35 years of age 0.37% in females, 0.09% in males.

Finally, we searched for “depression” (ICD-10: F32 and F33) and “fibromyalgia” (ICD-10: M97) [52-54]. According to the World Health Organization (WHO), “depression is a common mental disorder. Globally, an estimated 350 million people of all ages suffer from depression. Depression is the leading cause of disability worldwide and is a major contributor to the overall global burden of disease” [55]. The 12-month prevalence of major depression is estimated as 6.9% among all Europeans [52]. Fibromyalgia is a disorder both characterized basically by chronic widespread pain and mental symptoms like fatigue, cognitive disturbances, and other symptoms, and likewise controversially or dogmatically or ideologically discussed as (late) whiplash injury. Thus, fibromyalgia appears to be a common condition in most European countries affecting up to 2% of the general population [56-60].

The Web browser we used was Mozilla Firefox 11ff [61]. The search window in Google Trends was, where not otherwise stated, restricted to the 6-year time period between January 1, 2005 and December 31, 2010: during 2011, Google updated the categorization taxonomy and modified their geographical assignment, which may lead to contorted results — therefore we skipped the year 2011 for this analysis. Moreover, we did not want to include ongoing Google searches in 2012, as this would change the data, although we cannot guarantee that Google will or has changed the Google Trends algorithms that would affect analysis of the retrospective data shown here. Finally, all of the data presented here can be most easily validated by individually using Google Trends repeating our analysis. With regard to data consistency, all data was obtained in Germany starting in March 2012 with at least annual verifications ending in August 2016, always with continuous consistent results for the time period specified previously.

Technical Background of Internet Search Engine Analytics via Google Trends

The current world-leading search engine provider, Google Inc, provides, since August 2008, a publicly free available Internet search analytics tool based on Google search queries currently named Google Trends until September 27, 2012, known as Google Insights for Search (GIS) [62]. Google, in each case in its nationalized version, is, if uncensored, the most visited website internationally, with by far the highest market share in the search engine market in Europe [63]. For more information in the audience demographics for Google in each of mentioned countries see, for example, Alexa Internet, Inc [64].

According to information on the Google Trends website, one “can explore ‘trending stories’ in real time by category and location” using the Google Trends homepage [65]. In the appropriate frequently asked questions (FAQ) section, Google elaborates:

A trending story is a collection of Knowledge Graph topics, Search interest, trending YouTube videos, and/or Google News articles detected by our algorithms. Trending Stories rely on technology from the Knowledge Graph across Google Search, Google News, and YouTube to detect when topics are trending on these three platforms. The Knowledge Graph enables our technology to connect searches with real-world things and places. The algorithm for trending stories groups topics together that are trending at the same time on Google News, Google Search, and YouTube and ranks stories based on the relative spike in volume and the absolute volume of searches. (...) Google Trends analyzes a percentage of Google web searches to figure out how many searches were done over a certain period of time. Trends only analyzes data for popular terms, so search terms with low volume appear as 0, eliminates repeated searches from the same person over a short period of time, and filters out queries with apostrophes and other special characters. (...) Google Trends adjusts search data to make comparisons between terms easier. Otherwise, places with the most search volume would always be ranked highest. To do this, each data point is divided by the total searches of the geography and time range it represents, to compare relative popularity. The resulting numbers are then scaled to a range of 0 to 100. Data is relative across regions, i.e., just because two regions show the same number of searches for a term doesn't mean that their total search volumes are the same.

The calculation of search numbers is performed using the spelling, exactly as entered, and appropriate language [66] for Google search queries over a given period of time. The data do not contain personal information. The Internet protocol (IP) addresses of the protocols establish an educated guess on the search origins. Google elaborates that the data are normalized by dividing the datasets by a common variable to remove the effect of that variable on the data. This normalization allows a comparison of the underlying dataset characteristics. Thus, this tool does not provide absolute numbers of searches but rather a relative estimation based on search activity for the time period under study [67]. The analysis can compare 5 search terms simultaneously. If Google displayed the absolute rankings, data from regions generating the most search volume would always be ranked high (for details see [22,68]). Wikipedia states that query analysis in the context of geographical and temporal parameters produces so-called “vectors,” which may partially represent the life- and interest-space of the respective searchers [69]. Separate searches in a common context are feasible in many cases, which provide more differentiated vectors. Informative relationships and common motivators can be

determined using parallel search volumes and cross-comparisons, which may be profitable as forecasts and may be retrospectively instructive for both research and marketing.

“Top searches” are search terms with the most significant levels of interest. Google states: “These terms are related to the term you’ve entered; if you didn’t enter in a search term, the top searches will be related to the category or country/territory you’ve chosen. Google determines relativity by examining searches that have been conducted by a large group of users preceding the search term you’ve entered, as well as after.”

Furthermore, as “Insights for Search examines the past values for the terms you’ve entered, it can extrapolate the future values, creating a forecast of search trends for those terms. This prediction model doesn’t take into account the context of the search term or its category, nor does it account for any business cycles that may be driving a specific market (for details, see [70]).”

Finally, Google warns on their website that the analytical data provided “aims to provide insights into broad search patterns. Several approximations are used to compute these results. The Insights for Search (or Google Trends) map is intended for general analysis of volume patterns. Borders are an

approximation and may not be accurate.” Thus, Google releases its own data only in an aggregated way and often without assigning absolute values, such as the number of visitors to its graphs.

Results

Main Results

A comparison of the normalized data for the countrywide Google searches revealed that Google top searches for whiplash injury in Germany and the United Kingdom showed sufficient search volume and were frequently accompanied by searches for “compensation.” In other words, the concatenation of national search interest between these 2 topics such as “whiplash injury compensation” or its German counterpart “schmerzensgeld schleudertrauma,” was apparently common (see [Multimedia Appendix 1](#)). The top searches for whiplash injury in the “health” and “law and government” category for Germany and the United Kingdom can be found in [Multimedia Appendix 1](#), respectively. Searching for “whiplash” (without “injury”) in the United Kingdom revealed the following top searches (spelling not corrected), where 18 out of 47 or ~38% of the top search results were at least semantically associated with *compensation* ([Table 2](#)).

Table 2. Google Trends-ranked “top searches” for “whiplash” in the United Kingdom.

Rank (#)	Top searches
1	symptoms whiplash
2	whiplash injury
3	<i>compensation</i>
4	<i>compensation</i> whiplash
5	whiplash <i>claim</i>
6	whiplash injuries
7	accident whiplash
8	miss whiplash
9	whiplash <i>claims</i>
10	neck whiplash
11	<i>compensation</i> for whiplash
12	symptoms of whiplash
13	<i>claim</i> for whiplash
14	iron man whiplash
15	car accident whiplash
16	whiplash injury <i>compensation</i>
17	whiplash treatment
18	what is whiplash
19	whiplash <i>payout</i>
20	whiplash injury <i>claim</i>
21	neck pain
22	whiplash
23	whiplash injury symptoms
24	whiplash lyrics
25	whiplash injury <i>claims</i>
26	whiplash neck injury
27	neck injury
28	<i>claims</i> for whiplash
29	whiplash trash
30	whiplash <i>claiming</i> whiplash
31	whiplash scooter
32	whiplash effects
33	average whiplash <i>payout</i>
34	whiplash marvel
35	accident <i>claims</i>
36	whiplash guidelines
37	whiplash symptoms
38	symptoms for whiplash
39	whiplash monkey
40	whiplash <i>payouts</i>
41	whiplash braid
42	Berkley whiplash

Rank (#)	Top searches
43	average whiplash <i>claim</i>
44	treatment for whiplash
45	whiplash syndrome
46	<i>compensation</i> calculator
47	whiplash <i>compensation</i> uk

The mentioned combination of Google users search interest for “whiplash” with “compensation” was not detected in the nationalized search queries in Finland (“not enough search volume to show graphs”), France (“top searches: not enough search volume to show results”), Greece (“not enough search volume to show graphs”), or Lithuania (“not enough search volume to show graphs”).

Searching for “compensation” under the Google category “health” in the United Kingdom revealed that “whiplash” and “whiplash compensation” were ranked third and fourth under top searches, after “injury compensation” (rank 1) and “compensation act” (rank 2). Searching in the United Kingdom for “compensation” in “all categories” revealed “whiplash” for the first time at rank 14, and searching for “compensation” in “law and government category” revealed “whiplash” at rank 8 ([Multimedia Appendix 1](#)). Searching for “injury compensation” in the Google category “health” in the United Kingdom revealed “whiplash injury” at rank 1 of concatenated top searches ([Multimedia Appendix 1](#)).

Looking for analog searches in “all categories” in Germany for “schmerzensgeld” (injury compensation) shows that “schleudertrauma” (whiplash injury) is at rank 3 of “top searches” ([Multimedia Appendix 1](#)). The differing distribution of “whiplash” in the different Google categories with respect to Germany and the United Kingdom may be due to categorization inconsistencies by Google.

However, no similar correlation could be detected for Lithuania searching for “atlyginimas už kūno sužalojimą,” “žalos atlyginimas,” or “žalos kompensacija,” for “kipuraha+vamman korvaukset” (injury compensation) in Finland or “αποζημίωση τραυματισμών” in Greece, as these searches revealed “not enough search volume to show graphs.” Searching for “indemnisation+dommages et intérêts” in France revealed the following results: “indemnisation chômage, chômage, indemnisation assedic, indemnisation accident, indemnisation maladie, indemnisation assurance, accident travail indemnisation, chômage partiel, indemnisation chômage partiel, assedic indemnisation chômage, indemnisation pole employ, indemnisation licenciement, accident du travail, (...)”.

Comparison With Other Diseases

Checking for top searches in all categories for “arthritis” in Switzerland (German-speaking part of Switzerland) and Germany, “niveltulehdus” in Finland, “arthritis” in France, “αρθρίτιδα” in Greece, “artritas” in Lithuania, and “arthritis” in the United Kingdom revealed the results shown in [Multimedia Appendix 1](#). Searching for arthritis, the Internet users were transnationally most interested in terms like arthritis symptoms or arthritis treatment. As an aside, we did not find a

concatenation of top searches or rising searches of arthritis with compensation in any of the 3 countries.

Checking for top searches for “headache” in the United Kingdom, “kopfschmerzen” in Switzerland and Germany, “pänsärky” in Finland, “mal de tête” in France, “πονοκέφαλο” in Greece (“not enough search volume to show graphs”), and “galvos skausmas” in Lithuania (“not enough search volume to show results”) revealed the results presented in [Multimedia Appendix 1](#).

Checking for top searches in all categories for “hüftfraktur” (hip fracture) were done for Switzerland (German-speaking part of Switzerland) and Germany (both: “not enough search volume to show graphs”), “fracture de la hanche” in France (“not enough search volume to show results”), “lonkkamurtuman” in Finland (“not enough search volume to show graphs”), “κάταγμα ισχίου” in Greece (“not enough search volume to show graphs”), “šlaunikaulio lūžis” in Lithuania (“not enough search volume to show graphs”), and “hip fracture” in the United Kingdom (ranked geographic information system [GIS] top searches in the United Kingdom: “fracture of hip, hip fractures, neck of femur, hip replacement, hip fracture database, hip pain, hip fracture treatment, fractured hip, hip fracture guidelines, hip fracture classification, and hip fracture management”).

We also checked the top searches in all categories for “radius fracture” in the United Kingdom, “radiusfraktur” in Switzerland and Germany, “säde murtuma” in Finland (“not enough search volume to show results”), “fracture du radius” in France, “ακτίνα κάταγμα” in Greece (“not enough search volume to show results”), and “spindulys lūžis” in Lithuania (“not enough search volume to show results”); [Multimedia Appendix 1](#).

We checked for top searches in all categories for “depression” in Switzerland, Germany, France, and the United Kingdom (same spelling in mentioned countries), “masennus” in Finland, “κατάθλιψη” in Greece, and “depresija” in Lithuania (we did not rule out the economic meaning of this term choosing the “health” category because this category isn’t, eg, available for Lithuania) and the results are shown in [Multimedia Appendix 1](#). Searching for depression, the Internet users were transnationally most interested in terms like “depression symptoms” or “depression tests” ([Multimedia Appendix 1](#)). As an aside, we did not find a concatenation of top searches or rising searches with “compensation” in any of the 3 countries.


We checked for “fibromyalgia” in the United Kingdom, Finland, and France, “fibromyalgie” in Switzerland and Germany, “ινομυαλγία” in Greece (“not enough search volume to show graphs”), and “fibromialgija” in Lithuania (“not enough search volume to show graphs”). Remarkably, no concatenation of

“fibromyalgia” and “compensation” (or their respective translations) could be found (Multimedia Appendix 1).

Searching for “schleudertrauma+coup de fouet cervicale+coup du lapin+colpo di frusta” (German, French, and Italian search

term) in the health category for Switzerland for the years 2007-2009 and 2010-2011, respectively, revealed a decline in Google search queries for “whiplash” (Figure 2).

Figure 2. Google Insights Screenshot of cumulative regional interest for “schleudertrauma+coup de fouet cervicale+coup du lapin+colpo di frusta” in Switzerland during the period from 2005 to 2010. Regional Interest (search volume) was highest in 1. Bern, 2. Zurich and 3. Vaud. Left Upper Inset: The search interest for mentioned search terms declined ~ 40% comparing the years 2007-2009 (mean: 33) with 2010-2011 (mean: 20).



Supplementary Notes

Searching for the number of advertisers for the respective national queries for “whiplash” on Alexa—the leading provider of global Web metrics—revealed that attorneys offering their assistance in law suits on personal injuries are among the top 5 in search ads for “whiplash” searches on major search engines in Germany and the United Kingdom, but not Lithuania [64].

Searching for “whiplash injury” on YouTube [71], which also belongs to Google, Inc, shows that there are about 43,700 unfiltered results (as on November 27, 2016), in which the top videos are more or less advertisements for nonevidence-based methods of “treatment” (in particular, showing the “benefits” or “secrets” of chiropractic care) and “whiplash injury compensation.”

Discussion

Principal Findings

This study shows that, in general, Lithuanians, Greeks, and Finns use the Internet to search for health information on

conditions such as arthritis and depression in much the same way as do those from Germany, the United Kingdom, and Switzerland. However, there is a marked difference in the patterns of searches for whiplash injury or similar terms in the two former countries and Germany, the United Kingdom, and Switzerland, countries known to have high compensation rates for whiplash injury. Searches in Germany, the United Kingdom, and Switzerland for whiplash are high ranking when one examines searches combined with terms like “compensation.”

One main result is that the aforementioned combination of Google searches reflecting combined consumer interests in “whiplash injury” and “compensation” was not detected in Lithuania and other European countries (Finland, France, Greece) where cultural and psychosocial factors, including expectations, and insurance systems, have been described as significantly different from countries in which the problem of chronic whiplash is highly prevalent [72]. Actually, in Lithuania there is no formal compensation system for late whiplash injury pain and suffering, and this fact may, amongst others, influence the coping styles of the respective persons concerned [73]. Moreover, our findings reflect the low incidence of late whiplash

in Finland [6,34], where total socioeconomic costs are estimated as about 1.5 million euro per annum, France [6,74], and Greece [75,76]. There has been a 70% rise in motor insurance injury claims over the 6 years leading up to 2012 in the United Kingdom, despite a 23% drop in the number of casualties actually caused by road accidents—and whiplash accounted for 70% of the total. That amounted to roughly 554,000 whiplash claims from 2010-11, that is more than 1500 claims a day. The whiplash injury costs in the United Kingdom are approximately 4.6 billion euro per annum [77]. In Germany, whiplash injury is number 1 of consequences after vehicle accidents with about 20,000 cases per year, and costs the insurance companies at least 500 million euro per annum, “official” compensation for pain and suffering due to whiplash is about 2000 euro (higher regional court (OLG) Frankfurt VRS 90, 254).

These different signs synoptically suggest that a biopsychosocial [78] understanding of chronic whiplash is important [79], especially in the “social” aspect, and the Internet is a social medium. Despite many years of research, the evidence regarding unquestionable risk factors for late whiplash is sparse but seems to include personal, societal, medicolegal, and environmental factors [80]. Against this background one should also mention an experimental study in 2001 in which participants were placed in a stationary vehicle with a curtain blocking their rear view, and exposed to a simulated rear-end collision [81]. Twenty percent of patients had symptoms at 3 days, despite the fact that no collision actually occurred [38].

Until now, the Internet search statistics [82] and social media [83] in medicine are mainly used for outbreak detection and/or the monitoring of transmissible [22,68,84,85], whereupon noninfectious diseases noticeably gain attention (eg, [86,87]). Google Insights for Search (now: Google Trends), initially developed by Google’s research and development center in Israel, is the most important freely available application on the World Wide Web. These systems are growing, and they provide multifaceted information concerning old and emerging disorders [88]. This intrinsic “predictive power” is associated with the phenomena of “swarm intelligence” [89] and typical, sometimes enigmatic properties of “social networks” [90]. However, the impact and reliability of these systems on medical and public health and individual physicians is not certain [91]. Information overload [92] in times of “Health 2.0” [93], incorrect reports (as has been shown, eg, for psychological trauma-related [94]), or Web-based information on low back pain [95]), the lack of signal specificity [96], information filtered by Internet search engine providers (ie, economic [97], political or social search engine bias) [98], media or marketing [99] interest, differing search strategies (eg, [100]), misspellings, Internet availability and local specialties [101,102], age-related differences in the accuracy of Web query-based predictions [103], seasonal effects [104], problems with incidence peaks [82,105], the unforeseeable or undisclosed evolution of search algorithms or models [106], noise [107], and (last, not least) statistical issues

concerning the analysis of time series [108,109] are among the manifold confounding factors that may interfere with the development and reliability of the Internet search engine analytics, even in the medical sector [91,110]. Moreover, and from a more clinical point of view, Web-based information gathering may foster greater patient engagement in health maintenance and care [111]. Conversely, there is a relationship between searching for health information on the Web and health anxiety, a phenomenon recently named “cyberchondria,” which may inversely influence the health of the respective Internet searchers [112,113]. Against this background, we may have to develop feasible models and tools for consumers to assess and filter health information on the Internet [114]. It is of note that other societal factors appear to also play significant roles in the rate of development of late whiplash disorder as it has been shown that, at least within Canada, regions with similar compensation systems have large differences in rates occurrence [11].

Perspective

Future investigations will deepen our knowledge in the growing field of search engine analytics as kind of infodemiology [85,115] (or “i-epidemiology”) of the worldwide social network named Internet. Google Correlate [116], for example, enables one to find queries with a similar pattern to a target data series. The target can either be a real-world trend that one provides (eg, a dataset of event counts over time) or a query that one enters. In this context, Google Correlate uses Web search activity data to find queries with a similar pattern to a target data series. The results can be viewed on the Google Correlate website or downloaded as a comma-separated values (CSV) file [117] for further analysis. In other words, Google Correlate is like Google Trends in reverse. With Google Trends, one types in a query and receives a data series of activity. With Google Correlate, one enters a data series (the target) and receives a list of queries with a data series that follows a similar pattern.

These investigations should include the evolving impact of other social media, such as “Google+,” “Facebook” (eg, [118]), “YouTube” [119], “Wikipedia” [120], “Twitter” [121-124], and “IBM Watson” [125-127] on public health and reflect on the dark sides of the aforementioned developments, for example in terms of the possible impact of search engine analytics (on the companies behind them) on our privacy (eg, [128,129]). Second, the evolving contagiousity of ambient awareness, that is “awareness created through regular and constant reception, and/or exchange of information fragments through social media” (see [130]) has been neglected in public health thus far and could be a new form of “viral environment” for the upcoming generation. Finally, successful communication among health care providers and their patients from various sociocultural backgrounds depends on developing awareness of the normative cultural values of patients, how concepts of health and disease develop [131], and how these differ.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Supplementary tables S1 - S11.

[[PDF File \(Adobe PDF File\), 172KB-Multimedia Appendix 1](#)]

References

- Ferrari R, Shorter E. From railway spine to whiplash--the recycling of nervous irritation. *Med Sci Monit*. Nov 2003;9(11):HY27-HY37. [Medline: [14586283](#)]
- Ferrari R, Kwan O, Russell AS, Pearce JM, Schrader H. The best approach to the problem of whiplash? one ticket to Lithuania, please. *Clin Exp Rheumatol*. 1999;17(3):321-326. [Medline: [10410265](#)]
- Kwan O, Friel J. A new strategy to support the 'chronic injury' model of whiplash: ignore Lithuania. *Med Sci Monit*. Feb 2002;8(2):LE3-LE5. [Medline: [11913356](#)]
- Otte A. The 'railway spine'--a precursor for the 'whiplash syndrome'? *Med Sci Monit*. 2001;7(5):1064-1065. [Medline: [11535958](#)]
- Pearce JM. Whiplash injury: a reappraisal. *J Neurol Neurosurg Psychiatry*. Dec 1989;52(12):1329-1331. [[FREE Full text](#)] [Medline: [2693615](#)]
- Chappuis G, Soltermann B, CEA, AREDOC, CEREDOC. Number and cost of claims linked to minor cervical trauma in Europe: results from the comparative study by CEA, AREDOC and CEREDOC. *Eur Spine J*. Oct 2008;17(10):1350-1357. [[FREE Full text](#)] [doi: [10.1007/s00586-008-0732-8](#)] [Medline: [18704519](#)]
- European_Transport_Safety_Council. URL: <http://archive.etsc.eu/documents/ETS%20008-071.pdf> [accessed 2017-03-13] [[WebCite Cache ID 6ovquyV2j](#)]
- NCBI. 2016. URL: <https://www.ncbi.nlm.nih.gov/pubmed> [accessed 2016-11-27] [[WebCite Cache ID 6mL89pTOj](#)]
- Schrader H, Stovner LJ, Eisenmenger W. [Doubtful nosological validity of the chronic whiplash syndrome]. *Orthopade*. Feb 2012;41(2):147-152. [doi: [10.1007/s00132-011-1868-5](#)] [Medline: [22124728](#)]
- Schrader H, Obelieniene D, Bovim G, Surkiene D, Mickeviciene D, Miseviciene I, et al. Natural evolution of late whiplash syndrome outside the medicolegal context. *Lancet*. May 4, 1996;347(9010):1207-1211. [Medline: [8622449](#)]
- Cassidy JD, Carroll LJ, Côté P, Lemstra M, Berglund A, Nygren A. Effect of eliminating compensation for pain and suffering on the outcome of insurance claims for whiplash injury. *N Engl J Med*. Apr 20, 2000;342(16):1179-1186. [doi: [10.1056/NEJM200004203421606](#)] [Medline: [10770984](#)]
- Nuti SV, Wayda B, Ranasinghe I, Wang S, Dreyer RP, Chen SI, et al. The use of google trends in health care research: a systematic review. *PLoS One*. 2014;9(10):e109583. [[FREE Full text](#)] [doi: [10.1371/journal.pone.0109583](#)] [Medline: [25337815](#)]
- Jansen BJ. Search log analysis: what it is, what's been done, how to do it. *Libr Inf Sci Res*. Sep 2006;28(3):407-432. [doi: [10.1016/j.lisr.2006.06.005](#)]
- Eysenbach G. Infodemiology: tracking flu-related searches on the web for syndromic surveillance. *AMIA Annu Symp Proc*. 2006.:244-248. [[FREE Full text](#)] [Medline: [17238340](#)]
- Eysenbach G. Infodemiology and infoveillance: framework for an emerging set of public health informatics methods to analyze search, communication and publication behavior on the Internet. *J Med Internet Res*. 2009;11(1):e11. [[FREE Full text](#)] [doi: [10.2196/jmir.1157](#)] [Medline: [19329408](#)]
- Wikipedia. URL: https://en.wikipedia.org/wiki/Search_analytics [accessed 2016-11-27] [[WebCite Cache ID 6mL7bZr0h](#)]
- Internet World Stats. URL: <http://www.internetworldstats.com/stats.htm> [accessed 2016-11-27] [[WebCite Cache ID 6mL7THJEx](#)]
- Santana S, Lausen B, Bujnowska-Fedak M, Chronaki CE, Prokosch H, Wynn R. Informed citizen and empowered citizen in health: results from an European survey. *BMC Fam Pract*. 2011;12:20. [[FREE Full text](#)] [doi: [10.1186/1471-2296-12-20](#)] [Medline: [21496309](#)]
- Cross M. How the internet is changing health care. *BMJ*. Jul 22, 2008;337:a883. [Medline: [18647776](#)]
- Dufton JA, Bruni SG, Kopec JA, Cassidy JD, Quon J. Delayed recovery in patients with whiplash-associated disorders. *Injury*. Jul 2012;43(7):1141-1147. [doi: [10.1016/j.injury.2012.03.006](#)] [Medline: [22475071](#)]
- Noll-Hussong M, Lahmann C. [Whiplash and Werther effect: the potential of google insights for search for medical research and public health]. *Fortschr Neurol Psychiatr*. Jun 2011;79(6):340-344. [doi: [10.1055/s-0031-1273293](#)] [Medline: [21598205](#)]
- Ginsberg J, Mohebbi MH, Patel RS, Brammer L, Smolinski MS, Brilliant L. Detecting influenza epidemics using search engine query data. *Nature*. Feb 19, 2009;457(7232):1012-1014. [doi: [10.1038/nature07634](#)] [Medline: [19020500](#)]
- Breyer BN, Eisenberg ML. Use of Google in study of noninfectious medical conditions. *Epidemiology*. Jul 2010;21(4):584-585. [[FREE Full text](#)] [doi: [10.1097/EDE.0b013e3181e080cd](#)] [Medline: [20539114](#)]
- Rossignol L, Pelat C, Lambert B, Flahault A, Chartier-Kastler E, Hanslik T. A method to assess seasonality of urinary tract infections based on medication sales and google trends. *PLoS One*. 2013;8(10):e76020. [[FREE Full text](#)] [doi: [10.1371/journal.pone.0076020](#)] [Medline: [24204587](#)]

25. Nguyen T, Tran T, Luo W, Gupta S, Rana S, Phung D, et al. Web search activity data accurately predict population chronic disease risk in the USA. *J Epidemiol Community Health*. Jul 2015;69(7):693-699. [doi: [10.1136/jech-2014-204523](https://doi.org/10.1136/jech-2014-204523)] [Medline: [25805603](https://pubmed.ncbi.nlm.nih.gov/25805603/)]
26. Ling R, Lee J. Disease monitoring and health campaign evaluation using Google search activities for HIV and AIDS, stroke, colorectal cancer, and marijuana use in Canada: a retrospective observational study. *JMIR Public Health Surveill*. Oct 12, 2016;2(2):e156. [FREE Full text] [doi: [10.2196/publichealth.6504](https://doi.org/10.2196/publichealth.6504)] [Medline: [27733330](https://pubmed.ncbi.nlm.nih.gov/27733330/)]
27. Kelly CM, Jorm AF. Stigma and mood disorders. *Curr Opin Psychiatry*. Jan 2007;20(1):13-16. [doi: [10.1097/YCO.0b013e3280113cf5](https://doi.org/10.1097/YCO.0b013e3280113cf5)] [Medline: [17143076](https://pubmed.ncbi.nlm.nih.gov/17143076/)]
28. Elbers NA, Hulst L, Cuijpers P, Akkermans AJ, Bruinvels DJ. Do compensation processes impair mental health? A meta-analysis. *Injury*. May 2013;44(5):674-683. [FREE Full text] [doi: [10.1016/j.injury.2011.11.025](https://doi.org/10.1016/j.injury.2011.11.025)] [Medline: [22244996](https://pubmed.ncbi.nlm.nih.gov/22244996/)]
29. Fishbain DA, Rosomoff HL, Cutler RB, Rosomoff RS. Secondary gain concept: a review of the scientific evidence. *Clin J Pain*. Mar 1995;11(1):6-21. [Medline: [7787338](https://pubmed.ncbi.nlm.nih.gov/7787338/)]
30. Bernardo TM, Rajic A, Young I, Robiadek K, Pham MT, Funk JA. Scoping review on search queries and social media for disease surveillance: a chronology of innovation. *J Med Internet Res*. 2013;15(7):e147. [FREE Full text] [doi: [10.2196/jmir.2740](https://doi.org/10.2196/jmir.2740)] [Medline: [23896182](https://pubmed.ncbi.nlm.nih.gov/23896182/)]
31. Alexa. 2016. URL: <http://www.alexa.com/topsites> [accessed 2016-11-27] [WebCite Cache ID 6mL7YX2rk]
32. InternetLiveStats. 2016. URL: <http://www.internetlivestats.com/google-search-statistics/> [accessed 2016-11-27] [WebCite Cache ID 6mL7yKNrd]
33. Marx P. [Assessment of whiplash and cervical spine injury]. *Nervenarzt*. Dec 2011;82(12):1525-1532. [doi: [10.1007/s00115-011-3286-7](https://doi.org/10.1007/s00115-011-3286-7)] [Medline: [22105883](https://pubmed.ncbi.nlm.nih.gov/22105883/)]
34. Miettinen T, Lindgren KA, Airaksinen O, Leino E. Whiplash injuries in Finland: a prospective 1-year follow-up study. *Clin Exp Rheumatol*. 2002;20(3):399-402. [Medline: [12102479](https://pubmed.ncbi.nlm.nih.gov/12102479/)]
35. Obelieniene D, Schrader H, Bovim G, Miseviciene I, Sand T. Pain after whiplash: a prospective controlled inception cohort study. *J Neurol Neurosurg Psychiatry*. Mar 1999;66(3):279-283. [FREE Full text] [Medline: [10084524](https://pubmed.ncbi.nlm.nih.gov/10084524/)]
36. Haneline MT. The notion of a 'whiplash culture': a review of the evidence. *J Chiropr Med*. Sep 2009;8(3):119-124. [FREE Full text] [doi: [10.1016/j.jcm.2009.04.001](https://doi.org/10.1016/j.jcm.2009.04.001)] [Medline: [19703667](https://pubmed.ncbi.nlm.nih.gov/19703667/)]
37. Richter M, Ferrari R, Otte D, Kuensebeck H, Blauth M, Krettek C. Correlation of clinical findings, collision parameters, and psychological factors in the outcome of whiplash associated disorders. *J Neurol Neurosurg Psychiatry*. May 2004;75(5):758-764. [FREE Full text] [Medline: [15090574](https://pubmed.ncbi.nlm.nih.gov/15090574/)]
38. Castro WH, Meyer SJ, Becke ME, Nentwig CG, Hein MF, Ercan BI, et al. No stress--no whiplash? Prevalence of 'whiplash' symptoms following exposure to a placebo rear-end collision. *Int J Legal Med*. 2001;114(6):316-322. [Medline: [11508796](https://pubmed.ncbi.nlm.nih.gov/11508796/)]
39. Atherton K, Wiles NJ, Lecky FE, Hawes SJ, Silman AJ, Macfarlane GJ, et al. Predictors of persistent neck pain after whiplash injury. *Emerg Med J*. Mar 2006;23(3):195-201. [FREE Full text] [doi: [10.1136/emj.2005.027102](https://doi.org/10.1136/emj.2005.027102)] [Medline: [16498156](https://pubmed.ncbi.nlm.nih.gov/16498156/)]
40. Schmand B, Lindeboom J, Schagen S, Heijt R, Koene T, Hamburger HL. Cognitive complaints in patients after whiplash injury: the impact of malingering. *J Neurol Neurosurg Psychiatry*. Mar 1998;64(3):339-343. [FREE Full text] [Medline: [9527145](https://pubmed.ncbi.nlm.nih.gov/9527145/)]
41. Regenauer A. [Contentious diseases--a medico-social phenomenon from an insurance medicine perspective]. *Versicherungsmedizin*. Mar 1, 2008;60(1):3-7. [Medline: [18405228](https://pubmed.ncbi.nlm.nih.gov/18405228/)]
42. Jaracz J, Gattner K, Jaracz K, Górna K. Unexplained painful physical symptoms in patients with major depressive disorder: prevalence, pathophysiology and management. *CNS Drugs*. Apr 2016;30(4):293-304. [FREE Full text] [doi: [10.1007/s40263-016-0328-5](https://doi.org/10.1007/s40263-016-0328-5)] [Medline: [27048351](https://pubmed.ncbi.nlm.nih.gov/27048351/)]
43. Tribunal federal Switzerland V. Servat. URL: http://www.servat.unibe.ch/dfr/bger/100830_9C_510-2009.html [accessed 2017-03-13] [WebCite Cache ID 6ovvHHKsc]
44. Tobón GJ, Youinou P, Saraux A. The environment, geo-epidemiology, and autoimmune disease: rheumatoid arthritis. *J Autoimmun*. Aug 2010;35(1):10-14. [doi: [10.1016/j.jaut.2009.12.009](https://doi.org/10.1016/j.jaut.2009.12.009)] [Medline: [20080387](https://pubmed.ncbi.nlm.nih.gov/20080387/)]
45. Stovner LJ, Andree C. Prevalence of headache in Europe: a review for the Eurolight project. *J Headache Pain*. Aug 2010;11(4):289-299. [FREE Full text] [doi: [10.1007/s10194-010-0217-0](https://doi.org/10.1007/s10194-010-0217-0)] [Medline: [20473702](https://pubmed.ncbi.nlm.nih.gov/20473702/)]
46. Hartholt KA, Oudshoorn C, Zielinski SM, Burgers PT, Panneman MJ, van Beeck EF, et al. The epidemic of hip fractures: are we on the right track? *PLoS One*. 2011;6(7):e22227. [FREE Full text] [doi: [10.1371/journal.pone.0022227](https://doi.org/10.1371/journal.pone.0022227)] [Medline: [21799797](https://pubmed.ncbi.nlm.nih.gov/21799797/)]
47. Court-Brown CM, Wood AM, Aitken S. The epidemiology of acute sports-related fractures in adults. *Injury*. Dec 2008;39(12):1365-1372. [doi: [10.1016/j.injury.2008.02.004](https://doi.org/10.1016/j.injury.2008.02.004)] [Medline: [18514656](https://pubmed.ncbi.nlm.nih.gov/18514656/)]
48. Richter M, Otte D, Jahanyar K, Blauth M. Upper extremity fractures in restrained front-seat occupants. *J Trauma*. May 2000;48(5):907-912. [Medline: [10823535](https://pubmed.ncbi.nlm.nih.gov/10823535/)]
49. O'Neill TW, Cooper C, Finn JD, Lunt M, Purdie D, Reid DM, et al. UK Colles' Fracture Study Group. Incidence of distal forearm fracture in British men and women. *Osteoporos Int*. 2001;12(7):555-558. [Medline: [11527052](https://pubmed.ncbi.nlm.nih.gov/11527052/)]

50. Thompson PW, Taylor J, Dawson A. The annual incidence and seasonal variation of fractures of the distal radius in men and women over 25 years in Dorset, UK. *Injury*. May 2004;35(5):462-466. [doi: [10.1016/S0020-1383\(03\)00117-7](https://doi.org/10.1016/S0020-1383(03)00117-7)] [Medline: [15081322](#)]
51. Cheng SY, Levy AR, Lefavre KA, Guy P, Kuramoto L, Sobolev B. Geographic trends in incidence of hip fractures: a comprehensive literature review. *Osteoporos Int*. Oct 2011;22(10):2575-2586. [doi: [10.1007/s00198-011-1596-z](https://doi.org/10.1007/s00198-011-1596-z)] [Medline: [21484361](#)]
52. Wittchen HU, Jacobi F, Rehm J, Gustavsson A, Svensson M, Jönsson B, et al. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol*. Sep 2011;21(9):655-679. [doi: [10.1016/j.euroneuro.2011.07.018](https://doi.org/10.1016/j.euroneuro.2011.07.018)] [Medline: [21896369](#)]
53. Gustavsson A, Svensson M, Jacobi F, Allgulander C, Alonso J, Beghi E, et al. CDBE2010Study Group. Cost of disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol*. Oct 2011;21(10):718-779. [doi: [10.1016/j.euroneuro.2011.08.008](https://doi.org/10.1016/j.euroneuro.2011.08.008)] [Medline: [21924589](#)]
54. Olesen J, Gustavsson A, Svensson M, Wittchen H, Jönsson B, CDBE2010 SG, et al. The economic cost of brain disorders in Europe. *Eur J Neurol*. Jan 2012;19(1):155-162. [doi: [10.1111/j.1468-1331.2011.03590.x](https://doi.org/10.1111/j.1468-1331.2011.03590.x)] [Medline: [22175760](#)]
55. WHO. 2016. URL: <http://www.who.int/mediacentre/factsheets/fs369/en/> [accessed 2016-11-27] [WebCite Cache ID [6mL8FwmIj](#)]
56. Branco JC, Bannwarth B, Failde I, Abello CJ, Blotman F, Spaeth M, et al. Prevalence of fibromyalgia: a survey in five European countries. *Semin Arthritis Rheum*. Jun 2010;39(6):448-453. [doi: [10.1016/j.semarthrit.2008.12.003](https://doi.org/10.1016/j.semarthrit.2008.12.003)] [Medline: [19250656](#)]
57. Neumann L, Buskila D. Epidemiology of fibromyalgia. *Curr Pain Headache Rep*. Oct 2003;7(5):362-368. [Medline: [12946289](#)]
58. de Girolamo G. Epidemiology and social costs of low back pain and fibromyalgia. *Clin J Pain*. 1991;7 Suppl 1:S1-S7. [Medline: [1839774](#)]
59. Tzellos TG, Toulis KA, Goulis DG, Papazisis G, Zampeli VA, Vakfari A, et al. Gabapentin and pregabalin in the treatment of fibromyalgia: a systematic review and a meta-analysis. *J Clin Pharm Ther*. Dec 2010;35(6):639-656. [doi: [10.1111/j.1365-2710.2009.01144.x](https://doi.org/10.1111/j.1365-2710.2009.01144.x)] [Medline: [21054455](#)]
60. Mäkelä M, Heliövaara M. Prevalence of primary fibromyalgia in the Finnish population. *BMJ*. Jul 27, 1991;303(6796):216-219. [FREE Full text] [Medline: [1884057](#)]
61. Mozilla. 2016. URL: <https://www.mozilla.org/en-US/> [accessed 2016-11-27] [WebCite Cache ID [6mL84Z59b](#)]
62. Google. 2012. URL: <https://www.google.com/trends/> [accessed 2016-11-27] [WebCite Cache ID [6mL7IS3L2](#)]
63. StatCounter. 2016. URL: <http://gs.statcounter.com/> [accessed 2016-11-27] [WebCite Cache ID [6mL8COPyd](#)]
64. Alexa. 1996. URL: <http://www.alexa.com/> [accessed 2016-11-27] [WebCite Cache ID [6mL7enmau](#)]
65. Google. 2016. URL: https://support.google.com/trends/answer/6248105?hl=en&ref_topic=6248052,%2027Jul2015 [accessed 2015-07-27] [WebCite Cache ID [6mL7rpv1n](#)]
66. Singh PM, Wight CA, Sercinoglu O, Wilson DC, Boytsov A, Raizada MN. Language preferences on websites and in Google searches for human health and food information. *J Med Internet Res*. Jun 28, 2007;9(2):e18. [FREE Full text] [doi: [10.2196/jmir.9.2.e18](https://doi.org/10.2196/jmir.9.2.e18)] [Medline: [17613488](#)]
67. Metcalfe D, Price C, Powell J. Media coverage and public reaction to a celebrity cancer diagnosis. *J Public Health (Oxf)*. Mar 2011;33(1):80-85. [FREE Full text] [doi: [10.1093/pubmed/fdq052](https://doi.org/10.1093/pubmed/fdq052)] [Medline: [20679285](#)]
68. Eysenbach G. Infodemiology: tracking flu-related searches on the web for syndromic surveillance. *AMIA Annu Symp Proc*. 2006.;244-248. [FREE Full text] [Medline: [17238340](#)]
69. Wikipedia. 2016. URL: https://de.wikipedia.org/wiki/Google_Insights_for_Search [accessed 2016-11-27] [WebCite Cache ID [6mL8Y0psS](#)]
70. Shimshoni Y, Efron N, Matias Y, Wellesley. 2009. URL: http://cs.wellesley.edu/~cs315/Papers/Matias%20et%20al%20google_trends_predictability.pdf [accessed 2017-03-13] [WebCite Cache ID [6ovtMTfPO](#)]
71. Youtube. 2016. URL: https://www.youtube.com/results?search_query=whiplash+injury [accessed 2016-11-27] [WebCite Cache ID [6mL8SjALf](#)]
72. Ferrari R. The Whiplash Encyclopedia: The Facts and Myths of Whiplash. Sudbury, MA. Jones and Bartlett Publishers; 2006.
73. Ferrari R, Pieschl S. An examination of coping styles and expectations for whiplash injury in Germany: comparison with Canadian data. *Clin Rheumatol*. Sep 2011;30(9):1209-1214. [doi: [10.1007/s10067-011-1736-z](https://doi.org/10.1007/s10067-011-1736-z)] [Medline: [21448615](#)]
74. Benoist M, Rouaud J. Whiplash: myth or reality? *Joint Bone Spine*. Jun 2002;69(4):358-362. [Medline: [12184430](#)]
75. Ferrari R, Constantoyannis C, Papadakis N. Laypersons' expectation of the sequelae of whiplash injury: a cross-cultural comparative study between Canada and Greece. *Med Sci Monit*. Mar 2003;9(3):CR120-CR124. [Medline: [12640340](#)]
76. Partheni M, Constantoyannis C, Ferrari R, Nikiforidis G, Voulgaris S, Papadakis N. A prospective cohort study of the outcome of acute whiplash injury in Greece. *Clin Exp Rheumatol*. 2000;18(1):67-70. [Medline: [10728446](#)]
77. Bannister G, Amirfeyz R, Kelley S, Gargan M. Whiplash injury. *J Bone Joint Surg Br*. Jul 2009;91(7):845-850. [doi: [10.1302/0301-620X.91B7.22639](https://doi.org/10.1302/0301-620X.91B7.22639)] [Medline: [19567844](#)]

78. Ferrari R, Schrader H. The late whiplash syndrome: a biopsychosocial approach. *J Neurol Neurosurg Psychiatry*. Jun 2001;70(6):722-726. [FREE Full text] [Medline: [11385003](#)]
79. Wenzel HG, Mykletun A, Nilsen TI. Symptom profile of persons self-reporting whiplash: a Norwegian population-based study (HUNT 2). *Eur Spine J*. Sep 2009;18(9):1363-1370. [FREE Full text] [doi: [10.1007/s00586-009-1106-6](#)] [Medline: [19669172](#)]
80. Holm LW, Carroll LJ, Cassidy JD, Hogg-Johnson S, Côté P, Guzman J, et al. BoneJoint Decade 2000-2010 Task Force on Neck Pain/Its Associated Disorders. The burden and determinants of neck pain in whiplash-associated disorders after traffic collisions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)*. Feb 15, 2008;33(4 Suppl):S52-S59. [doi: [10.1097/BRS.0b013e3181643ece](#)] [Medline: [18204401](#)]
81. McLean SA, Clauw DJ. Predicting chronic symptoms after an acute 'stressor'--lessons learned from 3 medical conditions. *Med Hypotheses*. 2004;63(4):653-658. [doi: [10.1016/j.mehy.2004.03.022](#)] [Medline: [15325010](#)]
82. Ortiz JR, Zhou H, Shay DK, Neuzil KM, Fowlkes AL, Goss CH. Monitoring influenza activity in the United States: a comparison of traditional surveillance systems with Google Flu Trends. *PLoS One*. Apr 2011;6(4):e18687. [FREE Full text] [doi: [10.1371/journal.pone.0018687](#)] [Medline: [21556151](#)]
83. Corley CD, Cook DJ, Mikler AR, Singh KP. Using Web and social media for influenza surveillance. *Adv Exp Med Biol*. 2010;680:559-564. [doi: [10.1007/978-1-4419-5913-3_61](#)] [Medline: [20865540](#)]
84. Althouse BM, Ng YY, Cummings DA. Prediction of dengue incidence using search query surveillance. *PLoS Negl Trop Dis*. Aug 2011;5(8):e1258. [FREE Full text] [doi: [10.1371/journal.pntd.0001258](#)] [Medline: [21829744](#)]
85. Eysenbach G. Infodemiology and infoveillance: framework for an emerging set of public health informatics methods to analyze search, communication and publication behavior on the Internet. *J Med Internet Res*. 2009;11(1):e11. [FREE Full text] [doi: [10.2196/jmir.1157](#)] [Medline: [19329408](#)]
86. Yang AC, Huang NE, Peng C, Tsai S. Do seasons have an influence on the incidence of depression? the use of an internet search engine query data as a proxy of human affect. *PLoS One*. Oct 28, 2010;5(10):e13728. [FREE Full text] [doi: [10.1371/journal.pone.0013728](#)] [Medline: [21060851](#)]
87. Breyer BN, Sen S, Aaronson DS, Stoller ML, Erickson BA, Eisenberg ML. Use of Google Insights for Search to track seasonal and geographic kidney stone incidence in the United States. *Urology*. Aug 2011;78(2):267-271. [FREE Full text] [doi: [10.1016/j.urology.2011.01.010](#)] [Medline: [21459414](#)]
88. Wang L, Wang J, Wang M, Li Y, Liang Y, Xu D. Using internet search engines to obtain medical information: a comparative study. *J Med Internet Res*. 2012;14(3):e74. [FREE Full text] [doi: [10.2196/jmir.1943](#)] [Medline: [22672889](#)]
89. Krause J, Ruxton GD, Krause S. Swarm intelligence in animals and humans. *Trends Ecol Evol*. Jan 2010;25(1):28-34. [doi: [10.1016/j.tree.2009.06.016](#)] [Medline: [19735961](#)]
90. Christakis NA, Fowler JH. Social network sensors for early detection of contagious outbreaks. *PLoS One*. 2010;5(9):e12948. [FREE Full text] [doi: [10.1371/journal.pone.0012948](#)] [Medline: [20856792](#)]
91. Brownstein JS, Freifeld CC, Madoff LC. Digital disease detection--harnessing the Web for public health surveillance. *N Engl J Med*. May 21, 2009;360(21):2153-5, 2157. [FREE Full text] [doi: [10.1056/NEJMp0900702](#)] [Medline: [19423867](#)]
92. Fava GA, Guidi J. Information overload, the patient and the clinician. *Psychother Psychosom*. 2007;76(1):1-3. [doi: [10.1159/000096358](#)] [Medline: [17170558](#)]
93. Van De Belt TH, Engelen LJ, Berben SA, Schoonhoven L. Definition of Health 2.0 and Medicine 2.0: a systematic review. *J Med Internet Res*. 2010;12(2):e18. [FREE Full text] [doi: [10.2196/jmir.1350](#)] [Medline: [20542857](#)]
94. Bremner JD, Quinn J, Quinn W, Veledar E. Surfing the net for medical information about psychological trauma: an empirical study of the quality and accuracy of trauma-related websites. *Med Inform Internet Med*. Sep 2006;31(3):227-236. [FREE Full text] [doi: [10.1080/14639230600887866](#)] [Medline: [16954059](#)]
95. Butler L, Foster NE. Back pain online: a cross-sectional survey of the quality of web-based information on low back pain. *Spine (Phila Pa 1976)*. Feb 15, 2003;28(4):395-401. [doi: [10.1097/01.BRS.0000048497.38319.D3](#)] [Medline: [12590218](#)]
96. Eysenbach G. Infodemiology: the epidemiology of (mis)information. *Am J Med*. Dec 15, 2002;113(9):763-765. [Medline: [12517369](#)]
97. Dong X, Bollen J. Computational models of consumer confidence from large-scale online attention data: crowd-sourcing econometrics. *PLoS One*. 2015;10(3):e0120039. [FREE Full text] [doi: [10.1371/journal.pone.0120039](#)] [Medline: [25826692](#)]
98. Pariser E. *The Filter Bubble*. London. Penguin Books Ltd; 2012.
99. Hermeking M. Culture and internet consumption: contributions from cross-cultural marketing and advertising research. *J Comp Mediated Comm*. Nov 2005;11(1):192-216. [doi: [10.1111/j.1083-6101.2006.tb00310.x](#)]
100. Stuart D. Webometrics. 2008. URL: <http://blog.webometrics.org.uk/2008/09/google-insights-for-search-term-order-is-all-important/> [accessed 2017-03-13] [WebCite Cache ID 6ovu3t4ZL]
101. Cho S, Sohn CH, Jo MW, Shin S, Lee JH, Ryoo SM, et al. Correlation between national influenza surveillance data and google trends in South Korea. *PLoS One*. 2013;8(12):e81422. [FREE Full text] [doi: [10.1371/journal.pone.0081422](#)] [Medline: [24339927](#)]
102. Kang M, Zhong H, He J, Rutherford S, Yang F. Using Google trends for influenza surveillance in South China. *PLoS One*. 2013;8(1):e55205. [FREE Full text] [doi: [10.1371/journal.pone.0055205](#)] [Medline: [23372837](#)]

103. Domnich A, Panatto D, Signori A, Lai PL, Gasparini R, Amicizia D. Age-related differences in the accuracy of web query-based predictions of influenza-like illness. *PLoS One*. 2015;10(5):e0127754. [FREE Full text] [doi: [10.1371/journal.pone.0127754](https://doi.org/10.1371/journal.pone.0127754)] [Medline: [26011418](https://pubmed.ncbi.nlm.nih.gov/26011418/)]
104. Salek KM, Mamun MA, Parvin N, Ahmed SM, Khan MM, Rijvi AN, et al. Fluctuation of pain by weather change in musculoskeletal disorders. *Mymensingh Med J*. Oct 2011;20(4):645-651. [Medline: [22081184](https://pubmed.ncbi.nlm.nih.gov/22081184/)]
105. Olson DR, Konty KJ, Paladini M, Viboud C, Simonsen L. Reassessing Google flu trends data for detection of seasonal and pandemic influenza: a comparative epidemiological study at three geographic scales. *PLoS Comput Biol*. 2013;9(10):e1003256. [FREE Full text] [doi: [10.1371/journal.pcbi.1003256](https://doi.org/10.1371/journal.pcbi.1003256)] [Medline: [24146603](https://pubmed.ncbi.nlm.nih.gov/24146603/)]
106. Cook S, Conrad C, Fowlkes AL, Mohebbi MH. Assessing Google flu trends performance in the United States during the 2009 influenza virus A (H1N1) pandemic. *PLoS One*. 2011;6(8):e23610. [FREE Full text] [doi: [10.1371/journal.pone.0023610](https://doi.org/10.1371/journal.pone.0023610)] [Medline: [21886802](https://pubmed.ncbi.nlm.nih.gov/21886802/)]
107. Butler D. When Google got flu wrong. *Nature*. Feb 14, 2013;494(7436):155-156. [doi: [10.1038/494155a](https://doi.org/10.1038/494155a)] [Medline: [23407515](https://pubmed.ncbi.nlm.nih.gov/23407515/)]
108. Brockwell P, Davis R. *Introduction to Time Series and Forecasting* (Springer Texts in Statistics). New York. Springer-Verlag; 2016.
109. Chatfield C. *The Analysis of Time Series: An Introduction*, Sixth Edition. London. Chapman & Hall/CRC; 2003.
110. Eysenbach G, Diepgen TL. Patients looking for information on the Internet and seeking teleadvice: motivation, expectations, and misconceptions as expressed in e-mails sent to physicians. *Arch Dermatol*. Feb 1999;135(2):151-156. [Medline: [10052399](https://pubmed.ncbi.nlm.nih.gov/10052399/)]
111. Iverson SA, Howard KB, Penney BK. Impact of internet use on health-related behaviors and the patient-physician relationship: a survey-based study and review. *J Am Osteopath Assoc*. Dec 2008;108(12):699-711. [Medline: [19075034](https://pubmed.ncbi.nlm.nih.gov/19075034/)]
112. Muse K, McManus F, Leung C, Meghreblian B, Williams JM. Cyberchondriasis: fact or fiction? A preliminary examination of the relationship between health anxiety and searching for health information on the Internet. *J Anxiety Disord*. Jan 2012;26(1):189-196. [doi: [10.1016/j.janxdis.2011.11.005](https://doi.org/10.1016/j.janxdis.2011.11.005)] [Medline: [22137465](https://pubmed.ncbi.nlm.nih.gov/22137465/)]
113. Eastin MS, Guinsler NM. Worried and wired: effects of health anxiety on information-seeking and health care utilization behaviors. *Cyberpsychol Behav*. Aug 2006;9(4):494-498. [doi: [10.1089/cpb.2006.9.494](https://doi.org/10.1089/cpb.2006.9.494)] [Medline: [16901253](https://pubmed.ncbi.nlm.nih.gov/16901253/)]
114. Eysenbach G, Thomson M. The FA4CT algorithm: a new model and tool for consumers to assess and filter health information on the Internet. *Stud Health Technol Inform*. 2007;129(Pt 1):142-146. [Medline: [17911695](https://pubmed.ncbi.nlm.nih.gov/17911695/)]
115. Eysenbach G. Infodemiology and infoveillance tracking online health information and cyberbehavior for public health. *Am J Prev Med*. May 2011;40(5 Suppl 2):S154-S158. [doi: [10.1016/j.amepre.2011.02.006](https://doi.org/10.1016/j.amepre.2011.02.006)] [Medline: [21521589](https://pubmed.ncbi.nlm.nih.gov/21521589/)]
116. Google. 2016. URL: <https://www.google.com/trends/correlate/> [accessed 2016-11-27] [WebCite Cache ID 6mL7ukf14]
117. Wikipedia. 2016. URL: https://en.wikipedia.org/wiki/Comma-separated_values [accessed 2016-11-27] [WebCite Cache ID 6mL8PZwrH]
118. Moreno MA, Christakis DA, Egan KG, Brockman LN, Becker T. Associations between displayed alcohol references on Facebook and problem drinking among college students. *Arch Pediatr Adolesc Med*. Feb 2012;166(2):157-163. [FREE Full text] [doi: [10.1001/archpediatrics.2011.180](https://doi.org/10.1001/archpediatrics.2011.180)] [Medline: [21969360](https://pubmed.ncbi.nlm.nih.gov/21969360/)]
119. Stamelou M, Edwards MJ, Espay AJ, Fung VS, Hallett M, Lang AE, et al. Movement disorders on YouTube--caveat spectator. *N Engl J Med*. Sep 22, 2011;365(12):1160-1161. [FREE Full text] [doi: [10.1056/NEJMc1107673](https://doi.org/10.1056/NEJMc1107673)] [Medline: [21992142](https://pubmed.ncbi.nlm.nih.gov/21992142/)]
120. McIver DJ, Brownstein JS. Wikipedia usage estimates prevalence of influenza-like illness in the United States in near real-time. *PLoS Comput Biol*. Apr 2014;10(4):e1003581. [FREE Full text] [doi: [10.1371/journal.pcbi.1003581](https://doi.org/10.1371/journal.pcbi.1003581)] [Medline: [24743682](https://pubmed.ncbi.nlm.nih.gov/24743682/)]
121. Chew C, Eysenbach G. Pandemics in the age of twitter: content analysis of tweets during the 2009 H1N1 outbreak. *PLoS One*. 2010;5(11):e14118. [FREE Full text] [doi: [10.1371/journal.pone.0014118](https://doi.org/10.1371/journal.pone.0014118)] [Medline: [21124761](https://pubmed.ncbi.nlm.nih.gov/21124761/)]
122. Heavilin N, Gerbert B, Page JE, Gibbs JL. Public health surveillance of dental pain via Twitter. *J Dent Res*. Sep 2011;90(9):1047-1051. [FREE Full text] [doi: [10.1177/0022034511415273](https://doi.org/10.1177/0022034511415273)] [Medline: [21768306](https://pubmed.ncbi.nlm.nih.gov/21768306/)]
123. Paul MJ, Dredze M, Broniatowski D. Twitter improves influenza forecasting. *PLoS Curr*. 2014;6 [FREE Full text] [doi: [10.1371/currents.outbreaks.90b9ed0f59bae4ccaa683a39865d9117](https://doi.org/10.1371/currents.outbreaks.90b9ed0f59bae4ccaa683a39865d9117)] [Medline: [25642377](https://pubmed.ncbi.nlm.nih.gov/25642377/)]
124. Weeg C, Schwartz HA, Hill S, Merchant RM, Arango C, Ungar L. Using Twitter to Measure Public Discussion of Diseases: A Case Study. *JMIR Public Health Surveill*. 2015;1(1):e6. [FREE Full text] [doi: [10.2196/publichealth.3953](https://doi.org/10.2196/publichealth.3953)] [Medline: [26925459](https://pubmed.ncbi.nlm.nih.gov/26925459/)]
125. Chen Y, Elene A, Weber G. IBM Watson: how cognitive computing can be applied to big data challenges in life sciences research. *Clin Ther*. Apr 2016;38(4):688-701. [FREE Full text] [doi: [10.1016/j.clinthera.2015.12.001](https://doi.org/10.1016/j.clinthera.2015.12.001)] [Medline: [27130797](https://pubmed.ncbi.nlm.nih.gov/27130797/)]
126. Hoyt RE, Snider D, Thompson C, Mantravadi S. IBM Watson analytics: automating visualization, descriptive, and predictive statistics. *JMIR Public Health Surveill*. Oct 11, 2016;2(2):e157. [FREE Full text] [doi: [10.2196/publichealth.5810](https://doi.org/10.2196/publichealth.5810)] [Medline: [27729304](https://pubmed.ncbi.nlm.nih.gov/27729304/)]
127. Tang PC, Smith MD. Democratization of Health Care. *JAMA*. Oct 25, 2016;316(16):1663-1664. [doi: [10.1001/jama.2016.14060](https://doi.org/10.1001/jama.2016.14060)] [Medline: [27669345](https://pubmed.ncbi.nlm.nih.gov/27669345/)]

128. Greist J. The Internet: every good thing has a dark side. *World Psychiatry*. Jun 2010;9(2):95-96. [[FREE Full text](#)] [Medline: [20671895](#)]
129. Nelson SB. Privacy and medical information on the Internet. *Respir Care*. Feb 2006;51(2):183-187. [[FREE Full text](#)] [Medline: [16441962](#)]
130. Wikipeda. 2016. URL: https://en.wikipedia.org/wiki/Ambient_awareness [accessed 2016-11-27] [[WebCite Cache ID 6mL8KDC42](#)]
131. Boruchovitch E, Mednick BR. Cross-cultural differences in children's concepts of health and illness. *Rev Saude Publica*. Oct 1997;31(5):448-456. [Medline: [9629721](#)]

Abbreviations

CSV: comma-separated values
FAQ: frequently asked questions
GIS: Google Insights for Search
IP: Internet protocol

Edited by G Eysenbach; submitted 27.11.16; peer-reviewed by R Ferrari, D Snider; comments to author 05.01.17; revised version received 22.01.17; accepted 20.02.17; published 27.03.17

Please cite as:

Noll-Hussong M

Whiplash Syndrome Reloaded: Digital Echoes of Whiplash Syndrome in the European Internet Search Engine Context

JMIR Public Health Surveill 2017;3(1):e15

URL: <http://publichealth.jmir.org/2017/1/e15/>

doi: [10.2196/publichealth.7054](https://doi.org/10.2196/publichealth.7054)

PMID: [28347974](https://pubmed.ncbi.nlm.nih.gov/28347974/)

©Michael Noll-Hussong. Originally published in JMIR Public Health and Surveillance (<http://publichealth.jmir.org>), 27.03.2017. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.