

Letter to the Editor

Effectivity of Awareness Months in Increasing Internet Search Activity for Top Malignancies Among Women

Dhruvika Mukhija¹, MD; Anand Venkatraman², MD; Sajan Jiv Singh Nagpal³, MD

¹Cleveland Clinic, Internal Medicine, Cleveland, OH, United States

²University of Alabama, Neurology, Birmingham, AL, United States

³Mayo Clinic, Rochester, MN, United States

Corresponding Author:

Sajan Jiv Singh Nagpal, MD

Mayo Clinic

200 First St SW

Mayo 9E

Rochester, MN, 55905

United States

Phone: 1 5073191959

Fax: 1 (844) 668 6880

Email: nagpal.sajan@mayo.edu

Related Article:

Comment on: <http://www.publichealth.jmir.org/2016/2/e156/>

(*JMIR Public Health Surveill* 2017;3(3):e55) doi: [10.2196/publichealth.7714](https://doi.org/10.2196/publichealth.7714)

KEYWORDS

colorectal cancer, lung cancer, breast cancer, cancer awareness month, infoveillance

Letter

In a recent article, Ling et al. hypothesized that following the launch of a campaign for a medical condition, information seeking behavior pertaining to the condition would increase as well [1]. They used data from Google Trends (Google Inc., CA) on 4 different diseases (including Colon Cancer) to conclude that the use of infoveillance (type of public health surveillance based on online content analysis), shows promise as an alternative and inexpensive solution for disease surveillance and health care campaign evaluation. While a number of health campaigns are rolled out by the government and professional societies, there are limited means and tools to evaluate the effectivity of these campaigns. Using infoveillance, the ‘digital impact’ of various health- and disease-related campaigns can be assessed.

Cancer awareness has massively benefited from rapid growth of internet and mass media and the evolution of social marketing strategies around the promotion of healthcare [2,3]. This has resulted in the development of cancer oriented societies, websites, public campaigns and specifically earmarked Cancer Awareness Months (CAMs) directed at changing public attitudes towards prevention, screening, treatment and informed decision making. However, despite the significant impact of cancer awareness on screening of preventable cancers [3], the impact of CAMs on cancer-related internet search activity has not been

well studied. Breast (BC), Lung (LC) and Colorectal Cancers (CRC) are the leading causes of cancer incidence and mortality among women [4] and have their respective CAMs during October, November and March [5].

Using Google Trends, a public web facility of Google Inc. based on Google Search, we compared the relative frequency of search of terms ‘Breast Cancer’, ‘Lung Cancer’ and ‘Colon Cancer’ between 1st January 2004 and 31st January 2017 (n=158 months). The program assigns a reference value of 100 for the point of maximum popularity from among the search terms, and provides relative monthly scores for all terms, which we termed interest scores (IS). IS were then compared among cancers for the overall period (n=158 months) and specifically during their CAMs (n=13 months). Within each cancer, IS were then compared during the CAMs (n=13 months) as compared to the remaining months (n=145 months). Parametric and non-parametric analyses were carried out (wherever applicable) using ANOVA and Kruskal-Wallis tests respectively. A *P*-value of <.05 was considered significant.

We found that BC had higher IS (mean± S.D) than LC and CRC for the entire study period (38.83±14.46 vs 14.71 ±4.56 and 11.98±2.13 respectively, *P*<0.001*), including a peak IS of 100 in October, 2004. BC also had significantly higher IS during its CAM (October) than the CAMs for LC (November) and CRC (March); 69.92±11.75 vs 15.38±4.54 and 13.53±2.43 respectively, *P*<0.001*. While BC (69.92±11.02 vs

36.04±11.02; $P<0.001^*$) and CRC (13.53±11.84 vs 11.85 ± 2.06; $P=0.036^*$) had higher IS during their CAMs as compared to other months, LC did not (15.38 ±4.53 vs 14.65±4.57; $P=0.3019$) (Table 1).

We concluded that ongoing campaigns for BC awareness are very effective at driving internet search activity, not only at baseline (2.5-3 times) but even more so also during its CAM (4-5 times) as compared to the other two leading malignancies among women (CRC and LC). Despite having a higher mortality

than CRC, the campaign for LC was unable to significantly impact internet search activity during its CAM. Reasons behind the success of the BC awareness campaign in driving internet search activity should be further explored and applied to those for other malignancies such as LC and CRC, which also continue to have high mortality. Therefore, as highlighted by Ling et al [1], the use of infoveillance can serve as a cost-effective solution to evaluate the effects of health campaigns. However, further research is needed to definitively establish Google Trends as a valid and reliable tool for this purpose.

Conflicts of Interest

None declared.

References

1. Ling R, Lee J. Disease Monitoring and Health Campaign Evaluation Using Google Search Activities for HIV and AIDS, Stroke, Colorectal Cancer, and Marijuana Use in Canada: A Retrospective Observational Study. *JMIR Public Health Surveill*. Oct 12, 2016;2(2):e156. [FREE Full text] [doi: [10.2196/publichealth.6504](https://doi.org/10.2196/publichealth.6504)] [Medline: [27733330](https://pubmed.ncbi.nlm.nih.gov/27733330/)]
2. Cole GE, Keller PA, Reynolds J, Schaur M, Krause D. CDC MessageWorks: Designing and Validating a Social Marketing Tool to Craft and Defend Effective Messages. *Soc Mar Q*. Mar 2016;22(1):3-18. [FREE Full text] [doi: [10.1177/1524500415614817](https://doi.org/10.1177/1524500415614817)] [Medline: [26877714](https://pubmed.ncbi.nlm.nih.gov/26877714/)]
3. Wichachai S, Songserm N, Akakul T, Kuasiri C. Effects of Application of Social Marketing Theory and the Health Belief Model in Promoting Cervical Cancer Screening among Targeted Women in Sisaket Province, Thailand. *Asian Pac J Cancer Prev*. 2016;17(7):3505-3510. [FREE Full text] [Medline: [27510000](https://pubmed.ncbi.nlm.nih.gov/27510000/)]
4. Centers for Disease Control. URL: <https://www.cdc.gov/cancer/dcpc/data/women.htm> [accessed 2017-07-09] [WebCite Cache ID [6rqG6BKiu](https://webcache.googleusercontent.com/search?q=cache:https://www.cdc.gov/cancer/dcpc/data/women.htm)]
5. American Cancer Society. URL: <https://www.cancer.org/about-us/who-we-are/cancer-awareness-calendar.html> [accessed 2017-07-10] [WebCite Cache ID [6rqGDu5W5](https://webcache.googleusercontent.com/search?q=cache:https://www.cancer.org/about-us/who-we-are/cancer-awareness-calendar.html)]

Edited by T Sanchez; submitted 19.Mar.2017; peer-reviewed by N Amoyal Pensak, Y-M Schoenberger, SFA Shah; comments to author 23.Jun.2017; revised version received 09.Jul.2017; accepted 10.Jul.2017; published 21.Aug.2017

Please cite as:

Mukhija D, Venkatraman A, Nagpal SJS

Effectivity of Awareness Months in Increasing Internet Search Activity for Top Malignancies Among Women

JMIR Public Health Surveill 2017;3(3):e55

URL: <http://publichealth.jmir.org/2017/3/e55/>

doi: [10.2196/publichealth.7714](https://doi.org/10.2196/publichealth.7714)

PMID: [28827213](https://pubmed.ncbi.nlm.nih.gov/28827213/)

©Dhruvika Mukhija, Anand Venkatraman, Sajan Jiv Singh Nagpal. Originally published in JMIR Public Health and Surveillance (<http://publichealth.jmir.org>), 21.Aug.2017. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.